

단 신

벤젠 다이올과 알파-클로로-알파-(메틸싸이오)아세톤으로부터 치환 벤조다이푸란 유도체의 합성

장지봉 · 서필자 · 손병화[†] · 최홍대^{*}

동의대학교 화학과

[†]부경대학교 화학과

(2006. 3. 20 접수)

Synthesis of Substituted Benzodifuran Derivatives from Benzene Diols and α -Chloro- α -(methylthio)acetone

Ji Bong Jang, Pil Ja Seo, Byeng Wha Son[†], and Hong Dae Choi^{*}

Department of Chemistry, Donggeui University, Busan 614-714, Korea

[†]Department of Chemistry, Pukyong National University, Busan 608-737, Korea

(Received March 20, 2006)

주제어: 치환 벤조다이푸란 유도체, 벤젠 다이올류, 알파-클로로-알파-(메틸싸이오)아세톤

Keywords: Substituted Benzodifuran Derivatives, Benzene Diols, α -Chloro- α -(methylthio)acetone

The benzodifuran ring system has been known to exhibit interesting biological properties,¹ and another important applications of these compounds are in the field of photosensitizers² and fluorescent probes.³


With our ongoing research for the synthetic method of various arene ring-fused furan derivatives,^{4,6} we describe here one-pot synthesis of substituted benzodifurans from benzene diols and α -chloro- α -(methylthio)acetone (**1**).

The chloride **1** was prepared from α -(methylthio)acetone by chlorination with *N*-chlorosuccinimide according to the reported method.⁷ Initially, a methylene chloride-tetrahydrofuran solution of hydroquinone and an excess of the chloride **1** was treated with zinc chloride at room temperature. The crude material was chromatographed to give substituted benzodifurans (**2,3**) in 25% and 33% yields, respectively. The formation of isomers (**2,3**) could be due to the presence of the two hydroxyl groups on the *para* position of hydroquinone ring. Therefore, we concluded that the type **A** and **B** denoted in

Scheme 1 lead to 2,5-dimethyl-3,4-dimethylthiobenzo[1,2-*b*:4,3-*b'*]difuran (**2**) and 2,6-dimethyl-3,7-dimethylthiobenzo[1,2-*b*:4,5-*b'*]difuran (**3**).

Next, we examined the reaction of other benzene diols and methylresorcinols on the base of reaction conditions for the compounds (**2,3**). The results are summarized in *Scheme 1*. The reactions of resorcinol and 2-methylresorcinol took place smoothly at room temperature to give the adducts (**4,6**) in *ca.* 50% yields. In the case of catechol, 2,7-dimethyl-3,6-dimethylthiobenzo[1,2-*b*:6,5-*b'*]difuran (**5**) was obtained in 17% yield. We assumed that the attack of incoming groups toward electrophilic substitution could not easily be accomplished due to the adjacent two hydroxyl groups in catechol.

The reaction of 5-methylresorcinol with an excess of the chloride **1** afforded substituted benzodifurans (**7,8**) in 27% and 24% yields, respectively. We could confirmed that the type **C** and **D** denoted in *Scheme 1* lead to 2,4,6-trimethyl-3,5-dimethylthiobenzo[1,2-*b*:5,4-*b'*]difuran (**7**) and 2,4,7-trimethyl-3,8-dimethylthiobenzo[1,2-*b*:5,6-*b'*]difuran (**8**).



Scheme 1.

Like our previous report,⁴⁻⁵ The mechanism for the one-pot synthesis of benzodifurans **2-8** is similarly explained by a successive dehydrocyclization of Friedel-Crafts intermediate being obtained under Lewis acid conditions. The structures of the above new compounds **2-8** were determined by well-defined ^1H and ^{13}C NMR spectra, and mass spectra (see "Experimental").







As shown in Table 1, the adducts (**2-4**, **6-8**) obtained by the Lewis acid catalyzed reaction could easily be desulfurized into the corresponding benzodifurans (**9-14**) by heating with excess Raney nickel in ethanol.

A few synthetic methods for benzodifuran nucleus have so far been reported in the literature. The fol-

lowing methods are representative: 1) alkylation of 4,6-diacetylresorcinol with ethyl bromoacetate followed by saponification of the resulting ester and decarboxylative cyclization,^{2a} 2) photocyclization of 1,2-di(2-furyl)ethene,⁸ 3) the double photocyclization of 1,5-dibenzoyl-2,4-dialkoxybenzene followed by dehydration,⁹ 4) the base-catalyzed dehydrative cyclization of an appropriate dihydroxydibenzoylbenzene prepared from dimethoxybenzene with benzoyl chloride.¹⁰ Thus, these reports are rather limited on the synthesis of various benzodifuran isomers.

In summary, we developed a new route for synthesizing substituted benzodifuran derivatives using benzene diols with excess α -chloro- α -(methylthio)-

Table 1. The synthesis of desulfurized benzodifuran derivatives 9-14^a

Substrate	Product (%) ^b
	9 (83)
	10 (85)
	11 (82)
	12 (87)
	13 (88)
	14 (85)

^aReaction conditions: excess Raney nickel, EtOH, 60-70°C, 3 h. ^bIsolated yields.

acetone (**1**) in the presence of zinc chloride. The advantages of the present method are as follows: the diverse and available benzene diols are directly utilized as the starting materials, and the sequence of reactions can be performed under rather mild conditions.

EXPERIMENTAL

General. Melting points were measured by an electrothermal digital melting point apparatus IA 9100 and uncorrected. IR spectra were obtained on a JASCO FT IR-300E spectrometer. NMR spectra were recorded on a JNM-ECP 400 spectrometer using tetramethylsilane as an internal standard. Mass spectra were obtained by electron impact (EI) method using a Hewlett Packard 5970 GC-MS system. TLC was run on a Merck precoated silica gel plates. Silica gel 60 (70-230 mesh, E. Merck) was

used for all column chromatographic separations.

α -Chloro- α -(methylthio)acetone (1**):** *N*-Chlorosuccinimide (10.5 g, 0.08 mol) was added to a stirred solution of α -(methylthio)acetone (8.2 g, 0.08 mol) in CCl₄ (40 mL) in small portions at 0 °C and the stirring was continued at room temperature for 3 h. The precipitated succinimide was filtered off and the solvent was removed *in vacuo*. The residual oil was distilled to give **1** (6.8 g, 63%), bp 55-56 °C/7 mmHg (lit.⁷ 76-77 °C/15 mmHg); ¹H NMR (400 MHz, CDCl₃) δ 2.19 (s, 3H), 2.37 (s, 3H), 5.38 (s, 1H).

Typical procedure for the synthesis of substituted benzodifurans (2,3): To a stirred solution of **1** (1.53 g, 11.0 mmol) and hydroquinone (550 mg, 5.0 mmol) in CH₂Cl₂ (30 mL) and THF (5 mL) was added ZnCl₂ (1.50 g, 11.0 mmol) at room temperature, and the stirring was continued at the same temperature for 40 min. The mixture was quenched with water and organic layer was separated. The organic layer was dried over anhydrous MgSO₄ and concentrated *in vacuo*. The residue was purified by column chromatography (CCl₄) to afford **2** (348 mg) and **3** (458 mg) as white solids in 25% and 33%, respectively. The other compounds **4-8** were synthesized similarly and the spectroscopic data of the products **2-8** are as follows.

2,5-Dimethyl-3,4-dimethylthiobenzo[1,2-b:4,3-b']difuran (2**):** *R*_f=0.68 (CCl₄); 25%; white solid, mp 170-171 °C; IR (KBr) 2982, 2914, 1572, 1406, 1308, 1267, 1216, 1143, 1038 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 2.43 (s, 6H), 2.62 (s, 6H), 7.30 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 12.65, 22.05, 106.99, 108.97, 122.28, 150.80, 159.67; EI-MS *m/z* 278 (M⁺, 100%), 263, 247, 230, 217, 199, 188, 139, 115, 106, 89, 75.

2,6-Dimethyl-3,7-dimethylthiobenzo[1,2-b:4,5-b']difuran (3**):** *R*_f=0.60 (CCl₄); 33%; white solid, mp 179-180 °C; IR (KBr) 2917, 1588, 1424, 1351, 1176, 1137, 1052 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 2.31 (s, 6H), 2.57 (s, 6H), 7.55 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 12.61, 18.50, 100.32, 108.63, 127.56, 151.08, 158.78; EI-MS *m/z* 278 (M⁺, 100%), 263, 248, 216, 203, 187, 139, 131, 124, 115, 95, 77.

2,6-Dimethyl-3,5-dimethylthiobenzo[1,2-b:5,4-

b']difuran (4): $R_f=0.65$ (CCl_4); 52%; white solid, mp 94-95 °C; IR (KBr) 2983, 2919, 1595, 1431, 1374, 1274, 1173, 1113, 1067 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.35 (s, 6H), 2.56 (s, H), 7.46 (s, 1H), 7.68 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 12.56, 18.58, 94.20, 107.65, 108.39, 126.52, 151.96, 158.34; EI-MS (m/z) 278 (M^+ , 100%), 263, 248, 216, 203, 187, 159, 131, 124, 115, 98, 69.

2,7-Dimethyl-3,6-dimethylthiobenzo[1,2-b:6,5-b']difuran (5): $R_f=0.68$ (CCl_4); 17%; white solid, mp 96-97 °C; IR (KBr) 2919, 2859, 1644, 1442, 1390, 1354, 1282, 1182, 1124 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.19 (s, 6H), 2.38 (s, 6H), 7.36 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 8.11, 11.93, 92.99, 106.02, 109.37, 126.57, 150.03; EI-MS m/z 278 (M^+ , 100%), 263, 247, 214, 199, 171, 155, 128, 115, 92, 77.

2,6,8-Trimethyl-3,5-dimethylthiobenzo[1,2-b:5,4-b']difuran (6): $R_f=0.67$ (CCl_4); 53%; white solid, mp 130-131 °C; IR (KBr) 2920, 1588, 1429, 1372, 1276, 1198, 1140, 1095, 1043 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.34 (s, 6H), 2.57 (s, 6H), 2.61 (s, 3H), 7.52 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 8.36, 12.60, 18.58, 104.62, 104.78, 108.56, 126.06, 150.76, 158.09; EI-MS m/z 292 (M^+ , 100%), 277, 262, 245, 230, 217, 201, 171, 131, 115, 95, 69.

2,4,6-Trimethyl-3,5-dimethylthiobenzo[1,2-b:5,4-b']difuran (7): $R_f=0.71$ (CCl_4); 27%; white solid, mp 121-122 °C; IR (KBr) 2916, 1592, 1430, 1382, 1215, 1136, 1072 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.30 (s, 6H), 2.55 (s, 6H), 3.37 (s, 3H), 7.28 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 11.44, 12.37, 20.99, 91.62, 108.55, 123.28, 123.58, 151.86, 159.18; EI-MS m/z 292 (M^+ , 100%), 277, 262, 245, 229, 201, 187, 171, 146, 115, 89, 59.

2,4,7-Trimethyl-3,8-dimethylthiobenzo[1,2-b:5,6-b']difuran (8): $R_f=0.66$ (CCl_4); 24%; white solid, mp 102-103 °C; IR (KBr) 2919, 1609, 1496, 1336, 1282, 1236, 1171, 1113, 1007 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.29 (s, 3H), 2.47 (s, 3H), 2.53 (s, 3H), 2.64 (s, 3H), 2.89 (s, 3H), 7.09 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 12.25, 12.41, 18.23, 19.29, 21.04, 106.47, 108.28, 109.35, 113.27, 122.59, 127.10, 145.70, 152.66, 155.89, 158.36; EI-MS m/z 292 (M^+ , 100%), 277, 261, 245, 231, 201, 187, 171,

146, 115, 87, 59.

Typical procedure for the synthesis of desulfurized benzodifuran (9):

Compound **2** (195 mg, 0.7 mmol) was heated under reflux in ethanol (40 mL) containing Raney nickel (W-2, ca. 2.5 g) for 3h. The Raney nickel was removed by filtration and the solvent was evaporated off. The residue was purified by column chromatography (CCl_4) to afford **9** (108 mg) as white solid in 83%. The other compounds **10-14** were prepared similarly from substituted benzodifurans (**3-4**, **6-8**) and the spectroscopic data **9-14** are as follows.

2,5-Dimethylbenzo[1,2-b:4,3-b']difuran (9): $R_f=0.66$ (CCl_4); 83%; white solid, mp 109-110 °C; IR (KBr) 3114, 2913, 1585, 1419, 1269, 1221, 1155, 1043 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.49 (s, 6H), 6.50 (s, 2H), 7.25 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.22, 101.60, 105.74, 120.79, 151.16, 155.40; EI-MS m/z 186 (M^+ , 100%), 171, 157, 128, 115, 92, 77.

2,6-Dimethylbenzo[1,2-b:4,5-b']difuran (10): $R_f=0.61$ (CCl_4); 85%; white solid, mp 117-118 °C; IR (KBr) 2957, 2912, 1612, 1428, 1374, 1283, 1171, 1112 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.45 (s, 6H), 6.38 (s, 2H), 7.40 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.27, 100.46, 102.74, 125.98, 151.74, 155.49; EI-MS m/z 186 (M^+ , 100%), 171, 157, 128, 115, 92, 77.

2,6-Dimethylbenzo[1,2-b:5,4-b']difuran (11): $R_f=0.71$ (CCl_4); 82%; white solid, mp 83-84 °C; IR (KBr) 2964, 2914, 1609, 1435, 1349, 1277, 1125, 1084 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.45 (s, 6H), 6.37 (s, 2H), 7.42 (s, 1H), 7.43 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.18, 93.41, 102.31, 109.04, 125.45, 152.47, 155.21; EI-MS m/z 186 (M^+ , 100%), 171, 157, 143, 128, 115, 92, 77.

2,6,8-Trimethylbenzo[1,2-b:5,4-b']difuran (12): $R_f=0.69$ (CCl_4); 87%; white solid, mp 94-95 °C (lit.^{2a} 91.5-92 °C); IR (KBr) 2917, 1603, 1436, 1372, 1270, 1202, 1144, 1099 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.45 (s, 6H), 2.62 (s, CH_3 , 3H), 6.35 (s, 2H), 7.25 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 8.62, 14.25, 102.61, 103.89, 105.99, 125.01, 151.32, 154.95; EI-MS m/z 200 (M^+ , 100%), 185,

171, 159, 141, 128, 115, 99, 77.

2,4,6-Trimethylbenzo[1,2-b:5,4-b']difuran (13): $R_f=0.64$ (CCl_4); 86%; white solid, mp 88-90°C; IR (KBr) 2916, 2856, 1599, 1436, 1366, 1269, 1211, 1092, 1049 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.45 (s, 6H), 2.54 (s, 3H), 6.40 (s, 2H), 7.28 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.23, 15.42, 90.92, 100.84, 118.97, 124.76, 152.28, 154.47; EI-MS m/z 200 (M^+ , 100%), 185, 171, 167, 128, 115, 100, 77.

2,4,7-Trimethylbenzo[1,2-b:5,6-b']difuran (14): $R_f=0.65$ (CCl_4); 85%; white solid, mp 64-65°C; IR (KBr) 2918, 2736, 1587, 1439, 1395, 1271, 1200, 1067 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.48 (s, 3H), 2.49 (s, 3H), 2.52 (s, 3H), 6.42 (s, 1H), 6.54 (s, 1H), 7.09 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.06, 14.09, 18.88, 98.64, 101.75, 106.32, 111.89, 123.39, 124.60, 144.17, 153.25, 153.56, 154.02; EI-MS m/z 200 (M^+ , 100%), 185, 171, 157, 128, 115, 100, 84, 63.

Acknowledgement. This work was supported by the Dong-eui University grant (2005AA094).

REFERENCES

- (a) Rene, L.; Buisson, J. P.; Royer, R. *Eur. J. Med. Chem.* **1977**, *12*, 31. (b) Bilger, C.; Demerseman, P.; Buisson, J. P.; Royer, R.; Gayral, P.; Fourniat, J. *Eur. J. Med. Chem.* **1987**, *22*, 213. (c) Chambers, J. J.; Kurash-Orbaugh, D. M.; Parker, M. A.; Nichols, D. E. *J. Med. Chem.* **2001**, *44*, 1003.
- (a) Worden, L. R.; Burgstahler, A. W.; Kaufman, K. D.; Weis, J. A.; Schaaf, T. K. *J. Heterocycl. Chem.* **1969**, *6*, 191. (b) Rene, L.; Buisson, J. P.; Royer, R. *Eur. J. Med. Chem.* **1978**, *13*, 485.
- Abdul-Aziz, M.; Auping, J. V.; Meador, M. A. *J. Org. Chem.* **1995**, *60*, 1303.
- Choi, H. D.; Seo, P. J.; Son, B. W. *J. Korean Chem. Soc.*, **1999**, *43*, 237.
- (a) Choi, H. D.; Seo, P. J.; Son, B. W. *J. Korean Chem. Soc.*, **1999**, *43*, 494. (b) Choi, H. D.; Seo, P. J. *J. Korean Chem. Soc.*, **2000**, *44*, 485.
- (a) Choi, H. D.; Ha, M. C.; Seo, P. J.; Son, B. W.; Song, J. C. *Arch. Pharm. Res.*, **2000**, *23*, 438. (b) Choi, H. D.; Seo, P. J.; Son, B. W. *Arch. Pharm. Res.*, **2002**, *25*, 786. (c) Choi, H. D.; Seo, P. J.; Son, B. W.; Kang, B. W. *Arch. Pharm. Res.*, **2003**, *26*, 985. (d) Choi, H. D.; Seo, P. J.; Son, B. W.; Kang, B. W. *Arch. Pharm. Res.*, **2004**, *27*, 19.
- Bohme, H.; Krack, W. *Justus Liebigs Ann. Chem.*, **1977**, 51.
- Kellogg, R. M.; Groen, M. B.; Wynberg, H. *J. Org. Chem.*, **1967**, *32*, 3093.
- Park, K. K.; Lim, H.; Kim, S. H.; Bae, D. H. *J. Chem. Soc., Perkin Trans. I*, **2002**, 310.
- Park, K. K.; Jeong, J. S. *Tetrahedron*, **2005**, *61*, 545.