

licable to various primary and secondary alcohols, (b) the reagent, chlorotrimethylsilane/sodium iodide/zinc, is less expensive and more easily handled than other known reducing agents, (c) the reaction or work-up procedure is simple.


When the reaction of *p*-anisyl alcohol with chlorotrimethylsilane/sodium iodide was carried out in acetonitrile, followed by treatment with an excess of zinc dust¹³ with stirring at 75–85 °C for 6 h, *p*-methylanisole was obtained in 61% yield by distillation after working up. A similar reaction of decyl alcohol with the reagent gave decane in 63% yield, accompanied by a considerable amount of residue in the distillation vessel. However, addition of a little acetic acid to the zinc-acetonitrile mixture in the reduction step results in a dramatic enhancement of the product yield (83% and 86% yield, respectively), owing probably to its effectiveness as a proton donor.

Chlorotrimethylsilane/Sodium Iodide/Zinc as a Simple and Convenient Reducing System; One-Pot Deoxygenation of Alcohols and Ethers

Tsuyoshi MORITA*, Yoshiki OKAMOTO, Hiroshi SAKURAI

The Institute of Scientific and Industrial Research, Osaka University, Suita, Osaka 565, Japan

Deoxygenation of aliphatic alcohols into the corresponding alkanes is an important transformation in organic synthesis. A common route to the alkanes is reduction using various metal hydrides under anhydrous conditions^{1,2}. In such a case, the process necessarily requires the initial conversion of the hydroxy function into suitable derivatives containing good leaving groups such as tosylate, mesylate, halide etc. More recently, several new methods for the one-step reduction of alcohols have been described using lithium/ammonia³, organosilane/boron trifluoride⁴, sodium borohydride/trifluoroacetic acid⁵, aluminium chloride/palladium⁶, diiododimethylsilane⁷, and iron pentacarbonyl⁸. However, these procedures are almost limited to the α -aryl-substituted alcohols which are capable of forming relatively stable carbanions or carbenium ions. During the course of our continuing work utilizing chlorotrimethylsilane/sodium iodide reagent as an iodotrimethylsilane equivalent^{9–12}, we have found a novel and alternative system for one-pot conversion of alcohols and ethers **1** into the alkanes **2** via *in situ* generated iodides. The advantages of our method are as followed: (a) it is app-



The results are summarized in the Table. The products were not essentially contaminated with any alkene. All alkanes were identified by comparison (N.M.R., G.L.C., b.p.) with authentic samples. On comparison with the known reducing system of alcohols via the iodides by the use of methyltriphenoxyphosphonium iodide/sodium cyanoborohydride in hexamethylphosphoric triamide¹⁴, our present procedure is simple and convenient, producing volatile hexamethyldisiloxane as the only by-product.

On the other hand, no convenient and satisfactory route to alkanes from ethers was previously known¹. To demonstrate further the utility of our method, we applied it for one-pot deoxygenation of alkyl or trimethylsilyl ethers **1**. Interestingly, the reaction of decyl methyl ether with chlorotrimethylsilane/sodium iodide and zinc gave decane in 82% yield. Some data for the ether reductions are also given in the Table.

Deoxygenation of Alcohols **1** ($\text{R}^2 = \text{H}$); General Procedure:

To a mixture of the alcohol **1** (20 mmol) and anhydrous sodium iodide (7.5 g, 50 mmol) in dry acetonitrile (20 ml), chlorotrimethylsilane (5.4 g, 40 mmol) is added dropwise with stirring over 20 min. During the addition, the reaction vessel is heated at 30–55 °C. After 0.5–2.5 h, the reaction mixture is diluted with additional acetonitrile (10 ml) and acetic acid

Table. Deoxygenation of Alcohols and Ethers **1** into Alkanes **2**

Substrate 1 R^1	R^2	Reaction 1		Reaction 2		Yield [%] ^a	b.p. [°C]/torr	
		Temp. [°C]	Time [h]	Zn (equiv.)	Time [h]		observed	Lit. ¹⁵
<i>n</i> -C ₁₀ H ₂₁	H	30–35°	1.0	3	6.0	80	67–69°/16	57°/10
<i>n</i> -C ₁₀ H ₂₁	H	30–35°	1.0	7	6.0	86	71–73°/20	57°/10
<i>n</i> -C ₁₂ H ₂₅	H	30–35°	1.0	7	5.0	93	93–100°/14	91°/10
<i>s</i> -C ₉ H ₁₉	H	40–45°	1.5	7	6.0	61	70–72°/54	39°/10
4- <i>t</i> -butylcyclohexyl (<i>cis</i> and <i>trans</i>)	H	50–55°	2.5	7	5.0	40 ^b	59–62°/16	171°
4-H ₃ CO—C ₆ H ₄ —CH ₂	H	30–35°	0.5	7	5.0	83	63–65°/13	176°
(C ₆ H ₅) ₂ CH	H	30–35°	0.5	7	4.0	91	122–123°/10	125°/10
<i>n</i> -C ₁₀ H ₂₁	CH ₃	70–75°	1.5	7	4.5	82	69–71°/17	57°/10
<i>s</i> -C ₉ H ₁₉	CH ₃	70–75°	1.5	7	5.0	64	73–75°/60	39°/10
<i>n</i> -C ₁₀ H ₂₁	Si(CH ₃) ₃	70–75°	1.0	6	4.0	82	70–72°/18	57°/10
4-H ₃ CO—C ₆ H ₄ —CH ₂	Si(CH ₃) ₃	50–55°	0.5	6	5.0	81	68–69°/17	176°
(C ₆ H ₅) ₂ CH	Si(CH ₃) ₃	50–55°	0.3	6	4.0	85	130–132°/17	125°/10

^a Yield of isolated product of >98% purity as determined by ¹H-N.M.R. spectrometry and G.L.C. analysis (conditions: 1 m × 4 mm column, 5% Silicone OV-17 on Chromosorb W).

^b Contains ~5% of 4-*t*-butylcyclohexene.

(2 ml). Next zinc dust (3–7 equiv) is added portionwise with stirring to the resulting dark-yellow reaction mixture at room temperature. A slight evolution of heat immediately occurs and after 15 min the mixture is heated to 75–85 °C, followed by stirring vigorously for an additional 4–6 h. After cooling to room temperature, the mixture is filtered through a Büchner funnel and washed with diethyl ether (5 × 25 ml). The filtrate is washed with dilute aqueous sodium hydrogen carbonate solution (3 × 30 ml) and subsequently with aqueous sodium hydrogen sulfite solution (2 × 25 ml), then dried with anhydrous sodium sulfate. After the removal of low-boiling materials on an evaporator, the residue is distilled in vacuo to afford the pure alkane 2.

Deoxygenation of Ethers 1 ($R^2 = CH_3$ or $Si(CH_3)_3$); General Procedure:

A mixture of the methyl ether 1 (20 mmol), chlorotrimethylsilane (5.4 g, 50 mmol), and anhydrous sodium iodide (7.5 g, 50 mmol) in dry acetonitrile (20 ml) is heated at 50–75 °C with stirring. In the case of the trimethylsilyl ether 1 (20 mmol), chlorotrimethylsilane (3.3 g, 30 mmol) and sodium iodide (6.0 g, 40 mmol) are used in acetonitrile (20 ml). After the reaction is complete, the mixture is diluted with additional acetonitrile (10 ml) and acetic acid (4 ml). Then it is reduced with zinc and worked up in a similar manner to that described for the alcohol deoxygenation. After work-up, the pure alkane 2 is obtained by fractional distillation (column: 10 cm Vigreux).

The authors wish to thank Shin-Etsu Chemical Industry Ltd. for a gift of chlorotrimethylsilane.

Received: June 5, 1980

(Revised form: August 4, 1980)

- ¹ For recent reviews, see: H. O. House, *Modern Synthetic Reactions*, W. A. Benjamin, New York, 1972, Chapter 2.
- I. T. Harrison, S. Harrison, *Compendium of Organic Synthetic Methods*, Wiley-Interscience, New York, 1971; 1974, Vol. 2; 1977, Vol. 3.
- ² R. O. Hutchins et al., *J. Org. Chem.* **43**, 2259 (1978), and references therein.
- ³ G. H. Small, A. E. Minnella, S. S. Hall, *J. Org. Chem.* **40**, 3151 (1975).
- ⁴ M. G. Adlington, M. Orfanopoulos, J. L. Fry, *Tetrahedron Lett.* **1976**, 2955.
- ⁵ G. W. Gribble, R. M. Leese, B. E. Evans, *Synthesis* **1977**, 172.
- ⁶ G. A. Olah, G. K. S. Prakash, *Synthesis* **1978**, 397.
- ⁷ W. Ando, M. Ikeno, *Tetrahedron Lett.* **1979**, 4941.
- ⁸ H. Alper, M. Salisova, *Tetrahedron Lett.* **1980**, 801.
- ⁹ T. Morita, Y. Okamoto, H. Sakurai, *Tetrahedron Lett.* **1978**, 2523.
- ¹⁰ T. Morita, Y. Okamoto, H. Sakurai, *J. Chem. Soc. Chem. Commun.* **1978**, 874.
- ¹¹ T. Morita, S. Yoshida, Y. Okamoto, H. Sakurai, *Synthesis* **1979**, 379.
- ¹² T. Morita, Y. Okamoto, H. Sakurai, *Chem. Lett.* **1980**, 436.
See also: G. A. Olah, S. C. Narang, B. G. B. Gupta, R. Malhotra, *J. Org. Chem.* **44**, 1247 (1979).
- ³ Zinc (85% purity) was activated by a brief treatment with dilute aqueous HCl solution as described in the literature: K. Tsuda, E. Ohki, S. Nozoe, *J. Org. Chem.* **28**, 783 (1963).
- ⁴ R. O. Hutchins, B. E. Maryanoff, C. A. Milewski, *J. Chem. Soc. Chem. Commun.* **1971**, 1097.
- ⁵ *Handbook of Chemistry and Physics*, 54th Edn., Chemical Rubber Co., Cleveland, Ohio, 1973–1974.