

S0040-4039(96)00496-0

The Isolation from *Nicandra physalodes* and Identification of the 3-O- β -D-glucopyranoside of 1 α ,2 β ,3 α ,6 α -tetrahydroxy-*nor*-tropane (Calystegine B₁).

Rhodri C Griffiths^a, Alison A Watson^b, Haruhisa Kizu^c, Naoki Asano^c, Hazel J Sharp^b, M. George Jones^a, Mark R. Wormald^d, George W. J. Fleet^e and Robert J Nash^{b*}

^aInstitute of Biological Sciences, University of Wales, Aberystwyth, Dyfed SY23 1DB, UK ^bInstitute of Grassland and Environmental Research, Plas Gogerddan, Aberystwyth, Dyfed SY23 3EB, UK ^cFaculty of Pharmaceutical Sciences, Hokuriku University, Kanagawa-machi, Kanazawa 920-11, Japan. ^dGlycobiology Institute, Biochemistry Department, South Parks Road, Oxford OX1 3QU, UK ^eDyson Perrins Laboratory, South Parks Road, Oxford, OX1 3QY, UK

Abstract: The isolation and identification of 3-O- β -D-glucopyranosyl-1 α ,2 β ,3 α ,6 α -tetrahydroxy-*nor*-tropane from *Nicandra physalodes* Boehm. fruits (Solanaceae) is reported.


Copyright © 1996 Elsevier Science Ltd

Polyhydroxylated mono- and bicyclic nitrogen heterocycles are an important class of glycosidase inhibitors¹. Polyhydroxy-*nor*-tropane alkaloids are the most recent naturally-occurring class of these inhibitors to be discovered and they have been shown to be potent inhibitors of β -glucosidases and β -galactosidases². These alkaloids were first found in bindweeds³ (Convolvulaceae) and given the trivial name calystegines but have since been found in human foods such as potato tubers (*Solanum tuberosum*) and aubergine fruits (*Solanum melongena*)⁴. Calystegines are clearly widespread and their significance in the human diet remains to be explored. We now report the first isolation and identification of a glucoside of a calystegine.

Nicandra physalodes Boehm. (Solanaceae) fruits (230g fresh weight) were homogenised in 70% aqueous ethanol. The filtrate was applied to the cation exchange resin Dowex 50W-X2 (H⁺ form) and the bound compounds displaced with 2M ammonia solution. The tropane alkaloid calystegine B₁ (1 α ,2 β ,3 α ,6 α -tetrahydroxy-*nor*-tropane) and a glycoside of it were determined to be the major alkaloids present by GC-MS of the trimethylsilyl-derivatives. The alkaloids were readily separated from amino acids in the extracted material by ion exchange chromatography using Amberlite CG120 (NH₄⁺ form) with the glycoside displaced with 2M pyridine and the aglycone eluted before arginine with 0.1M ammonia solution. The glycoside was then purified on the anion exchange resin Dowex 1-X2 (OH form) and washed off with water (yield 2.1mg).

The structure of the glycoside was determined to be 3-O- β -D-glucopyranosylcalystegine B₁ **1** on the basis of ¹H and ¹³C NMR data, including 2D HMQC and HMBC spectral data⁵. The complete carbon and hydrogen atom connectivity of both the aglycone and glycone was defined. From comparison with previously reported NMR data^{2,6}, the aglycone was identified as calystegine B₁. The large vicinal *J* values of the glycone H-2', H-3', and H-4' and coupling constant of the anomeric proton (H-1', δ 4.50, *J*_{1,2'} = 7.8 Hz) indicate that the glycone part of this glycoside is the pyranose form of β -glucose. It was shown that D-glucose is contained in the filtrate after acid hydrolysis of this glycoside using Dowex 50W-X2 (H⁺) resin by the D-glucose-oxidase peroxidase method. The aglycone part was eluted with 0.5M ammonia solution from the resin, concentrated to dryness, and confirmed as calystegine B₁ by GC-MS of the trimethylsilylated eluate. The HMBC spectrum showed a correlation peak between the anomeric proton}

of the glucone and the aglycone C-3 carbon, defining the linkage site. The ^{13}C -NMR data for the calystegine component shows a 7.6 ppm downfield shift for C-3 and 2.0 and 2.7 ppm upfield shifts for C-2 and C-4 respectively, compared to the free calystegine, also consistent with a 3-O- linkage.



3-O- β -D-glucopyranosyl-(calystegine B₁) 1

3-O- β -D-Glucopyranosylcalystegine B₁: ^1H -NMR (400 MHz, D₂O); δ : 1.40 (m, 1H, H-7_{exo}); 1.49 (ddd, 1H, $J_{3,4\text{ax}}=10.7$, $J_{4\text{ax},4\text{eq}}=13.4$, $J_{4\text{ax},5}=3.9\text{Hz}$, H-4_{ax}); 2.19 (ddd, 1H, $J_{3,4\text{eq}}=6.4$, $J_{4\text{ax},4\text{eq}}=13.4$, $J_{4\text{eq},5}=2.7\text{ Hz}$, H-4_{eq}); 2.53 (dd, 1H, $J_{6,7\text{endo}}=7.3$, $J_{7\text{endo},7\text{exo}}=14.4\text{ Hz}$, H-7_{endo}); 3.26 (dd, 1H, $J_{1,2}=7.8$, $J_{2,3}=9.5\text{ Hz}$, H-2'); 3.29 (m, 1H, H-5); 3.37 (dd, 1H, $J_{3',4}=9.0$, $J_{4',5}=9.8\text{ Hz}$, H-4'); 3.44 (ddd, 1H, $J_{4',5}=9.8$, $J_{5',6\text{a}}=6.1$, $J_{5',6\text{b}}=2.2\text{ Hz}$, H-5'); 3.45 (dd, 1H, $J_{2,3}=8.5$, $J_{2,7\text{exo}}=1.7\text{ Hz}$, H-2); 3.47 (t, 1H, $J_{2,3}=J_{3,4}=9.0\text{ Hz}$, H-3'); 3.63 (ddd, 1H, $J_{2,3}=8.5$, $J_{3,4\text{ax}}=10.7$, $J_{3,4\text{eq}}=6.4\text{ Hz}$, H-3); 3.70 (dd, 1H, $J_{5',6\text{a}}=6.1$, $J_{6\text{a},6\text{b}}=12.2\text{ Hz}$, H-6'a); 3.92 (dd, 1H, $J_{5',6\text{b}}=2.2$, $J_{6\text{a},6\text{b}}=12.2\text{ Hz}$, H-6'b); 4.09 (dd, 1H, $J_{6,7\text{endo}}=7.3$, $J_{6,7\text{exo}}=2.7\text{ Hz}$, H-6); 4.50 (d, 1H, $J_{1,2}=7.8\text{ Hz}$, H-1'); ^{13}C -NMR (100 MHz, D₂O); δ : 36.2 (C-4); 43.5 (C-7); 62.7 (C-5); 63.6 (C-6'); 72.5 (C-4'); 75.7 (C-6); 75.8 (C-2'); 78.4 (C-3'); 78.7 (C-5'); 79.3 (C-2); 80.3 (C-3); 93.7 (C-1), 102.9 (C-1'). HRFAB-MS⁷ m/z 338.1446 [M + H] (C₁₃H₂₄O₉N requires 338.1451) measured on a Jeol JMS-SX 102A spectrometer with glycerol matrix⁸.

REFERENCES.

1. Fleet, G., Winchester, B. *Glycobiology* 1992, 2, 199-210.
2. Asano, N., Kato, A., Oseki, K., Kizu, H., Matsui, K. *Eur. J. Biochem.* 1995, 229, 369-376.
3. Tepfer, D., Goldmann, A., Pamboukdjian, N., Maille, M., Lepingle, A., Chevalier, D., Denarie, J. Rosenberg, C. *J. Bacteriol.* 1988, 170, 1153-1161.
4. Nash, R., Rothschild, M., Porter, E., Watson, A., Waigh, R., Waterman, P. *Phytochemistry* 1993, 34, 1281-1283.
5. NMR abbreviations: HMQC, heteronuclear multiple quantum correlation spectroscopy; HMBC, heteronuclear multiple bond correlation spectroscopy.
6. Goldmann, A., Milat, M.L., Ducrot, P.H., Lallemand, J.Y., Maille, M., Lepingle, A., Charpin, I. Tepfer, D. *Phytochemistry* 1990, 29, 2125-2127.
7. MS abbreviations: HR, high resolution; FAB, fast atom bombardment.
8. This work was partly funded by a BBSRC grant to RCG.

(Received in UK 4 March 1996; accepted 14 March 1996)