

The catalytic N-alkylation procedure can also be applied to ring synthesis. Thus the amines (**3a**) and (**3b**) are cyclised to the pyrrolidines (**4a**) (56% isolated yield) and (**4b**) (82% by g.l.c.), respectively, by 5 mol % RhH(PPh₃)₄ in boiling dioxan. In a similar manner butane-1,4-diol and

benzylamine (10:1 mol. ratio) in boiling dioxan give (**4b**) (31% by g.l.c.).

We thank Queen's University for support.

(Received, 30th March 1981; Com. 360.)

¹ C. F. Winan and H. Adkins, *J. Am. Chem. Soc.*, 1932, **54**, 306.

² D. C. Best, U.S. P. 4,123,462.

³ M. V. Klynev and M. L. Khidekel, *Russ. Chem. Rev.*, 1980, **49**, 28; S. Baron, U.S. P. 4,076,649.

⁴ Y. Motoyama, H. Naibu, and H. Nirose, *Jpn. Kokai Tokkyo Koho*, 78/90,227 (*Chem. Abstr.*, 1979, **90**, 22552s).

⁵ E. Csaszar, L. Horvath, O. Hesser, L. Kisgergely, J. Jesztl, G. Kincses, C. Nagy, B. Szeiler, J. Schneider, and E. Szentgyorgyvari, Hungarian P. 14,908 (*Chem. Abstr.*, 1978, **89**, 179532n).

⁶ S. V. Dobrovol'skii, A. F. Grigorov, L. I. Blyakhman, R. M. Grizik, N. I. Sidorova, I. G. Shipilova, N. P. Buldakova, I. G. Kronich, and V. F. Kashina, U.S.S.R. P. 644,526.

⁷ C. E. Bibby, R. Grigg, and R. Price, *J. Chem. Soc., Dalton Trans.*, 1977, 872; R. Grigg, T. R. B. Mitchell, and S. Sutthivaiyakit, *Tetrahedron Lett.*, 1979, 1067.

⁸ J. Hine, R. C. Dempsey, R. A. Evangelista, E. T. Jarvi, and J. M. Wilson, *J. Org. Chem.*, 1977, **42**, 1593.

⁹ R. Grigg, T. R. B. Mitchell, and N. Tongpenyai, *Synthesis*, 1981, in the press.