



US 20060258861A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2006/0258861 A1****Anderskewitz et al.**(43) **Pub. Date: Nov. 16, 2006**(54) **COMPOUNDS FOR THE TREATMENT OF INFLAMMATORY DISEASES****Publication Classification**

(75) Inventors: **Ralf Anderskewitz**, Laupheim (DE);
Horst Dollinger, Schemmerhofen (DE);
Rolf Goeggel, Biberach (DE); **Birgit Jung**, Laupheim (DE); **Joerg Kley**,
Mittelbiberach (DE); **Juergen Mack**,
Biberach-Mettenberg (DE); **Peter Nickolaus**, Warthausen (DE); **Rainer Walter**, Biberach (DE)

(51) **Int. Cl.**
A61K 31/5377 (2006.01)
A61K 31/454 (2006.01)
A61K 31/4184 (2006.01)
C07D 403/02 (2006.01)
C07D 417/02 (2006.01)
C07D 413/02 (2006.01)

(52) **U.S. Cl.** **544/144**; 546/201; 544/373;
548/305.7

Correspondence Address:
MICHAEL P. MORRIS
BOEHRINGER INGELHEIM CORPORATION
900 RIDGEBURY ROAD
P. O. BOX 368
RIDGEFIELD, CT 06877-0368 (US)


(73) Assignee: **Boehringer Ingelheim International GmbH**, Ingelheim (DE)(21) Appl. No.: **11/409,482**(22) Filed: **Apr. 21, 2006**(30) **Foreign Application Priority Data**

Apr. 28, 2005 (DE)..... 10 2005 019 718
May 20, 2005 (DE)..... 10 2005 023 207
Jul. 29, 2005 (DE)..... 10 2005 035 575

(57) **ABSTRACT**

Compounds of formula 1 and hetero derivatives thereof

1



and the pharmacologically acceptable salts, enantiomers, racemates, hydrates, or solvates thereof, which are suitable for the treatment of respiratory or gastrointestinal complaints or diseases, inflammatory diseases of the joints, skin, or eyes, diseases of the peripheral or central nervous system or cancers, as well as pharmaceutical compositions which contain these compounds.


COMPOUNDS FOR THE TREATMENT OF INFLAMMATORY DISEASES

RELATED APPLICATIONS

[0001] This application claims priority to German Application No. DE 10 2005 019 718.3, filed Apr. 28, 2005, German Application No. DE 10 2005 023 207.8, filed May 20, 2005, and German Application No. DE 10 2005 035 575.7, filed Jul. 29, 2005, each of which are hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION


[0002] The invention relates to new compounds of formula 1 and hetero derivatives thereof, as well as pharmacologically acceptable salts, diastereomers, enantiomers, racemates, hydrates, or solvates thereof



which are suitable for the treatment of respiratory or gastrointestinal complaints or diseases, inflammatory diseases of the joints, skin, or eyes, diseases of the peripheral or central nervous system or cancers, as well as pharmaceutical compositions which contain these compounds.

BRIEF DESCRIPTION OF THE INVENTION

[0003] Surprisingly it has now been found that the compounds of formula 1 are suitable for the treatment of inflammatory diseases. The present invention therefore relates to compounds of formula 1



wherein:

[0004] A denotes CO, C=NH, C₁₋₆-alkylene, or C₃₋₈-cycloalkylene;

[0005] B¹ denotes phenyl or an aromatic or non-aromatic ring which may optionally contain one, two, or three

heteroatoms, selected from among oxygen, sulfur, and nitrogen and which may optionally be mono- or polysubstituted by one or more groups selected from among OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, C₁₋₆-alkyl, and C₁₋₆-haloalkyl;

[0006] B² denotes phenyl or a heteroaryl, which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen;

[0007] X denotes O, S, NR⁵, or CR⁶R⁷;

[0008] n denotes 0, 1, 2, or 3;


[0009] R¹ denotes H, C₁₋₆-alkyl, C₁₋₆-haloalkyl, COR^{1.1}, COOR^{1.1}, or CH₂COOR^{1.1}; preferably H, C₁₋₆-alkyl, or C₁₋₆-haloalkyl, wherein R^{1.1} denotes H or C₁₋₆-alkyl;

[0010] R² denotes H, C₁₋₆-alkyl, or C₁₋₆-haloalkyl; or

[0011] R¹ and R² together with the nitrogen form a non-aromatic heterocycle, which may contain one, two, or three heteroatoms selected from among oxygen and nitrogen; or

[0012] R², N, A, and B¹ together form a bicyclic group of formula (i)

wherein:



(i)

[0013] A denotes CO, C=NH, or C₁₋₃-alkyl,

[0014] m denotes 1, 2, or 3, and

[0015] R³ denotes H or a group selected from among OH, C₁₋₆-haloalkyl, a C₆₋₁₀-aryl, a C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, while the C₃₋₁₀-heterocycle and the C₅₋₁₀-heteroaryl may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen, which may optionally be substituted by one or more groups selected from among C₁₋₆-alkyl, C₆₋₁₀-aryl, optionally bridged C₃₋₈-cycloalkyl, and C₁₋₆-haloalkyl, which may optionally be substituted by a group selected from C₁₋₆-alkyl, C₁₋₆-haloalkyl, OH, halogen, and C₆₋₁₀-aryl; or


[0016] R³ denotes C₁₋₆-alkyl, which may optionally be substituted by one or more groups selected from among halogen, OH, CN, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, CO—C₆₋₁₀-aryl, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, SH, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, SO₂—C₁₋₆-alkanol; SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂, or

[0017] R³ denotes a group selected from among C₃₋₈-cycloalkyl, C₅₋₈-cycloalkenyl, C₁₋₆-alkyl, and C₁₋₆-alkanol, which may optionally be substituted by one or more groups selected from among C₆₋₁₀-aryl, C₃₋₈-cycloalkyl, a C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, which may optionally be substituted by one or more

groups selected from among C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, CN, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, COC₆₋₁₀-aryl, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, SH, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, SO₂—C₁₋₆-alkanol, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, N(C₁₋₆-alkyl)₂, a C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, while the C₅₋₁₀-heteroaryl and the C₃₋₁₀-heterocycle may optionally be substituted by a group selected from oxo, hydroxyl, halogen, C₁₋₆-alkyl, and C₁₋₆-haloalkyl; or

[0018] R³ denotes a group selected from among C₆₋₁₀-aryl, a C₆₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, which may optionally be substituted by one or more groups selected from among C₆₋₁₀-aryl, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, CO—C₆₋₁₀-aryl, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, SH, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-alkanol, SO₂—C₁₋₆-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, N(C₁₋₆-alkyl)₂, and N—(SO₂—C₁₋₄-alkyl)(R^{3,4}); or

[0019] R³ denotes a group selected from among C₁₋₆-alkyl, C₆₋₁₀-aryl-C₁₋₆-alkylene, C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, C₃₋₇-cycloalkyl, C₆₋₁₀-aryl, a C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, which may optionally be substituted by one or more groups selected from among B, halogen, OH, C₁₋₆-alkyl, and oxo, while B is a compound of formula 2



[0020] wherein Z¹ denotes H, OH, halogen, C₁₋₆-alkyl, C₁₋₆-alkanol, O(C₁₋₆-alkyl), C₆₋₁₀-aryl, O—C₆₋₁₀-aryl, NH₂, NH(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂, or C₃₋₇-cycloalkyl; and Z² denotes OH, NH₂, NH(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂, O(C₁₋₆-alkyl), mono- or bicyclic C₃₋₇-cycloalkyl, mono- or bicyclic C₃₋₁₀-heterocycle, mono- or bicyclic C₅₋₁₀-heteroaryl, or C₆₋₁₀-aryl; or

[0021] R³ denotes a group selected from among C₆₋₁₀-aryl and a C₅₋₁₀-heteroaryl, which may optionally be substituted by C₁₋₆-alkyl, which may optionally be substituted by a group selected from among COOR^{3,3}, NR^{3,3}R^{3,4}, NHCOR^{3,3}, NHCOOR^{3,3}, phenyl, while phenyl may optionally be substituted by a group selected from among C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, CN, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, SH, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-

haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂, a C₃₋₁₀-heterocycle, and a C₅₋₁₀-heteroaryl, while the C₃₋₁₀-heterocycle and the C₅₋₁₀-heteroaryl may optionally be substituted by an oxo group or a methyl group, wherein:

[0022] R^{3,3} denotes H or C₁₋₆-alkyl, and

[0023] R^{3,4} denotes H, C₁₋₆-alkyl, C₆₋₁₀-aryl-C₁₋₆-alkylene, or C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, or

[0024] R³ denotes a group selected from among C₆₋₁₀-aryl and a C₅₋₁₀-heteroaryl, which may optionally be substituted by NR^{3,1}R^{3,2}, wherein:

[0025] R^{3,1} denotes H, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₆-haloalkyl, COR^{3,1,1}, COOR^{3,1,1}, CONR^{3,1,1}R^{3,1,2}, or SO₂—R^{3,1,1}, wherein: R^{3,1,1} denotes H, C₁₋₆-alkyl, C₁₋₆-haloalkyl, C₃₋₆-cycloalkyl, or C₆₋₁₀-aryl; and R^{3,1,2} denotes H, C₁₋₆-alkyl, C₁₋₆-haloalkyl, C₃₋₆-cycloalkyl, or C₆₋₁₀-aryl; and

[0026] R^{3,2} denotes H or a group selected from among C₁₋₆-alkyl, C₃₋₇-cycloalkyl, and C₁₋₆-haloalkyl, which may optionally be substituted by one or more groups selected from among NH₂, NH(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂, oxo, and a non-aromatic C₃₋₁₀-heterocycle, which may contain one or two heteroatoms selected from among nitrogen, oxygen, and sulfur, while the non-aromatic C₃₋₁₀-heterocycle may optionally be substituted by C₁₋₄-alkyl; or

[0027] R³ denotes a group selected from among C₆₋₁₀-aryl and a C₅₋₁₀-heteroaryl, which may optionally be substituted by a C₃₋₁₀-heterocycle, which may optionally be substituted by one or more groups selected from among C₆₋₁₀-aryl, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, CN, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, oxo, OH, O—C₁₋₆-alkyl, halogen, SH, S—C₁₋₆-alkyl, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂; or

[0028] R³ denotes benzimidazolyl, which may optionally be substituted by a group selected from among C₁₋₆-alkyl, C₁₋₆-haloalkyl, and C₃₋₆-cycloalkyl;

[0029] R⁴ denotes C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-haloalkyl, OR^{4,1}, NR^{4,1}R^{4,2}, CN, or halogen, wherein: R^{4,1} denotes H, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, or C₁₋₆-haloalkyl; and R^{4,2} denotes H, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, or C₁₋₆-haloalkyl;

[0030] R⁵ denotes C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-haloalkyl, COR^{5,1}, CONHR^{5,1}, CON(R^{5,1})₂, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-haloalkyl, SO₂—C₆₋₁₀-aryl, or a group selected from among R^{5,2}, SO₂—C₁₋₆-alkyl-R^{5,2}, and C₁₋₆-alkyl-R^{5,2}, which may optionally be substituted by one or more groups selected from among C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, CN, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, O—C₃₋₆-cycloalkyl, O—C₁₋₆-haloalkyl, O—C₁₋₆-alkyl, halogen, SO₂—C₁₋₆-alkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂, wherein: R^{5,1} denotes C₁₋₆-alkyl, C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, or C₆₋₁₀-aryl-C₁₋₆-alkylene, and R^{5,2} denotes C₆₋₁₀-aryl or a C₅₋₁₀-heteroaryl; and

[0031] R^6 denotes H, C_{1-6} -alkyl, or C_{1-6} -haloalkyl; and

[0032] R^7 denotes H, C_{1-6} -alkyl, or C_{1-6} -haloalkyl; or

[0033] R^6 and R^7 together form a 3-6 membered carbocycle,

optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0034] Preferred compounds of formula 1 above are those wherein:

[0035] A denotes CO, C=NH, C_{1-4} -alkylene, or C_{3-6} -cycloalkylene;

[0036] B^1 denotes phenyl or a C_{5-10} -heteroaryl, which may contain one, two, or three heteroatoms, selected from among oxygen, sulfur, and nitrogen;

[0037] B^2 denotes phenyl or pyridinyl;

[0038] X denotes O, S, NR^5 , or CR^6R^7 ;

[0039] n denotes 0, 1, 2, or 3;

[0040] R^1 denotes H, C_{1-4} -alkyl, C_{1-4} -haloalkyl;

[0041] R^2 denotes H, C_{1-4} -alkyl, or C_{1-4} -haloalkyl; or

[0042] R^1 and R^2 together with the nitrogen form a non-aromatic heterocycle, which may contain one or two nitrogen atoms;

[0043] R^3 denotes H, OH, C_{1-4} -haloalkyl, C_{6-10} -aryl, a C_{5-10} -heteroaryl, which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen, while the C_{6-10} -aryl and the C_{5-10} -heteroaryl may optionally be substituted by one or more groups selected from among C_{1-4} -alkyl, C_{3-6} -cycloalkyl, and C_{1-4} -haloalkyl; or

[0044] R^3 denotes a group selected from among C_{3-8} -cycloalkyl, C_{5-8} -cycloalkenyl, and C_{1-6} -alkyl, which may optionally be substituted by one or more groups selected from among C_{6-10} -aryl or a C_{5-10} -heteroaryl, which may optionally be substituted by one or more groups selected from among C_{1-6} -alkyl, C_{1-6} -haloalkyl, CN, OH, O— C_{1-6} -alkyl, O— C_{1-6} -haloalkyl, halogen, S— C_{1-6} -alkyl, S— C_{1-6} -haloalkyl, NO_2 , NH_2 , NH— C_{1-6} -alkyl, and N(C_{1-6} -alkyl)₂;

[0045] a group selected from among C_{6-10} -aryl and a heterocyclic, aromatic ring, substituted by one or more groups selected from among C_{6-10} -aryl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, C_{1-4} -haloalkyl, $CONH_2$, $CONH-C_{1-4}$ -alkyl, $CON(C_{1-4}$ -alkyl)₂, COOH, COO— C_{1-4} -alkyl, COH, CO— C_{1-4} -alkyl, CO— C_{6-10} -aryl, OH, O— C_{1-4} -alkyl, O— C_{1-4} -haloalkyl, halogen, SH, S— C_{1-4} -alkyl, S— C_{1-4} -haloalkyl, SO_2-C_{1-4} -alkyl, SO_2-C_{1-4} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-4} -alkyl, $SO_2-N(C_{1-4}$ -alkyl)₂, NO_2 , NH_2 , NH— C_{1-4} -alkyl, and N(C_{1-4} -alkyl)₂;

[0046] a group selected from among C_{6-10} -aryl, a C_{5-10} -heteroaryl, C_{6-10} -aryl- C_{1-4} -alkylene, and C_{5-10} -heteroaryl- C_{1-4} -alkylene, which may optionally be substituted by one or more groups selected from among $COOR^{3.3}$, $NR^{3.3}R^{3.4}$, $NHCOR^{3.3}$, $NHCOOR^{3.3}$, and phenyl, which may optionally be substituted by one or

more groups selected from among C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, CN, C_{1-4} -haloalkyl, $CONH_2$, $CONH-C_{1-4}$ -alkyl, $CON(C_{1-4}$ -alkyl)₂, COOH, COO— C_{1-4} -alkyl, COH, CO— C_{1-4} -alkyl, OH, O— C_{1-4} -alkyl, O— C_{1-4} -haloalkyl, halogen, SH, S— C_{1-4} -alkyl, S— C_{1-4} -haloalkyl, SO_2-C_{1-4} -alkyl, SO_2-C_{1-4} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-4} -alkyl, $SO_2-N(C_{1-4}$ -alkyl)₂, NO_2 , NH_2 , NH— C_{1-4} -alkyl, and N(C_{1-4} -alkyl)₂, and a C_{5-10} -heterocycle, which may contain one, two, or three heteroatoms selected from among oxygen, nitrogen, and sulfur and which may optionally be substituted by an oxo group, wherein: $R^{3.3}$ denotes H or C_{1-4} -alkyl, and $R^{3.4}$ denotes H, C_{1-4} -alkyl, C_{6-10} -aryl- C_{1-6} -alkylene, or C_{5-10} -heteroaryl- C_{1-6} -alkylene; or

[0047] R^3 denotes a group selected from among C_{6-10} -aryl and a C_{5-10} -heterocycle, which may be substituted by $NR^{3.1}R^{3.2}$, wherein:

[0048] $R^{3.1}$ denotes H, C_{1-4} -alkyl, C_{3-6} -cycloalkyl, C_{1-4} -haloalkyl, $COR^{3.1.1}$, $COOR^{3.1.1}$, $CONR^{3.1.1}R^{3.1.2}$, or $SO_2-R^{3.1.1}$, wherein: $R^{3.1.1}$ denotes H, C_{1-4} -alkyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, or C_{6-10} -aryl; and $R^{3.1.2}$ denotes H, C_{1-4} -alkyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, or C_{6-10} -aryl; and

[0049] $R^{3.2}$ denotes H or a group selected from among C_{1-4} -alkyl, C_{3-6} -cycloalkyl, and C_{1-4} -haloalkyl, which may optionally be substituted by one or more groups selected from among NH_2 , $NH(C_{1-4}$ -alkyl), N(C_{1-4} -alkyl)₂, oxo, and a C_{3-10} -heterocycle, which may contain one or two heteroatoms selected from among nitrogen, oxygen, and sulfur and which may optionally be substituted by C_{1-4} -alkyl; or

[0050] R^3 denotes C_{6-10} -aryl, which may be substituted by a C_{5-10} -heteroaryl which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen and which may optionally be substituted by one or more groups selected from among C_{6-10} -aryl, C_{1-4} -alkyl, C_{3-6} -cycloalkyl, CN, C_{1-4} -haloalkyl, $CONH_2$, $CONH-C_{1-4}$ -alkyl, $CON(C_{1-4}$ -alkyl)₂, COOH, COO— C_{1-4} -alkyl, COH, CO— C_{1-4} -alkyl, OH, O— C_{1-4} -alkyl, halogen, SH, S— C_{1-4} -alkyl, NH_2 , NH— C_{1-4} -alkyl, and N(C_{1-4} -alkyl)₂; or

[0051] R^3 denotes C_{6-10} -aryl, which may be substituted by a C_{3-10} -heterocycle which may contain one or two heteroatoms, selected from among oxygen, sulfur, and nitrogen and which may optionally be substituted by one or more groups selected from among C_{1-4} -alkyl and oxo; or

[0052] R^3 denotes benzimidazolyl, which may optionally be substituted by a group selected from among C_{1-4} -alkyl, C_{1-4} -haloalkyl, and C_{3-6} -cycloalkyl;

[0053] R^4 denotes C_{1-4} -alkyl, C_{3-6} -cycloalkyl, C_{1-4} -haloalkyl, $OR^{4.1}$, $NR^{4.1}R^{4.2}$, CN, or halogen; and

[0054] $R^{4.1}$ denotes H, C_{1-4} -alkyl, C_{3-6} -cycloalkyl, or C_{1-4} -haloalkyl, and

[0055] $R^{4.2}$ denotes H, C_{1-4} -alkyl, C_{3-6} -cycloalkyl, or C_{1-4} -haloalkyl;

[0056] R^5 denotes C_{1-4} -alkyl, C_{3-8} -cycloalkyl, C_{1-4} -haloalkyl, $COR^{5.1}$, $CONHR^{5.1}$, $CON(R^{5.1})_2$, SO_2-C_{1-4} -alkyl, SO_2-C_{1-4} -haloalkyl, SO_2-C_{6-10} -aryl, or a group

selected from among $R^{5.2}$, $SO_2-C_{1.4}$ -alkyl- $R^{5.2}$, and $C_{1.4}$ -alkyl- $R^{5.2}$, while this group may optionally be substituted by one or more groups selected from among $C_{1.4}$ -alkyl, $C_{2.4}$ -alkenyl, $C_{2.4}$ -alkynyl, CN, $C_{1.4}$ -haloalkyl, $CONH_2$, $CONH-C_{1.4}$ -alkyl, $CON(C_{1.4}$ -alkyl) $_2$, $COOH$, $COO-C_{1.4}$ -alkyl, COH , $CO-C_{1.4}$ -alkyl, $O-C_{1.4}$ -cycloalkyl, $O-C_{1.4}$ -haloalkyl, $O-C_{1.4}$ -alkyl, halogen, $SO_2-C_{1.4}$ -alkyl, SO_2-NH_2 , $SO_2-NH-C_{1.4}$ -alkyl, $SO_2-N(C_{1.4}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1.4}$ -alkyl, and $N(C_{1.4}$ -alkyl) $_2$, wherein: $R^{5.1}$ denotes $C_{1.4}$ -alkyl, C_{6-10} -aryl- $C_{1.6}$ -alkylene, or C_{5-10} -heteroaryl- $C_{1.6}$ -alkylene, and $R^{5.2}$ denotes C_{6-10} -aryl or a C_{5-10} -heteroaryl;

[0057] R^6 denotes H, $C_{1.4}$ -alkyl, or $C_{1.4}$ -haloalkyl; and

[0058] R^7 denotes H, $C_{1.4}$ -alkyl, or $C_{1.4}$ -haloalkyl; or

[0059] R^6 and R^7 together form a 3-6 membered carbocycle,

optionally in the form of the racemates, enantiomers, diastereomers thereof, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0060] Preferred compounds of formula 1 above are those wherein:

[0061] A denotes CO, $C=NH$, $C_{1.6}$ -alkylene, or C_{3-8} -cycloalkylene,

[0062] B^1 denotes phenyl or an aromatic or non-aromatic ring, which may optionally contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen and which may optionally be mono- or polysubstituted by one or more groups selected from among OH, $O-C_{1.6}$ -alkyl, $O-C_{1.6}$ -haloalkyl, halogen, $C_{1.6}$ -alkyl, and $C_{1.6}$ -haloalkyl;

[0063] B^2 denotes phenyl or pyridinyl;

[0064] X denotes O, S, NR^5 , or CR^6R^7 ;

[0065] n denotes 0, 1, 2, or 3;

[0066] R^1 denotes H, $C_{1.4}$ -alkyl, or $C_{1.4}$ -haloalkyl;

[0067] R^2 denotes H, $C_{1.4}$ -alkyl, or $C_{1.4}$ -haloalkyl; or

[0068] R^1 and R^2 together with the nitrogen form a non-aromatic heterocycle, which may contain one or two nitrogen atoms;

[0069] R^3 denotes H, OH, $C_{1.6}$ -haloalkyl, or a group selected from among C_{6-10} -aryl, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may optionally be substituted by a methyl group, oxo, or OH; or


[0070] R^3 denotes $C_{1.6}$ -alkyl, which may optionally be substituted by one or more groups selected from among halogen, OH, CN, $CONH_2$, $CONH-C_{1.6}$ -alkyl, $CON(C_{1.6}$ -alkyl) $_2$, $COOH$, $COO-C_{1.6}$ -alkyl, COH , $CO-C_{1.6}$ -alkyl, COaryl, OH, $O-C_{1.6}$ -alkyl, $O-C_{1.6}$ -haloalkyl, halogen, SH, $S-C_{1.6}$ -alkyl, $S-C_{1.6}$ -haloalkyl, $SO_2-C_{1.6}$ -alkanol; $SO_2-C_{1.6}$ -alkyl, $SO_2-C_{1.6}$ -haloalkyl, SO_2-NH_2 , $SO_2-NH-C_{1.6}$ -alkyl, $SO_2-N(C_{1.6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1.6}$ -alkyl, and $N(C_{1.6}$ -alkyl) $_2$; or

[0071] R^3 denotes a group selected from among C_{3-8} -cycloalkyl, a C_{3-8} -cycloalkyl bridged with $C_{1.3}$ -alkylene,

a C_{5-8} -cycloalkenyl, $C_{1.6}$ -alkyl, and $C_{1.6}$ -alkanol, which may optionally be substituted by one or more groups selected from among C_{6-10} -aryl, C_{3-8} -cycloalkyl, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may optionally be substituted by one or more groups selected from among $C_{1.6}$ -alkyl, $C_{1.6}$ -haloalkyl, CN, OH, $O-C_{1.6}$ -alkyl, $O-C_{1.6}$ -haloalkyl, halogen, $S-C_{1.6}$ -alkyl, $S-C_{1.6}$ -haloalkyl, NO_2 , NH_2 , $NH-C_{1.6}$ -alkyl, $N(C_{1.6}$ -alkyl) $_2$, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, while the C_{5-10} -heteroaryl and the C_{3-10} -heterocycle may optionally be substituted by a group selected from oxo, hydroxyl, halogen, $C_{1.6}$ -alkyl, and $C_{1.6}$ -haloalkyl; or

[0072] R^3 denotes a group selected from among $C_{1.6}$ -alkyl, C_{6-10} -aryl- $C_{1.6}$ -alkylene, C_{5-10} -heteroaryl- $C_{1.6}$ -alkylene, C_{3-7} -cycloalkyl, C_{6-10} -aryl, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which is substituted by one or more groups selected from among B, halogen, OH, $C_{1.6}$ -alkyl, oxo, where B is a compound of formula 2

wherein:




2

[0073] Z^1 denotes H, OH, halogen, $C_{1.6}$ -alkyl, $C_{1.6}$ -alkanol, $O(C_{1.6}$ -alkyl), C_{6-10} -aryl, $O-C_{6-10}$ -aryl, NH_2 , $NH(C_{1.6}$ -alkyl), $N(C_{1.6}$ -alkyl) $_2$, or C_{3-7} -cycloalkyl; and Z^2 denotes OH, NH_2 , $NH(C_{1.6}$ -alkyl), $N(C_{1.6}$ -alkyl) $_2$, $O(C_{1.6}$ -alkyl), mono- or bicyclic C_{3-7} -cycloalkyl, a mono- or bicyclic C_{3-10} -heterocycle, a mono- or bicyclic C_{5-10} -heteroaryl, or C_{6-10} -aryl; or

[0074] R^3 denotes phenyl, which is substituted by one or more groups selected from among C_{6-10} -aryl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, $C_{1.6}$ -haloalkyl, $CONH_2$, $CONH-C_{1.6}$ -alkyl, $CON(C_{1.6}$ -alkyl) $_2$, $COOH$, $COO-C_{1.6}$ -alkyl, COH , $CO-C_{1.6}$ -alkyl, COaryl, OH, $O-C_{1.6}$ -alkyl, $O-C_{1.4}$ -haloalkyl, halogen, SH, $S-C_{1.6}$ -alkyl, $S-C_{1.4}$ -haloalkyl, $SO_2-C_{1.6}$ -alkyl, $SO_2-C_{1.4}$ -haloalkyl, SO_2-NH_2 , $SO_2-NH-C_{1.6}$ -alkyl, $SO_2-N(C_{1.6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1.6}$ -alkyl, and $N(C_{1.6}$ -alkyl) $_2$; or

[0075] R^3 denotes phenyl, which is substituted by $C_{1.4}$ -alkyl, which may optionally be substituted by a group selected from among $COOR^{3.3}$, $NR^{3.3}R^{3.4}$, $NHCOR^{3.3}$, $NHCOOR^{3.3}$, and phenyl, which may optionally be substituted by one or more groups selected from among methyl, t-butyl, F, Cl, Br, CN, OH, and a heterocycle, which may contain one, two, or three heteroatoms selected from among oxygen and nitrogen, while the heterocycle may optionally be substituted by an oxo group or a methyl group, wherein: $R^{3.3}$ denotes H or $C_{1.6}$ -alkyl; $R^{3.4}$ denotes H, $C_{1.6}$ -alkyl, or C_{6-10} -aryl- $C_{1.6}$ -alkylene, C_{5-10} -heteroaryl- $C_{1.6}$ -alkylene; or

- [0076] R^3 denotes phenyl, substituted by $NR^{3.1}R^{3.2}$, wherein:
- [0077] $R^{3.1}$ denotes H, C_{1-4} -alkyl, C_{1-4} -haloalkyl, $COR^{3.1.1}$, $COOR^{3.1.1}$, $CONR^{3.1.1}R^{3.1.2}$, or $SO_2-R^{3.1.1}$, wherein $R^{3.1.1}$ denotes H, C_{1-4} -alkyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, or C_{6-10} -aryl; $R^{3.1.2}$ denotes H, C_{1-4} -alkyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, or C_{6-10} -aryl; and
- [0078] $R^{3.2}$ denotes H, C_{1-4} -alkyl, which may optionally be substituted by one or more groups selected from among NH_2 , $NH(C_{1-4}\text{-alkyl})$, $N(C_{1-4}\text{-alkyl})_2$, oxo, or a non-aromatic C_{3-10} -heterocycle, which may contain one or two nitrogen atoms and may optionally be substituted by a methyl group; or
- [0079] R^3 denotes C_{6-10} -aryl, which may be substituted by a C_{5-10} -heteroaryl containing one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen which may optionally be substituted by one or more groups selected from among C_{6-10} -aryl, C_{1-4} -alkyl, C_{3-6} -cycloalkyl, CN, C_{1-4} -haloalkyl, $CONH_2$, $CONH-C_{1-4}$ -alkyl, $CON(C_{1-4}\text{-alkyl})_2$, $COOH$, $COO-C_{1-4}$ -alkyl, COH , $CO-C_{1-4}$ -alkyl, OH, $O-C_{1-4}$ -alkyl, halogen, SH, $S-C_{1-4}$ -alkyl, NH_2 , $NH-C_{1-4}$ -alkyl, and $N(C_{1-4}\text{-alkyl})_2$; or
- [0080] R^3 denotes C_{6-10} -aryl, which may be substituted by a non-aromatic C_{3-10} -heterocycle, which may contain one or two heteroatoms, selected from among oxygen, sulfur, and nitrogen, while the C_{3-10} -heterocycle may optionally be substituted by one or more groups selected from among C_{1-4} -alkyl and oxo; or
- [0081] R^3 denotes benzimidazolyl, which may optionally be substituted by one group or several groups selected from among C_{1-4} -alkyl, C_{1-4} -haloalkyl, and C_{3-6} -cycloalkyl;
- [0082] R^4 denotes C_{1-4} -alkyl, C_{3-6} -cycloalkyl, C_{1-4} -haloalkyl, $OR^{4.1}$, $NR^{4.1}R^{4.2}$, CN, or halogen, wherein $R^{4.1}$ denotes H, C_{1-4} -alkyl, C_{3-6} -cycloalkyl, or C_{1-4} -haloalkyl; $R^{4.2}$ denotes H, C_{1-4} -alkyl, C_{3-6} -cycloalkyl, or C_{1-4} -haloalkyl;
- [0083] R^5 denotes C_{1-4} -alkyl, C_{3-8} -cycloalkyl, C_{1-4} -haloalkyl, $COR^{5.1}$, $CONHR^{5.1}$, $CON(R^{5.1})_2$, SO_2-C_{1-4} -alkyl, SO_2-C_{1-4} -haloalkyl, SO_2 -aryl, or a group selected from among $R^{5.2}$ SO_2-C_{1-4} -alkyl- $R^{5.2}$, and C_{1-4} -alkyl- $R^{5.2}$, which may optionally be substituted by C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, CN, C_{1-4} -haloalkyl, $CONH_2$, $CONH-C_{1-4}$ -alkyl, $CON(C_{1-4}\text{-alkyl})_2$, $COOH$, $COO-C_{1-4}$ -alkyl, COH , $CO-C_{1-4}$ -alkyl, $O-C_{1-4}$ -cycloalkyl, $O-C_{1-4}$ -haloalkyl, $O-C_{1-4}$ -alkyl, halogen, SO_2-C_{1-4} -alkyl, SO_2-NH_2 , SO_2-NH-C_{1-4} -alkyl, $SO_2-N(C_{1-4}\text{-alkyl})_2$, NO_2 , NH_2 , $NH-C_{1-4}$ -alkyl, and $N(C_{1-4}\text{-alkyl})_2$, wherein $R^{5.1}$ denotes C_{1-6} -alkyl, C_{5-10} -heteroaryl- C_{1-6} -alkylene, or C_{6-10} -aryl- C_{1-6} -alkylene, and $R^{5.2}$ denotes C_{6-10} -aryl, or a C_{5-10} -heteroaryl;
- [0084] R^6 denotes H, C_{1-6} -alkyl, or C_{1-6} -haloalkyl; and
- [0085] R^7 denotes H, C_{1-6} -alkyl, or C_{1-6} -haloalkyl; or
- [0086] R^6 and R^7 together form 3-6 membered carbocycle, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.
- [0087] Preferred compounds of formula 1 above are those wherein R^1 , R^2 , R^4 , R^5 , R^7 , and A, B^1 , B^2 , X, and n have the meanings given above, and wherein:
- [0088] R^3 denotes C_{1-10} -alkyl, which may optionally be substituted by one or more groups selected from among halogen, OH, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}\text{-alkyl})_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkanol; SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}\text{-alkyl})_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, and $N(C_{1-6}\text{-alkyl})_2$, or
- [0089] R^3 denotes a group selected from among C_{3-8} -cycloalkyl, a C_{3-8} -cycloalkyl bridged with C_{1-3} -alkylene, a C_{5-8} -cycloalkenyl, C_{1-6} -alkyl, and C_{1-6} -alkanol, which may optionally be substituted by one or more groups selected from among C_{6-10} -aryl, C_{3-8} -cycloalkyl, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which in turn may optionally be substituted by one or more groups selected from among C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}\text{-alkyl})_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}\text{-alkyl})_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, $N(C_{1-6}\text{-alkyl})_2$, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may optionally be substituted by one or more groups selected from oxo, hydroxyl, halogen, or C_{1-6} -alkyl, and C_{1-6} -haloalkyl; or
- [0090] R^3 denotes a group selected from among C_{6-10} -aryl, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may be substituted by one or more groups selected from among C_{6-10} -aryl, C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}\text{-alkyl})_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}\text{-alkyl})_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, $N(C_{1-6}\text{-alkyl})_2$, and $N-(SO_2-C_{1-4}\text{-alkyl})(R^{3.4})$ or
- [0091] R^3 denotes a group selected from among C_{1-6} -alkyl, C_{6-10} -aryl- C_{1-6} -alkylene, C_{5-10} -heteroaryl- C_{1-6} -alkylene, C_{3-7} -cycloalkyl, C_{6-10} -aryl, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may optionally be substituted by one or more groups selected from among B, halogen, OH, C_{1-6} -alkyl, and oxo, while B is a compound of formula 2



[0092] wherein:

[0093] Z^1 denotes H, OH, halogen, C_{1-6} -alkyl, C_{1-6} -alkanol, $O(C_{1-6}$ -alkyl), C_{6-10} -aryl, $O-C_{6-10}$ -aryl, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, or C_{3-7} -cycloalkyl, and

[0094] Z^2 denotes OH, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, $O(C_{1-6}$ -alkyl), mono- or bicyclic C_{3-7} -cycloalkyl, mono- or bicyclic C_{5-10} -heteroaryl, mono- or bicyclic C_{3-10} -heterocycle, or C_{6-10} -aryl,

optionally in the form of the racemates, enantiomers, diastereomers thereof, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0095] Preferred compounds of formula 1 above are those wherein R^3 and R^4 have the above-mentioned meanings and wherein

[0096] A denotes CH_2 , CD_2 , $C=NH$, CHMe, CMe $_2$, 1,1'-cyclopropylene, or 1,1'-cyclobutylidene;

[0097] B^1 denotes phenyl;

[0098] B^2 denotes phenyl;

[0099] X denotes O or NR^5 ; wherein R^5 denotes methyl, ethyl, cyclopropyl, cyclobutyl, $CONHCH_2$ -phenyl, CH_2CF_3 , or benzyl, which may optionally be substituted by F; and wherein

[0100] n denotes 0 or 1;

[0101] R^1 denotes H; and

[0102] R^2 denotes H,

optionally in the form of the racemates, enantiomers, diastereomers thereof, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.


[0103] Preferred are the above compounds of formula 1, wherein R^1 , R^2 , R^3 , R^6 , R^7 , and A, B^1 , B^2 , X, and n have the meanings given above and wherein:

[0104] R^3 denotes C_{1-6} -alkyl, which may optionally be substituted by one or more groups selected from among halogen, OH, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, COOH, $COO-C_{1-6}$ -alkyl, COH, $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, SO_2-C_{1-6} -alkanol; SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, and $N(C_{1-6}$ -alkyl) $_2$, or

[0105] R^3 denotes a group selected from among C_{3-8} -cycloalkyl, a C_{3-8} -cycloalkyl bridged by C_{1-3} -alkylene and C_{1-6} -alkyl, which may optionally be substituted by one or more groups selected from among C_{6-10} -aryl, C_{3-8} -cycloalkyl, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may in turn optionally be substituted by one or more groups selected from among C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, COOH, $COO-C_{1-6}$ -alkyl, COH, $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, $N(C_{1-6}$ -alkyl) $_2$, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may optionally be substituted by one or more groups selected from oxo, hydroxyl, halogen, or C_{1-6} -alkyl, and C_{1-6} -haloalkyl; or

[0106] R^3 denotes a group selected from among C_{6-10} -aryl, a C_{3-8} -heterocycle with 1 to 4 heteroatoms selected from N, O, S, and a C_{5-10} -heteroaryl with 1 to 2 heteroatoms selected from N, O, S, which may optionally be substituted by one or more groups selected from among C_{6-10} -aryl, C_{1-6} -alkyl, C_{1-6} -haloalkyl, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, COOH, $COO-C_{1-6}$ -alkyl, COH, $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, $N(C_{1-6}$ -alkyl) $_2$, and $N-(SO_2-C_{1-4}$ -alkyl)($R^{3,4}$), wherein $R^{3,4}$ is a C_{5-10} -heteroaryl- C_{1-6} -alkylene; or

[0107] R^3 denotes a group selected from among C_{1-6} -alkyl, C_{6-10} -aryl- C_{1-6} -alkylene, C_{5-10} -heteroaryl- C_{1-6} -alkylene, C_{3-7} -cycloalkyl, C_{6-10} -aryl, C_{3-8} -heterocycle with 1 to 4 heteroatoms selected from N, O, S, and a C_{5-10} -heteroaryl with 1 to 2 heteroatoms selected from N, O, S, which may optionally be substituted in each case by one or more groups selected from among B, halogen, OH, C_{1-6} -alkyl, oxo, wherein B is a compound of formula 2




[0108] wherein:

[0109] Z^1 is H, OH, halogen, C_{1-6} -alkyl, C_{1-6} -alkanol, or $O(C_{1-6}$ -alkyl) and

[0110] Z^2 is OH, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, $O(C_{1-6}$ -alkyl), mono- or bicyclic C_{3-7} -cycloalkyl,

- mono- or bicyclic C₅₋₁₀-heteroaryl, mono- or bicyclic C₃₋₁₀-heterocycle, or C₆₋₁₀-aryl; and
- [0111] R⁴ denotes H, F, or Cl,
- optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.
- [0112] Preferred are the above compounds of formula 1, wherein
- [0113] A denotes CO, C=NH, C₁₋₆-alkylene, or C₃₋₈-cycloalkylene,
- [0114] B¹ denotes phenyl or pyridinyl;
- [0115] B² denotes phenyl or pyridinyl;
- [0116] X denotes O or NR⁵;
- [0117] n denotes 0, 1, 2, or 3;
- [0118] R¹ denotes H, methyl, ethyl, or propyl;
- [0119] R² denotes H, methyl, ethyl, or propyl;
- [0120] R³ denotes H, OH, C₁₋₆-haloalkyl, or C₆₋₁₀-aryl, or a group selected from among a C₅₋₁₀-heteroaryl and a C₃₋₁₀-cycloalkyl, which may contain one, two, or three nitrogen atoms and which may optionally be substituted by a methyl group; or
- [0121] R³ denotes a group selected from among cyclopentyl, cyclohexyl, cyclopentenyl, cyclohexenyl, methyl, ethyl, propyl, and butyl, which may optionally be substituted by one or more groups selected from among C₆₋₁₀-aryl and a C₅₋₁₀-heterocycle, which may in turn optionally be substituted by one or more groups selected from among C₁₋₆-alkyl, C₁₋₆-haloalkyl, CN, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂; or
- [0122] R³ denotes phenyl, which may optionally be substituted by one or more groups selected from among C₆₋₁₀-aryl, C₂₋₄-alkenyl, C₂₋₄-alkynyl, C₁₋₄-haloalkyl, CONH₂, CONH—C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO—C₁₋₄-alkyl, COH, CO—C₁₋₄-alkyl, CO—C₆₋₁₀-aryl, OH, O—C₁₋₄-alkyl, O—C₁₋₄-haloalkyl, halogen, SH, S—C₁₋₄-alkyl, S—C₁₋₄-haloalkyl, SO₂—C₁₋₄-alkyl, SO₂—C₁₋₄-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₄-alkyl, SO₂—N(C₁₋₄-alkyl)₂, NO₂, NO₂, NH₂, NH—C₁₋₄-alkyl, and N(C₁₋₄-alkyl)₂; or
- [0123] R³ denotes phenyl, which may optionally be substituted by C₁₋₄-alkyl, which may in turn optionally be substituted by a group selected from among COOR^{3.3}, NR^{3.3}R^{3.4}, NHCOR^{3.3}, NHCOOR^{3.3}, p-fluorophenyl, and a heterocycle, which may contain one, two, or three heteroatoms selected from among oxygen and nitrogen and which may optionally be substituted by an oxo group, wherein R^{3.3} denotes H or C₁₋₄-alkyl; and R^{3.4} denotes H, C₁₋₄-alkyl, C₆₋₁₀-aryl-C₁₋₆-alkylene, or C₅₋₁₀-heteroaryl-C₁₋₆-alkylene; or
- [0124] R³ denotes phenyl, which may be substituted by NR^{3.1}R^{3.2}, wherein:
- [0125] R^{3.1} denotes H, C₁₋₄-alkyl, COR^{3.1.1}, COOR^{3.1.1}, CONR^{3.1.1}R^{3.1.2}, or SO₂—R^{3.1.1}, wherein
- [0126] R^{3.1.1} denotes H, C₁₋₄-alkyl, or C₆₋₁₀-aryl, and R^{3.1.2} denotes H, C₁₋₄-alkyl, or C₆₋₁₀-aryl, and
- [0127] R^{3.2} denotes H, C₁₋₄-alkyl, which may optionally be substituted by one or more groups selected from among NH₂, NH(C₁₋₄-alkyl), N(C₁₋₄-alkyl)₂, oxo, or a C₃₋₁₀-heterocycle, which may contain one or two nitrogen atoms and which may optionally be substituted by a methyl group; or
- [0128] R³ denotes phenyl, which may be substituted by a C₅₋₁₀-heteroaryl, which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen, while the C₅₋₁₀-heteroaryl may optionally be substituted by one or more groups selected from among C₆₋₁₀-aryl, C₁₋₄-alkyl, C₁₋₄-haloalkyl, C₃₋₆-cycloalkyl, CN, CONH₂, CONH—C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO—C₁₋₄-alkyl, COH, CO—C₁₋₄-alkyl, OH, O—C₁₋₄-alkyl, halogen, NH₂, and N(C₁₋₄-alkyl)₂; or
- [0129] R³ denotes phenyl, which may be substituted by a C₅₋₁₀-heteroaryl, which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen, while the C₅₋₁₀-heteroaryl may optionally be substituted by one or more groups selected from among C₁₋₄-alkyl and oxo; or
- [0130] R³ denotes benzimidazolyl, which may optionally be substituted by one or more groups selected from among methyl, ethyl, propyl, CF₃, CH₂CF₃, cyclopropyl, cyclopentyl, and cyclohexyl;
- [0131] R⁴ denotes C₁₋₄-alkyl, C₁₋₄-haloalkyl, or halogen; and
- [0132] R⁵ denotes a group selected from among C₁₋₄-alkyl, C₃₋₆-cycloalkyl, COR^{5.1}, CONHR^{5.1}, C₆₋₁₀-aryl, SO₂—C₆₋₁₀-aryl-C₁₋₆-alkylene, SO₂—C₆₋₁₀-aryl, or C₆₋₁₀-aryl-C₁₋₆-alkylene, and C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, which may optionally be substituted by one or more groups selected from among C₁₋₄-alkyl, C₂₋₄-alkenyl, C₂₋₄-alkynyl, CN, C₁₋₄-haloalkyl, CONH₂, CONH—C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO—C₁₋₄-alkyl, COH, CO—C₁₋₄-alkyl, OH, O—C₁₋₄-alkyl, halogen, SO₂—C₁₋₄-alkyl, SO₂—NH₂, SO₂—NH—C₁₋₄-alkyl, SO₂—N(C₁₋₄-alkyl)₂, NO₂, NH₂, NH—C₁₋₄-alkyl, and N(C₁₋₄-alkyl)₂; and R^{5.1} denotes C₁₋₄-alkyl, or C₆₋₁₀-aryl-C₁₋₆-alkylene,
- optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0133] Particularly preferred are the compounds of formula 1a



1a

wherein A, X, n, R¹, R², R³, and R⁴ are as hereinbefore defined, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0134] Particularly preferred are the above compounds of formula 1 or 1a, wherein R¹, R², R³, R⁴, and A, B¹, B², and n have the meanings given above and wherein X denotes O, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0135] Particularly preferred are the above compounds of formula 1 or 1a, wherein R¹, R², R³, R⁴, R⁵, and A, B¹, B², and n have the meanings given above and wherein X denotes NR⁵, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0136] Particularly preferred are the above compounds of formula 1 or 1a, wherein R¹, R², R³, R⁴, R⁶, and R⁷ as well as A, B¹, B², and n have the meanings given above and wherein X denotes CR⁶R⁷, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0137] Particularly preferred are the above compounds of formula 1 or 1a, wherein R¹, R², R³, R⁴, R⁵, R⁶, R⁷ as well as A, B¹, B², and X have the meanings given above and wherein n denotes 0, 1, or 2, preferably 0 or 1, particularly preferably 0, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0138] Particularly preferred are the above compounds of formula 1 or 1a, wherein:

[0139] A denotes CH₂, CHMe, CMe₂, C=NH, 1,1'-cyclopropylene, 1,1'-cyclobutylidene;

[0140] B¹ denotes phenyl;

[0141] B² denotes phenyl;

[0142] X denotes O or NR⁵;

[0143] n denotes 0, 1, or 2;

[0144] R¹ denotes H, methyl, or ethyl;

[0145] R² denotes H, methyl, or ethyl;

[0146] R³ denotes H, cyclopropyl, cyclobutyl, N-methylpiperidinyl, pyridinyl, phenyl, or 4-phenylcyclohexanyl, or

[0147] R³ denotes phenyl, which may optionally be substituted by one or more groups selected from among phenyl, methyl, ethyl, propyl, butyl, CF₃, CONH₂, CONHMe, CONMe₂, COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, SH, SO₂Me, SONH₂, SONMe₂, NO₂, NH₂, NHMe, and NMe₂; or

[0148] R³ denotes phenyl, which is optionally substituted by a group selected from among methyl and ethyl, which may optionally be substituted by one or more groups selected from among COOH, COOMe, NH₂, NMe₂, NHCOMe, NHCOO-tert-butyl, NMe(benzyl), p-fluorophenyl, pyrrolidinyl, piperidinyl, morpholinyl, pyrrolidin-2-onyl, imidazolyl, and triazolyl; or

[0149] R³ denotes phenyl, which is substituted by NR^{3.1}R^{3.2}, wherein R^{3.1} denotes H, methyl, COH, COMe, COOMe, CONH₂, CONMe₂, SO₂Me, SO₂CF₃, or SO₂-phenyl, and R^{3.2} denotes H or a group selected from among methyl and ethyl, which may optionally be substituted by one or more groups selected from among NH₂, NHMe, NMe₂, N-piperidinyl, N-morpholinyl, and N-methylpiperazinyl, wherein the N-piperidinyl, N-morpholinyl, and the N-methylpiperazinyl may optionally be substituted by a further oxo; or

[0150] R³ denotes phenyl, which may be substituted by a C₅₋₁₀-heteroaryl, which may contain one, is two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen, while the C₅₋₁₀-heteroaryl may optionally be substituted by one or more groups selected from among phenyl, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, CF₃, CN, CONH₂, CONMe₂, CONEt₂, COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, NH₂, NMe₂, NEt₂, and NPr₂; or

[0151] R³ denotes phenyl which is substituted by a C₃₋₁₀-heterocycle, which may contain one or two heteroatoms selected from among oxygen and nitrogen, while the C₃₋₁₀-heterocycle may optionally be substituted by one or more groups selected from among C₁₋₄-alkyl and oxo; or

[0152] R³ denotes benzimidazolyl, which may optionally be substituted by one or more groups selected from among methyl, propyl, CF₃, CH₂CF₃, cyclopropyl, and cyclohexyl;

[0153] R⁴ denotes methyl, ethyl, propyl, butyl, CF₃, CH₂CF₃, F, Cl, or Br; and

[0154] R⁵ denotes a group selected from among methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, CF₃, CH₂CF₃, COR^{5.1}, CONHR^{5.1}, phenyl, phenylsulfonyl, and benzyl, while benzyl may optionally be substituted by one or more groups selected from among methyl, ethyl, propyl, butyl, CF₃, CN, CONH₂, CONMe₂, CONEt₂, COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, SO₂Me, SONH₂, SONMe₂, NO₂, NH₂, NMe₂, NEt₂, and NPr₂, wherein R^{5.1} denotes methyl, ethyl, propyl, butyl, or benzyl,

optionally in the form of the racemates, enantiomers, diastereomers thereof, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0155] Particularly preferred are the above compounds of formula 1 or 1a, wherein:

[0156] A denotes CH_2 , CHMe, CMe_2 , CO, $\text{C}=\text{NH}$, 1,1'-cyclopropylene, or 1,1'-cyclobutylidene;

[0157] B^1 denotes phenyl;

[0158] B^2 denotes phenyl;

[0159] X denotes O or NR^5 ;

[0160] n denotes 0 or 1;

[0161] R^1 denotes H, methyl, or ethyl, preferably H;

[0162] R^2 denotes H, methyl, or ethyl, preferably H;

[0163] R^3 denotes H, OH, cyclopropyl, cyclobutyl, N-methylpiperidinyl, pyridinyl, phenyl, or 4-phenylcyclohexanyl, or

[0164] R^3 denotes phenyl, which may be substituted by a group selected from among phenyl, OH, F, and CONH_2 ; or

[0165] R^3 denotes phenyl, which is substituted by a group selected from among methyl and ethyl, which may optionally be substituted by a group selected from among COOH, COOMe, NH_2 , NMe_2 , NHCOMe , NHCOO-tert-butyl , $\text{NMe}(\text{benzyl})$, p-fluorophenyl, pyrrolidinyl, piperidinyl, morpholinyl, pyrrolidin-2-onyl, imidazolyl, and triazolyl; or

[0166] R^3 denotes phenyl, which is substituted by $\text{NR}^{3.1}\text{R}^{3.2}$, wherein $\text{R}^{3.1}$ denotes H, methyl, SO_2Me , SO_2CF_3 , or $\text{SO}_2\text{-phenyl}$; $\text{R}^{3.2}$ denotes H or a group selected from among methyl and ethyl, which may optionally be substituted by one or more groups selected from among NH_2 , NHMe , NMe_2 , oxo, N-piperidinyl, N-morpholinyl, and N-methylpiperazinyl; or

[0167] R^3 denotes phenyl, which may be substituted by a C_{5-10} -heteroaryl, which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen and which may optionally be substituted by one or more groups selected from among phenyl, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, CF_3 , CN, CONH_2 , CONMe_2 , CONEt_2 , COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, SO_2Me , SONH_2 , SONMe_2 , NO_2 , NH_2 , NMe_2 , NEt_2 , and NPr_2 , or

[0168] R^3 denotes phenyl, which is substituted by a C_{3-10} -heterocycle which may contain one or two heteroatoms selected from among oxygen and nitrogen and may optionally be substituted by one or more groups selected from among C_{1-4} -alkyl and oxo, or

[0169] R^3 denotes benzimidazolyl, which may optionally be substituted by methyl;

[0170] R^4 denotes F or Cl; and

[0171] R^5 denotes a group selected from among methyl, ethyl, cyclopropyl, COMe, $\text{CONHR}^{5.1}$, phenyl, phenylsulfonyl, and benzyl, which may optionally be substituted by F, wherein $\text{R}^{5.1}$ denotes butyl or benzyl,

optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0172] Particularly preferred are the above compounds of formula 1 or 1a, wherein:

[0173] A denotes CH_2 , CHMe, CMe_2 , 1,1'-cyclopropylene, 1,1'-cyclobutylidene;

[0174] B^1 denotes phenyl;

[0175] B^2 denotes phenyl;

[0176] X denotes O or NR^5 ;

[0177] n denotes 0 or 1;

[0178] R^1 denotes H;

[0179] R^2 denotes H;

[0180] R^3 denotes H or 4-phenylcyclohexanyl; or

[0181] R^3 denotes phenyl, which may optionally be substituted by $\text{NR}^{3.1}\text{R}^{3.2}$, wherein $\text{R}^{3.1}$ denotes H, methyl, SO_2Me , SO_2CF_3 , or $\text{SO}_2\text{-phenyl}$; and $\text{R}^{3.2}$ denotes H or a group selected from among methyl and ethyl, which may optionally be substituted by one or more groups selected from among NH_2 , NHMe , NMe_2 , oxo, N-piperidinyl, N-morpholinyl, and N-methylpiperazinyl, or

[0182] R^3 denotes phenyl, which is substituted by a C_{5-10} -heteroaryl, which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen, while the C_{5-10} -heteroaryl may optionally be substituted by one or more groups selected from among phenyl, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, CF_3 , CN, CONH_2 , CONMe_2 , CONEt_2 , COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, SO_2Me , SONH_2 , SONMe_2 , NO_2 , NH_2 , NMe_2 , NEt_2 , and NPr_2 , or

[0183] R^3 denotes phenyl, which is substituted by a C_{3-10} -heterocycle, which may contain one or two heteroatoms selected from among oxygen and nitrogen, while the C_{3-10} -heterocycle may optionally be substituted by one or more groups selected from among C_{1-4} -alkyl and oxo;

[0184] R^4 denotes H, F, or Cl; and

[0185] R^5 denotes methyl, ethyl, cyclopropyl, cyclobutyl, CH_2CF_3 , or benzyl, while the benzyl may optionally be substituted by F,

optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0186] Also particularly preferred are the above compounds of formula 1 or 1a, wherein:

[0187] A denotes a group selected from among CH_2 , CHMe, CMe_2 , CO, $\text{C}=\text{NH}$, and



[0188] B¹ denotes phenyl;

[0189] B² denotes phenyl;

[0190] X denotes O or NR⁵;











[0191] n denotes 0 or 1;

[0192] R¹ denotes H;

[0193] R² denotes H;


[0194] R³ denotes H, OH, 4-phenylcyclohexyl, or a group selected from among

-continued



[0195] R^4 denotes H, F, or Cl; and

[0196] R^5 denotes methyl or a group selected from among



optionally in the form of the racemates, enantiomers, diastereomers thereof, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0197] Also particularly preferred are the above compounds of formula 1 or 1a, wherein:

[0198] A denotes CH_2 , CD_2 , $\text{C}=\text{NH}$, CHMe , CMe_2 , 1,1'-cyclopropylene, 1,1'-cyclobutylidene;

[0199] B^1 denotes phenyl;

[0200] B^2 denotes phenyl;

[0201] X denotes O or NR^5 , wherein R^5 denotes methyl, ethyl, cyclopropyl, cyclobutyl, CONHCH_2 -phenyl, CH_2CF_3 , or benzyl, optionally substituted by F;

[0202] n denotes 0 or 1;

[0203] R^1 denotes H;

[0204] R^2 denotes H;


[0205] R^3 denotes C_{1-6} -alkyl, optionally substituted by one or more groups selected from among halogen, OH, CN, CONH_2 , $\text{CONH}-\text{C}_{1-6}$ -alkyl, $\text{CON}(\text{C}_{1-6}$ -alkyl) $_2$, COOH , $\text{COO}-\text{C}_{1-6}$ -alkyl, COH , $\text{CO}-\text{C}_{1-6}$ -alkyl, COaryl , OH , $\text{O}-\text{C}_{1-6}$ -alkyl, $\text{O}-\text{C}_{1-6}$ -haloalkyl, halogen, SH, $\text{S}-\text{C}_{1-6}$ -alkyl, $\text{S}-\text{C}_{1-6}$ -haloalkyl, $\text{SO}_2-\text{C}_{1-6}$ -alkanol; $\text{SO}_2-\text{C}_{1-6}$ -alkyl, $\text{SO}_2-\text{C}_{1-6}$ -haloalkyl, SO_2-NH_2 , $\text{SO}_2-\text{NH}-\text{C}_{1-6}$ -alkyl, $\text{SO}_2-\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $\text{NH}-\text{C}_{1-6}$ -alkyl, and $\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, or

[0206] R^3 denotes a group selected from among C_{3-8} -cycloalkyl, a C_{3-8} -cycloalkyl bridged by C_{1-3} -alkylene, or C_{5-8} -cycloalkenyl, and C_{1-6} -alkyl, C_{1-6} -alkanol, which may optionally be substituted in each case by one or more groups selected from among C_{6-10} -aryl, C_{3-8} -cycloalkyl, C_{5-10} -heteroaryl, and C_{3-10} -heterocycle, which may optionally be substituted by one or more groups selected from among C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, CN, CONH_2 , $\text{CONH}-\text{C}_{1-6}$ -alkyl, $\text{CON}(\text{C}_{1-6}$ -alkyl) $_2$, COOH , $\text{COO}-\text{C}_{1-6}$ -alkyl, COH , $\text{CO}-\text{C}_{1-6}$ -

alkyl, $\text{CO}-\text{C}_{6-10}$ -aryl, OH, $\text{O}-\text{C}_{1-6}$ -alkyl, $\text{O}-\text{C}_{1-6}$ -haloalkyl, halogen, SH, $\text{S}-\text{C}_{1-6}$ -alkyl, $\text{S}-\text{C}_{1-6}$ -haloalkyl, $\text{SO}_2-\text{C}_{1-6}$ -alkanol, $\text{SO}_2-\text{C}_{1-6}$ -alkyl, $\text{SO}_2-\text{C}_{1-6}$ -haloalkyl, SO_2-NH_2 , $\text{SO}_2-\text{NH}-\text{C}_{1-6}$ -alkyl, $\text{SO}_2-\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $\text{NH}-\text{C}_{1-6}$ -alkyl, $\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may optionally be substituted by a group selected from oxo, hydroxyl, halogen, or C_{1-6} -alkyl, and C_{1-6} -haloalkyl; or

[0207] R^3 denotes a group selected from among C_{6-10} -aryl, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may be substituted in each case by one or more groups selected from among C_{6-10} -aryl, C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, CONH_2 , $\text{CONH}-\text{C}_{1-6}$ -alkyl, $\text{CON}(\text{C}_{1-6}$ -alkyl) $_2$, COOH , $\text{COO}-\text{C}_{1-6}$ -alkyl, COH , $\text{CO}-\text{C}_{1-6}$ -alkyl, $\text{CO}-\text{C}_{6-10}$ -aryl, OH, $\text{O}-\text{C}_{1-6}$ -alkyl, $\text{O}-\text{C}_{1-6}$ -haloalkyl, halogen, SH, $\text{S}-\text{C}_{1-6}$ -alkyl, $\text{S}-\text{C}_{1-6}$ -haloalkyl, $\text{SO}_2-\text{C}_{1-6}$ -alkyl, $\text{SO}_2-\text{C}_{1-6}$ -alkanol, $\text{SO}_2-\text{C}_{1-6}$ -haloalkyl, SO_2-NH_2 , $\text{SO}_2-\text{NH}-\text{C}_{1-6}$ -alkyl, $\text{SO}_2-\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $\text{NH}-\text{C}_{1-6}$ -alkyl, $\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, and $\text{N}(\text{SO}_2-\text{C}_{1-4}$ -alkyl)($R^{3,4}$), wherein $R^{3,4}$ is a C_{6-10} -aryl, C_{6-10} -aryl- C_{1-6} -alkylene, or a C_{5-10} -heteroaryl- C_{1-6} -alkylene; or

[0208] R^3 denotes a group selected from among C_{1-6} -alkyl, C_{6-10} -aryl- C_{1-6} -alkylene, C_{5-10} -heteroaryl- C_{1-6} -alkylene, C_{3-7} -cycloalkyl, C_{6-10} -aryl, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, which may optionally be substituted in each case by one or more groups selected from among B, halogen, OH, C_{1-6} -alkyl, oxo, where B is a compound of formula 2



[0209] wherein Z^1 denotes H, OH, halogen, C_{1-6} -alkyl, C_{1-6} -alkanol, $\text{O}(\text{C}_{1-6}$ -alkyl), C_{6-10} -aryl, $\text{O}-\text{C}_{6-10}$ -aryl, NH_2 , $\text{NH}(\text{C}_{1-6}$ -alkyl), $\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, or C_{3-7} -cycloalkyl; and Z^2 OH, NH_2 , $\text{NH}(\text{C}_{1-6}$ -alkyl), $\text{N}(\text{C}_{1-6}$ -alkyl) $_2$, $\text{O}(\text{C}_{1-6}$ -alkyl), C_{6-10} -aryl; mono- or bicyclic C_{3-7} -cycloalkyl, mono- or bicyclic aromatic or non-aromatic C_{3-10} -heterocycle, and

[0210] R^4 denotes H, F, or Cl,

optionally in the form of the racemates, enantiomers, diastereomers thereof, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0211] Also particularly preferred are compounds of formula 1 or 1a, wherein:

[0212] A denotes CH_2 , CD_2 , $\text{C}=\text{NH}$, CHMe , CMe_2 , 1,1'-cyclopropylene, 1,1'-cyclobutylidene;

[0213] B^1 denotes phenyl;

[0214] B² denotes phenyl;
 [0215] X denotes O or NR⁵, wherein R⁵ denotes methyl, ethyl, cyclopropyl, cyclobutyl, CONHCH₂-phenyl, CH₂CF₃, or benzyl, optionally substituted with F;

[0216] n denotes 0 or 1;

[0217] R¹ denotes H;

[0218] R² denotes H;


[0219] R³ denotes C₁₋₆-alkyl, which may optionally be substituted by one or more groups selected from among halogen, OH, CN, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, phenyl, CO-phenyl, OH, O—C₁₋₆-alkyl, SO₂—C₁₋₆-alkanol; SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂; or

[0220] R³ denotes a group selected from among C₃₋₈-cycloalkyl, a C₃₋₈-cycloalkyl bridged by C₁₋₃-alkylene and C₁₋₆-alkyl, which may optionally be substituted in each case by one or more groups selected from among phenyl or another aromatic or non-aromatic C₃₋₈ ring, preferably a C₅₋₇ ring, more preferably a C₅₋₆ ring, which may optionally contain in each case 1 to 4 heteroatoms independently of one another selected from N, O, S, while each of these groups may optionally be substituted by one or more groups selected from among C₁₋₆-alkyl, C₁₋₆-haloalkyl, CN, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, phenyl, CO-phenyl, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, SH, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, SO₂—C₁₋₆-alkanol, SO₂—C₁₋₆-alkyl, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂, an aromatic or non-aromatic C₃₋₈ ring, preferably a C₅₋₇ ring, more preferably a C₅₋₆ ring, which optionally contains 1 to 4 heteroatoms selected independently of one another from N, O, S, while this aromatic or non-aromatic C₃₋₈ ring may optionally be substituted by a group selected from oxo, hydroxyl, halogen, or C₁₋₆-alkyl, and C₁₋₆-haloalkyl; or

[0221] R³ denotes a group selected from among phenyl and another aromatic or non-aromatic C₃₋₈ ring, preferably a C₅₋₇ ring, more preferably a C₅₋₆ ring, which may optionally contain 1 to 4 heteroatoms selected from N, O, S, and which may optionally be substituted by one or more groups selected from among phenyl, C₁₋₆-alkyl, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, CO-phenyl, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-alkanol, SO₂—C₁₋₆-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, N(C₁₋₆-alkyl)₂, and N—(SO₂—C₁₋₄-alkyl)(R^{3,4}), wherein R^{3,4} denotes a C₅₋₁₀-heteroaryl-C₁₋₆-alkylene;

[0222] R³ denotes a group selected from among C₁₋₆-alkyl, C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, C₆₋₁₀-aryl-C₁₋₆-alkylene, C₃₋₇-cycloalkyl, phenyl, and an aromatic or non-aromatic C₃₋₈ ring, preferably a C₅₋₇ ring, more preferably a C₅₋₆ ring, which may contain 1 to 4 heteroatoms selected from N, O, and S, while each of these groups may

optionally be substituted by one or more groups selected from among B, halogen, OH, C₁₋₆-alkyl, oxo, and B is a compound of formula 2



[0223] wherein Z¹ denotes H, OH, halogen, C₁₋₆-alkyl, C₁₋₆-alkanol, or O(C₁₋₆-alkyl), and Z² denotes OH, NH₂, NH(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂, O(C₁₋₆-alkyl), mono- or bicyclic C₃₋₇-cycloalkyl, mono- or bicyclic aromatic or non-aromatic C₃₋₁₀-heterocycle, or phenyl; and

[0224] R⁴ denotes H, F, or Cl,

optionally in the form of the racemates, enantiomers, diastereomers thereof, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

[0225] Also particularly preferred are compounds of the above formula 1 or 1a, wherein:

[0226] A denotes CH₂, CD₂, C=NH, CHMe, CMe₂, 1,1'-cyclopropylene, 1,1'-cyclobutylidene;

[0227] B¹ denotes phenyl;

[0228] B² denotes phenyl;


[0229] X denotes O or NR⁵, wherein R⁵ denotes methyl, ethyl, cyclopropyl, cyclobutyl, CONHCH₂-phenyl, CH₂CF₃, or benzyl, optionally substituted by F;

[0230] n denotes 0 or 1;


[0231] R¹ denotes H;

[0232] R² denotes H;


wherein R³ denotes a group selected from among





-continued



-continued



-continued



[0236] If there is no asterisk (*) in the structural formula of the substituent, each hydrogen atom may be removed from the substituent and the valency thus freed may act as a binding site to the rest of a molecule. Thus, for example, VI may represent 2-tolyl, 3-tolyl, 4-tolyl, and benzyl



[0233] R^4 denotes H, F, or Cl,

optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof, as well as deuterated forms thereof.

Terms and Definitions Used

[0234] Unless otherwise stated, all the substituents are independent of one another. If, for example, there are a plurality of C_{1-6} -alkyl groups as substituents in one group, in the case of three substituents C_{1-6} -alkyl, one may represent methyl, one n-propyl, and one tert-butyl.


[0235] Within the scope of this application, in the definition of possible substituents, these may also be represented in the form of a structural formula. An asterisk (*) in the structural formula of the substituent is to be understood as being the linking point to the rest of the molecule. Moreover, the atom of the substituent which follows the linking point is referred to as the atom in position number 1. Thus, for example, the groups N-piperidyl (I), 4-piperidyl (II), 2-tolyl (III), 3-tolyl (IV), and 4-tolyl (V) are shown as follows:

[0237] By the term " C_{1-6} -alkyl" (including those which are part of other groups) are meant branched and unbranched alkyl groups with 1 to 6 carbon atoms and by the term " C_{1-4} -alkyl" are meant branched and unbranched alkyl groups with 1 to 4 carbon atoms. Alkyl groups with 1 to 4 carbon atoms are preferred. Examples include: methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, or hexyl. The following abbreviations may optionally also be used for the abovementioned groups: Me, Et, n-Pr, i-Pr, n-Bu, i-Bu, t-Bu, etc. Unless stated otherwise, the definitions propyl, butyl, pentyl, and hexyl include all the possible isomeric forms of the groups in question. Thus, for example, propyl includes n-propyl and isopropyl, butyl includes isobutyl, sec-butyl and tert-butyl etc.

[0238] By the term " C_{1-6} -alkylene" (including those which are part of other groups) are meant branched and unbranched alkylene groups with 1 to 6 carbon atoms and by the term " C_{1-4} -alkylene" are meant branched and unbranched alkylene groups with 1 to 4 carbon atoms. Preferred are alkylene groups with 1 to 4 carbon atoms. Examples include: methylene, ethylene, propylene, 1-methylethylene, butylene, 1-methylpropylene, 1,1-dimethylethylene, 1,2-dimethylethylene, pentylene, 1,1-dimethylpropylene,

lene, 2,2-dimethylpropylene, 1,2-dimethylpropylene, 1,3-dimethylpropylene, or hexylene. Unless stated otherwise, the definitions propylene, butylene, pentylene, and hexylene include all the possible isomeric forms of the groups in question with the same number of carbons. Thus, for example, propyl also includes 1-methylethylene and butylene includes 1-methylpropylene, 1,1-dimethylethylene, 1,2-dimethylethylene.

[0239] If the carbon chain is substituted by a group which together with one or two carbon atoms of the alkylene chain forms a carbocyclic ring with 3, 5, or 6 carbon atoms, the following examples of rings are also included:



[0240] By the term “C₂₋₆-alkenyl” (including those which are part of other groups) are meant branched and unbranched alkenyl groups with 2 to 6 carbon atoms and by the term “C₂₋₄-alkenyl” are meant branched and unbranched alkenyl groups with 2 to 4 carbon atoms, provided that they have at least one double bond. Alkenyl groups with 2 to 4 carbon atoms are preferred. Examples include: ethenyl or vinyl, propenyl, butenyl, pentenyl, or hexenyl. Unless stated otherwise, the definitions propenyl, butenyl, pentenyl, and hexenyl include all the possible isomeric forms of the groups in question. Thus, for example, propenyl includes 1-propenyl and 2-propenyl, butenyl includes 1-, 2-, and 3-butenyl, 1-methyl-1-propenyl, 1-methyl-2-propenyl, etc.

[0241] By the term “C₂₋₆-alkenylene” (including those which are part of other groups) are meant branched and unbranched alkenylene groups with 2 to 6 carbon atoms and by the term “C₂₋₄-alkenylene” are meant branched and unbranched alkenylene groups with 2 to 4 carbon atoms. Preferred are alkenylene groups with 2 to 4 carbon atoms. Examples include: ethenylene, propenylene, 1-methylethenylene, butenylene, 1-methylpropenylene, 1,1-dimethylethenylene, 1,2-dimethylethenylene, pentenylene, 1,1-dimethylpropenylene, 2,2-dimethylpropenylene, 1,2-dimethylpropenylene, 1,3-dimethylpropenylene, or hexenylene. Unless stated otherwise, the definitions propenylene, butenylene, pentenylene, and hexenylene include all the possible isomeric forms of the groups in question with the same number of carbons. Thus, for example, propenyl also includes 1-methylethenylene and butenylene includes 1-methylpropenylene, 1,1-dimethylethenylene, and 1,2-dimethylethenylene.

[0242] By the term “C₂₋₆-alkynyl” (including those which are part of other groups) are meant branched and unbranched alkynyl groups with 2 to 6 carbon atoms and by the term “C₂₋₄-alkynyl” are meant branched and unbranched alkynyl groups with 2 to 4 carbon atoms, provided that they have at least one triple bond. Alkynyl groups with 2 to 4 carbon

atoms are preferred. Examples include: ethynyl, propynyl, butynyl, pentynyl, or hexynyl. Unless stated otherwise, the definitions propynyl, butynyl, pentynyl, and hexynyl include all the possible isomeric forms of the groups in question. Thus, for example, propynyl includes 1-propynyl and 2-propynyl, butynyl includes 1-, 2-, and 3-butenyl, 1-methyl-1-propynyl, 1-methyl-2-propynyl etc.


[0243] By the term “C₂₋₆-alkynylene” (including those which are part of other groups) are meant branched and unbranched alkynylene groups with 2 to 6 carbon atoms and by the term “C₂₋₄-alkynylene” are meant branched and unbranched alkylene groups with 2 to 4 carbon atoms. Preferred are alkynylene groups with 2 to 4 carbon atoms. Examples include: ethynylene, propynylene, 1-methylethynylene, butynylene, 1-methylpropynylene, 1,1-dimethylethynylene, 1,2-dimethylethynylene, pentynylene, 1,1-dimethylpropynylene, 2,2-dimethylpropynylene, 1,2-dimethylpropynylene, 1,3-dimethylpropynylene, or hexynylene. Unless stated otherwise, the definitions propynylene, butynylene, pentynylene, and hexynylene include all the possible isomeric forms of the groups in question with the same number of carbons. Thus, for example, propynyl also includes 1-methylethynylene and butynylene includes 1-methylpropynylene, 1,1-dimethylethynylene, 1,2-dimethylethynylene.


[0244] By the term “aryl” (including those which are part of other groups) are meant aromatic ring systems with 6 or 10 carbon atoms. Examples include: phenyl or naphthyl, the preferred aryl group being phenyl. Unless otherwise stated, the aromatic groups may be substituted by one or more groups selected from among methyl, ethyl, isopropyl, tert-butyl, hydroxy, fluorine, chlorine, bromine, and iodine.

[0245] By the term “aryl-C₁₋₆-alkylene” (including those which are part of other groups) are meant branched and unbranched alkylene groups with 1 to 6 carbon atoms, which are substituted by an aromatic ring system with 6 or 10 carbon atoms. Examples include: benzyl, 1- or 2-phenylethyl, or 1- or 2-naphthylethyl. Unless otherwise stated, the aromatic groups may be substituted by one or more groups selected from among methyl, ethyl, isopropyl, tert-butyl, hydroxy, fluorine, chlorine, bromine, and iodine.


[0246] By the term “heteroaryl-C₁₋₆-alkylene” (including those which are part of other groups) are meant, even though they are already included under “aryl-C₁₋₆-alkylene”, branched and unbranched alkylene groups with 1 to 6 carbon atoms, which are substituted by a heteroaryl.

[0247] A heteroaryl of this kind includes five- or six-membered heterocyclic aromatic groups or 5-10-membered, bicyclic heteroaryl rings which may contain one, two, or three heteroatoms selected from among oxygen, sulfur, and nitrogen, and contain so many conjugated double bonds that an aromatic system is formed. The following are examples of five- or six-membered heterocyclic aromatic groups:






[0248] Unless otherwise stated, these heteroaryls may be substituted by one or more groups selected from among methyl, ethyl, isopropyl, tert-butyl, hydroxy, fluorine, chlorine, bromine, and iodine. The following are examples of heteroaryl- C_{1-6} -alkylenes:




[0249] By the term " C_{1-6} -haloalkyl" (including those which are part of other groups) are meant branched and unbranched alkyl groups with 1 to 6 carbon atoms, which are substituted by one or more halogen atoms. By the term " C_{1-4} -alkyl" are meant branched and unbranched alkyl groups with 1 to 4 carbon atoms, which are substituted by one or more halogen atoms. Alkyl groups with 1 to 4 carbon atoms are preferred. Examples include: CF_3 , CHF_2 , CH_2F , and CH_2CF_3 .

[0250] By the term " C_{3-7} -cycloalkyl" (including those which are part of other groups) are meant cyclic alkyl groups with 3 to 7 carbon atoms. Examples include: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl. Unless otherwise stated, the cyclic alkyl groups may be substituted by one or more groups selected from among methyl, ethyl, isopropyl, tert-butyl, hydroxy, fluorine, chlorine, bromine, and iodine.


[0251] By the term "heterocyclic rings" or also "heterocycles" are meant five-, six-, or seven-membered, saturated or unsaturated heterocyclic rings which may contain one, two, or three heteroatoms, selected from among oxygen, sulfur, and nitrogen, while the ring may be linked to the molecule through a carbon atom or through a nitrogen atom, if there is one. Although included by the term "heterocyclic rings" or "heterocycles", the term "heterocyclic non-aromatic rings" refers to five-, six-, or seven-membered unsaturated rings. Examples include:




[0252] Although included by the term "heterocyclic rings" or "heterocycles", the term "heterocyclic aromatic rings" or "heteroaryl" refers to five- or six-membered heterocyclic aromatic groups or 5-10-membered, bicyclic heteroaryl rings which may contain one, two, or three heteroatoms, selected from among oxygen, sulfur, and nitrogen, and contain so many conjugated double bonds that an aromatic system is formed. Examples of five- or six-membered heterocyclic aromatic groups include:




[0253] Unless otherwise mentioned, a heterocyclic ring (or "heterocycle") may be provided with a keto group. Examples include:





[0254] By the term "bicyclic rings" are meant eight-, nine-, or ten-membered bicyclic rings which may optionally contain one or more heteroatoms, selected from among oxygen, sulfur, and nitrogen. The ring may be linked to the molecule through a carbon atom of the ring or through a nitrogen atom of the ring, if there is one. Examples include:




[0255] Although included by the term “bicyclic rings”, the term “fused bicyclic rings” denotes bicyclic rings wherein the bridge separating the rings denotes a direct single bond. The following are examples of a fused, bicyclic ring:



[0256] Although included by the term “bicyclic rings”, the term “fused bicyclic heterorings” denotes is bicyclic 5-10 membered heterorings which contain one, two, or three heteroatoms, selected from among oxygen, sulfur, and nitrogen, and wherein the bridge separating the rings denotes a direct single bond. Examples include pyrrolizine, indole, indolizine, isoindole, indazole, purine, quinoline, isoquinoline, benzimidazole, benzofuran, benzopyran, benzothiazole, benzothiazole, benzoisothiazole, pyridopyrimidine, pteridine, pyrimidopyrimidine,



[0257] By the term “heterocyclic spirorings” (spiro) are meant 5-10 membered, spirocyclic rings which may optionally contain one, two, or three heteroatoms, selected from among oxygen, sulfur, and nitrogen, while the ring may be linked to the molecule through a carbon atom or through a nitrogen atom, if there is one. Unless otherwise mentioned, a spirocyclic ring may be provided with an oxo, methyl, or ethyl group. Examples include:



[0258] “Halogen” within the scope of the present invention denotes fluorine, chlorine, bromine, or iodine. Unless stated to the contrary, fluorine, chlorine, and bromine are regarded as preferred halogens.

[0259] By the terms “ambient temperature” or “RT” is meant normal room temperature.

[0260] Compounds of general formula 1 may have acid groups, mainly carboxyl groups, and/or basic groups such as, e.g., amino functions. Compounds of general formula 1 may therefore be present as internal salts, as salts with pharmaceutically useable inorganic acids such as, for example, hydrochloric acid, sulfuric acid, phosphoric acid, sulfonic acid, or organic acids (such as, for example, maleic acid, fumaric acid, citric acid, tartaric acid, or acetic acid) or as salts with pharmaceutically useable bases such as alkali or alkaline earth metal hydroxides or carbonates, zinc, or ammonium hydroxides or organic amines such as e.g., diethylamine, triethylamine, or triethanolamine, inter alia.

[0261] As mentioned hereinbefore, the compounds of formula 1 may be converted into the salts thereof, particularly for pharmaceutical use, into the physiologically and pharmacologically acceptable salts thereof. These salts may on the one hand be in the form of the physiologically and pharmacologically acceptable acid addition salts of the compounds of formula 1 with inorganic or organic acids. On the other hand, if R is hydrogen, the compound of formula 1 may also be converted by reaction with inorganic bases into physiologically and pharmacologically acceptable salts with alkali or alkaline earth metal cations as counter ion. The acid addition salts may be prepared, for example, using hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, acetic acid, fumaric acid, succinic acid, lactic acid, citric acid, tartaric acid, or maleic acid. It is also possible to use mixtures of the abovementioned acids. The alkali and alkaline earth metal salts of the

compound of formula I are preferably prepared using the alkali and alkaline earth metal hydroxides and hydrides thereof, of which the hydroxides and hydrides of the alkaline earth metals, particularly of sodium and potassium, are preferred and sodium and potassium hydroxide are particularly preferred.

[0262] If desired, the compounds of general formula (1) may be converted into the salts thereof, particularly, for pharmaceutical use, into the pharmacologically acceptable acid addition salts with an inorganic or organic acid. Suitable acids include, for example, succinic acid, hydrobromic acid, acetic acid, fumaric acid, maleic acid, methanesulfonic acid, lactic acid, phosphoric acid, hydrochloric acid, sulfuric acid, tartaric acid, or citric acid. It is also possible to use mixtures of the abovementioned acids.

[0263] The invention relates to the compounds in question, optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates, in the form of the tautomers as well as in the form of the free bases or the corresponding acid addition salts with pharmacologically acceptable acids such as, for example, acid addition


salts with hydrohalic acids, for example, hydrochloric or hydrobromic acid or organic acids such as, for example, oxalic, fumaric, diglycolic, or methanesulfonic acid.

[0264] The compounds according to the invention may optionally occur as racemates, but they may also be obtained as pure enantiomers, i.e., in the (R) or (S) form. Preferred compounds are those which occur as racemates or as the (S) form.

[0265] The invention relates to the compounds in question, optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates, in the form of the tautomers as well as in the form of the free bases or the corresponding acid addition salts with pharmacologically acceptable acids such as, for example, acid addition salts with hydrohalic acids, for example, hydrochloric or hydrobromic acid or organic acids, such as, for example, oxalic, fumaric, diglycolic, or methanesulfonic acid.







[0266] The Tables of Examples A and B that follow list compounds of formula 1 prepared according to the invention.

EXAMPLES A









#	R ¹	R ²	R ³	R ⁴	R ⁵	A
1.	H	H		H	Me	CH ₂
2.	H	H		H		CH ₂











-continued

3.	H	H		H		
4.	H	H		H	Me	CH ₂
5.	H	H		H	Me	CH ₂
6.	H	H		H	Me	CH ₂
7.	H	H		H	Me	CH ₂










-continued

8.	H	H		H	Me	CH ₂
9.	H	H		H	Me	CH ₂
10.	H	H		H		CH ₂
11.	H	H		H		CH ₂
12.	H	H		H		CH ₂











-continued

13. H	H		H		CH ₂
14. H	H		H		C=NH
15. H	H		H		CH ₂
16. H	H	 	H		CH ₂
17. H	H		H		CH ₂





-continued

18. H	H		H		CH ₂
19. H	H		H		CH ₂
20. H	H		H		CH ₂
21. H	H		H		CH ₂
22. Et	Et		H		CH ₂







-continued

23.	H	H		H		CH ₂
24.	H	H		H		CH ₂
25.	H	H		H		C=NH
26.	H	H		H		CH ₂
27.	H	H		H		CH ₂






-continued

28.	H	H		H	*	CH ₂
29.	H	H		H	*	CH ₂
30.	H	H		H	*	CH ₂
31.	H	H		H	*	CH ₂
32.	H	H H		H	*	CH ₂
33.	H	H H		H	Me	CH ₂






-continued

34.		H		H	Me	CH ₂
35.	H	H		H	Me	CH ₂
36.	H	H		H	Me	CH ₂
37.	H	H		H	Me	CH ₂
37a.	H	H		H	Me	CH ₂

-continued


37b.	H	H		H	Me	CH ₂
37c.	H	H		7-CF ₃	Me	CH ₂
37d.	H	H		H	cBu	CH ₂
37e.	H	H		H	cBu	CH ₂
37f.	H	H		H	cBu	CH ₂

-continued


37g.	H	H		H	CH ₂ CF ₃	CH ₂
37h.	H	H		H	CH ₂ CF ₃	CH ₂
37i.	H	H		H	CONHCH ₂ Ph	CH ₂
37j.	H	H		H	CONHCH ₂ Ph	CH ₂
37k.	H	H		H	CH ₂ CF ₃	CH ₂

-continued


37l. H H CH₂CF₃ CH₂




37m. H H cBu CH₂



37n. H H CH₂CF₃ CH₂




37o. H H cBu CH₂



-continued


EXAMPLES B



#	R ¹	R ²	R ³	R ⁴	A
38.	H	H		H	C=NH
39.	H	H		H	CH ₂
40.	H	H		H	
41.	H	H		H	CH ₂


-continued

42.	H	H		H	CH ₂
-----	---	---	--	---	-----------------




43.	H	H	H	H	CH ₂
-----	---	---	---	---	-----------------


44.	H	H		H	CH ₂
-----	---	---	--	---	-----------------




45.	H	H		H	CH ₂
-----	---	---	--	---	-----------------








46.	H	H		H	CH ₂
-----	---	---	--	---	-----------------








47.	H	H		H	CH ₂
-----	---	---	--	---	-----------------




-continued

48. H H H CH₂49. H H H CH₂50. H H H CH₂51. H H H CH₂52. H H H CH₂


-continued

53.	H	H		H	CH ₂
54.	H	H		H	CH ₂
55.	H	H		H	CH ₂
56.	H	H		H	CMe ₂
57.	H	H		H	CH ₂
58.	H	H		H	CH ₂

-continued

59. H H H H CH₂

60. H H



H



61. H H H

H

H




62. H H



H

CH₂


63. H H



H

CH₂

64. H H



H









65. H H

OH

H




-continued

66.	H	H		H	CH ₂
67.	H	H		H	
68.	H	H		H	CH ₂
69.	H	H		H	CH ₂
70.	H	H	CH ₂ CF ₃	H	CH ₂
71.	H	H		H	CH ₂
72.	H	H		H	CH ₂






-continued

EXAMPLES C








#	R ¹	R ²	R ³	R ⁴	A
73.	H	H		H	CH ₂
74.	H	H		H	CMe ₂
75.	H	H		H	CMe ₂
76.	H	H		H	CH ₂





-continued

77. H H H CMe₂78. H H H CMe₂79. H H H CH₂80. H H H CH₂81. H H H CH₂






-continued

82. H H H CH₂83. H H H CH₂84. H H H CH₂85. H H H CH₂86. H H H CH₂






-continued

87. H H H CH₂88. H H H CD₂89. H H H CD₂90. H H H CH₂91. H H H CH₂









-continued

92. H H H CH₂93. H H H CH₂94. H H H CH₂95. H H H CH₂96. H H H CH₂






-continued

97.	H	H	 <chem>Cc1ncsc1[C@H]2CCCCC2</chem>	H	CH ₂
98.	H	H	 <chem>CN1CC[C@H]1N(C(=O)OC)c2ccc(S(=O)(=O)C)cc2</chem>	H	CH ₂
99.	H	H	 <chem>CCN(C)C(=O)CN(C(=O)OC)c1cccc(CC)c1</chem>	H	CH ₂
100.	CHO	H	 <chem>CN(C)C(=O)CN(C(=O)OC)c1ccc(S(=O)(=O)C)cc1</chem>	H	CH ₂
104.	H	H	 <chem>CCCSc1ccc(cc1)CC</chem>	H	CH ₂
105.	H	H	 <chem>CCCSc1ccc(cc1)CC</chem>	H	CH ₂






-continued

106.	H	H		H	CH ₂
107.	H	H		H	CH ₂
108.	H	H		H	CH ₂
109.	H	H		H	CH ₂
110.	H	H		H	CH ₂
111.	H	H		H	CH ₂
112.	H	H		H	CH ₂
113.	H	H		H	CH ₂


-continued

114. H H H CH₂115. H H H CH₂116. H H H CH₂117. H H H CH₂118. H H H CH₂119. H H H CH₂

-continued

120. H H H CH₂121. H H H CH₂122. H H H CH₂123. H H H CH₂124. H H H CH₂

-continued

125. H H H CH₂

EXAMPLES D


Example 103

[0267] Other Examples according to the invention are as follows:

[0270]



Example 101

[0268]



Example 102


[0269]




[0271] General synthesis scheme 1 is shown below, according to which the compounds of formula 1 according to the invention (coumaranones and indolinones) may be prepared. Instead of the allyloxycarbonyl protecting group used in Scheme 1 to protect the amino function, it is also possible to use other protecting groups, preferably the tert-butyloxycarbonyl protecting group (BOC protecting group).


Synthesis Scheme 1

1. Coupling of the tBu-O protecting group to the substituted benzoic acid




2. Coupling of the alkyloxycarbonyl protecting group to intermediate B






3. Cleaving of the tBu-O protecting group




4. Coupling of the intermediate D to coumaranone or indolinone




-continued


5. Coupling of the intermediate E to an R³-substituted amine



6. Cleaving of the allyloxycarbonyl protecting group from intermediate F



[0272] The present invention further relates to all intermediate products of the process according to Scheme 1, particularly the intermediate products according to formulae i, ii, or iii



wherein X, A, R², and R³ have the meanings given above and PG (=protective group) is hydrogen or a suitable protecting group, preferably the allyloxycarbonyl protecting group or the tert-butyloxycarbonyl protecting group.


[0273] The following are illustrations of methods of synthesis for preparing compounds of formula 1 according to synthesis Scheme 1 above.

Example 40

1. 3-[[4-(1-aminocyclopropyl)phenyl]-[4-(1-methyl-1H-imidazol-4-yl)phenylamino]-methylene]-3H-benzofuran-2-one (End Product)

1.1 tert-butyl 4-(1-aminocyclopropyl)benzoate (Intermediate B)

[0274]




[0275] 50 mL of a 2M lithium isopropoxide solution in THF was placed under an argon atmosphere and 13.4 g of anhydrous lithium iodide was added thereto. After 15 minutes, 45 mL of a 1M methyltitanium isopropoxide solution in THF was added. Then the mixture was combined with 8.1 g of tert-butyl 4-cyanobenzoate and stirred for 10 minutes.

Then within 60 minutes, 47.3 mL of a 15% diethylzinc solution in hexane was added thereto and the mixture was stirred for 12 hours at RT. After hydrolysis with 20 mL of water, it was stirred for a further 30 minutes, filtered off, and washed 3× with 100 mL of ether. After extraction of the organic phase with water, the mixture was evaporated down with silica gel and the substance was chromatographed with dichloromethane/methanol (98:2) at 20 mL/min on silica gel 70. Residue: 3 g oil.

1.2 tert-butyl
4-(1-allyloxycarbonylamino)cyclopropyl)benzoate
(Intermediate C)


[0276]



[0277] 3 g of tert-butyl 4-(1-aminocyclopropyl)benzoate was placed in 30 mL of dichloromethane and 1.8 mL of pyridine. The solution was cooled to 0° C. and 1.2 mL of allyl chloroformate (dissolved in 5 mL of dichloromethane) was added dropwise at 0° C.-5° C. The mixture was left to react for 30 minutes at 0° C. and 2 hours at RT. The solution was combined with silica gel and evaporated down to the residue. This was chromatographed with dichloromethane on silica gel 70. Yield: 1.46 g of oil.

1.3 4-(1-allyloxycarbonylamino)cyclopropyl)benzoic acid

[0278]




[0279] A solution of 1.46 g of tert-butyl 4-(1-allyloxycarbonylamino)cyclopropyl)benzoate in 10 mL of acetonitrile was combined with 1 g of montmorillonite and refluxed for 3 hours. Then more montmorillonite was added and the mixture was boiled for 4 hours. The suspension was filtered hot, the inorganic material was extracted 3× with hot aceto-

nitrile and filtered off, and the acetonitrile solutions were combined and evaporated down. Residue: 0.8 g of crystals.

1.4 allyl(1-{4-[hydroxy-(2-oxobenzofuran-3-ylidene)methyl]phenyl}cyclopropyl)-carbamate (Intermediate D)


[0280]



[0281] 0.8 g of 4-(1-allyloxycarbonylamino)cyclopropylbenzoic acid and 0.585 mL of triethylamine were dissolved in 10 mL of anhydrous DMF and combined with 1.16 g of TBTU, stirred for 15 minutes at RT, then combined with 0.4 g of coumaranone and stirred for a further 10 minutes. While cooling with the ice bath, 0.42 g of NaH was added batchwise as a 60% suspension in white oil, the mixture was stirred for 2 hours at RT and, once the reaction had ended, the mixture was diluted with water to a total volume of approximately 150 mL. The mixture was acidified with 2N acetic acid. The crystals were suction filtered and washed with water. Yield: 1.1 g of solid with a melting point of 125° C.-126° C.

1.5 allyl(1-{4-[methoxy-(2-oxobenzofuran-3-ylidene)methyl]phenyl}cyclopropyl)-carbamate (Intermediate E)


[0282]



[0283] 1 g of allyl(1-{4-[hydroxy-(2-oxobenzofuran-3-ylidene)methyl]phenyl}cyclopropyl)-carbamate, 0.78 g of trimethyloxonium tetrafluoroborate (Meerwein salt), and 1 mL of diisopropylethylamine (Hünig base) were refluxed in 20 mL of dichloromethane for 2 hours, then another 1 mL of Hünig base and 0.5 g of Meerwein salt were added and the mixture was refluxed again for 2 hours. After cooling, the mixture was extracted 3× with water, and the organic phase was dried over MgSO₄ and evaporated down to the residue. Yield: 1.05 g of crude.

1.6 allyl(1-{4-[[4-(1-methyl-1H-imidazol-4-yl)phenylamino]-(2-oxobenzofuran-3-ylidene)methyl]phenyl}cyclopropyl)carbamate (Intermediate F)


[0284]



[0285] 1.05 g of allyl(1-{4-[methoxy-(2-oxobenzofuran-3-ylidene)methyl]phenyl}cyclopropyl)-carbamate and 0.47 g of 4-(4-aminophenyl)-1-methylimidazole in 3 mL of DMPU were reacted in the microwave reactor at 180° C. After cooling, the mixture was diluted to 150 mL with ethyl acetate, extracted 2× with water, and the organic phase was dried over MgSO₄ and evaporated down to the residue with silica gel. The substance was chromatographed with dichloromethane/methanol (97:3) on silica gel 70 at a flow rate of 30 mL/min. Residue: 0.6 g of yellow oil.

1.7 3-{[4-(1-aminocyclopropyl)phenyl]-[4-(1-methyl-1H-imidazol-4-yl)phenylamino]-methylene}-3H-benzofuran-2-one (End Product)

[0286]



[0287] 0.58 g of allyl(1-{4-[[4-(1-methyl-1H-imidazol-4-yl)phenylamino]-(2-oxobenzofuran-3-ylidene)methyl]phenyl}cyclopropyl)carbamate was dissolved in 20 mL of dichloromethane. The solution was freed from oxygen using argon. Then at RT, 0.52 mL of diethylamine and 0.058 g of tetrakis(triphenylphosphine)palladium (0) were added. The mixture was stirred for 2 hours at RT, the suspension was evaporated down with silica gel and chromatographed with

dichloromethane/methanol (97:3) on silica gel 70 at a flow rate of 30 mL/min. The clean fractions were combined, evaporated down to the residue, dissolved in dichloromethane, filtered, and combined with diisopropyl ether. The dichloromethane was distilled off, the crystals formed were suction filtered and washed with diisopropyl ether. Yield: 0.26 g of yellow crystals, melting point 190° C.-191° C., as base. R_f value: 0.51 (dichloromethane/methanol (9:1)).


[0288] The following compounds were prepared analogously to Synthesis Scheme 1 or to Example 40:

Example	salt form	melting point (° C.)
60	x HCl	>250
61	x HCl	185-187
64	x HCl	261-262
65	x HCl	210-211
67	x HCl	254-255
74	x HCl	285-286


-continued

Example	salt form	melting point (° C.)
77	x HCl	275-276
78	x HCl	214-216
37k	x HCl	263-265
37m	x HCl	>270
68	x HCl	286-287 decomp.
97	x HCl	234-235
102	base	196-197

[0289] Scheme 2 shows the method of preparation for synthesizing the compound according to Example 47. The majority of the compounds of formula 1 according to the invention (coumaranones and indolinones) may also be prepared analogously to this preparation method according to Scheme 2, particularly the compounds according to the invention mentioned on the following pages.




-continued



[0290] The present invention further relates to all the intermediate products of the process according to Scheme 2, particularly the intermediate products according to formulae I, II, III, IV, or V

-continued




[0291] The methods of synthesizing a number of example compounds are described hereinafter; they are substantially similar to Synthesis Scheme 2 described above.

Example 36

2. 3-[(4-aminomethylphenyl)-[4-(1-methyl-1H-imidazol-4-yl)phenylamino]methylene]-1-methyl-1,3-dihydroindol-2-one

[0292]



[0293] Starting from 4-cyanobenzoic acid, synthesis was carried out analogously to Example 40. Only the final step differed as follows. 6.15 g of 3-[(4-cyanophenyl)-[4-(1-methyl-1H-imidazol-4-yl)phenylamino]methylene]-1-methyl-1,3-dihydroindol-2-one was dissolved in 200 mL of methanolic ammonia and THF and hydrogenated with 5 g of Raney nickel at RT and under a pressure of 50 psi. After about 10 hours, the catalyst was removed by suction filtering and the filtrate was again filtered through a glass filter, evaporated down, and the crystalline residue was filtered with diisopropyl ether suction. The crystals were suspended in 400 mL of methanol, adjusted to pH 1 with 10% ethanolic hydrochloric acid, and heated to boiling temperature. The mixture was filtered and the filtrate was evaporated down to a residual volume of about 100 mL and combined with 300 mL of ethanol. After further evaporation to approximately 100 mL residual volume, the substance crystallized out. After cooling, it was suction filtered and washed with ice-cold ethanol. Yield: 6.15 yellow crystals; melting point >300° C.; R_f value=0.32 (dichloromethane/methanol/ammonia (85:15:1)).

[0294] The following were prepared analogously:

Ex-ample	salt form	melting point (° C.)	R_f value
1	base		
2	x TFA		0.38 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 9:1:0.1)
3	x TFA		0.5 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 4:1:0.1)
4	base		0.34 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 9:1:0.1)
5	x TFA		0.25 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 4:1:0.1)
6	base		0.27 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 9:1:0.1)
7	base		0.54 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 4:1:0.1)
8	base		0.33 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 9:1:0.1)
9	base		0.21 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 9:1:0.1)
10	x HCl	294 decomp.	

-continued

Ex-ample	salt form	melting point (° C.)	R_f value
11	base		0.27 (CH ₂ Cl ₂ /CH ₃ OH/NH ₃ 9:1:0.1)
12	x HCl	272	
13	x HCl	>250	
15	x HCl	278	
16	x HCl	214–215 decomp.	
17	x HCl	>245 decomp.	
18	base	225 decomp.	
19	x HCl	295–296	
20	x HCl	265–267	
21	x HCl	274–275	
22	x HCl	259–260 decomp.	
23	x HCl	281 decomp.	
24	x HCl	245–246 decomp.	
26	x HCl	>260 decomp.	
27	base	230–232	
28	base	153–155	
29	base	168–171	
30	base	233–238	
31	base	194–197	
32	x HCl	210–212	
33	base	157–159	
34	base	229	
35	base	228	
36	base	214–217	
37a	base	204–206	
37b	base	214–216	
37c	base	186–188	
37d	base	227–229	
37e	base	263–265	
37f	base	198–200	
37g	x HCl	255–256	
37h	x HCl	284	
37i	x HCl	251–252	
37j	x HCl	226–227	
37l	base	172–173	
37n	x HCl	195–198	
37o	x HCl	297–298	
39	x HCl	>290	
41	base	167–169	
42	base	201–203	
43	x HCl	95–97 decomp.	
44	base	222–224	
45	base	208–210	
46	base	186–188	
47	x HCl	274	
48	x HCl	186–188	
49	base	171–173	
50	base	164–166	
51	base	177–179	
52	x HCl	232–233	
53	x HCl	>285	
54	x HCl	245–247	
55	base	149–151	
57	base	161–164	
58	base	168–172	
59	base	153–156	
62	x HCl	285–286	
63	x HCl	289–290	
66	base	215–217	
69	base	204–205	
70	x HCl	285–286	
71	x HCl	>290	
72	x HCl	289	
73	base	163–166	
76	x HCl	288	
79	base	182–184	
80	base	135–137	
81	x HCl	294–295	
82	x HCl	254–255	
83	x HCl	>275	
84	x HCl	215–220	
85	base	135–137	

-continued

Ex- am- ple	salt form	melting point (° C.)	R _f value
86	x HCl	207	
87	x HCl	253–254	
88	x HCl	>290	
89	x HCl	>270	
90	x HCl	85–88 (amorphous?)	
91	x HCl	283–284	
92	x TFA	amorphous	
93	base	161–162	
94	x HCl	271–272	
95	x HCl	188	
96	x HCl	268–269	
98	x HCl	241 decomposition	
99	x HCl	196	
100	—	140	
101	x HCl	253–254	
103	x TFA	amorphous	

[0295]

Example	m + H	R _f value	retention time [min]
104		0.5	
105	380		
106		0.5	
107		0.4	
108		0.5	
109		0.2	
110		0.4	
111		0.5	
112		0.3	
113	352		
114		0.3	
115		0.6	
116		0.4	
117		0.4	
118		0.2	
119		0.3	
120		0.3	
121		0.3	
122			1.7
123		0.6	
124	539		
125		0.2	

Method of Carrying Out Thin Layer Chromatography to Determine the R_f Values:

[0296] The solid phase used was silica gel 60 F254 (made by Merck) and the liquid phase, unless otherwise specified, was a (9:1:0.1) mixture of dichloromethane:methanol:ammonia.

Chromolith Method:

[0297] HPLC-MS-1 Waters ZMD, Alliance 2690/2695 HPLC, Waters 2700 Autosampler, Waters 996/2996 diode array detector. The mobile phase used was:

A: water with 0.10% TFA

[0298] B: acetonitrile with 0.10% TFA

time in min	% A	% B	flow rate in mL/min
0.00	95	5	2.00
0.10	95	5	2.00
2.10	2	98	2.00
3.00	2	98	2.00
3.25	95	5	2.00


[0299] The stationary phase used was a Merck Chromolith™ SpeedROD RP-18e column, 4.6 mm×50 mm (column temperature: constant at 25° C.).

[0300] The diode array detection took place in a wavelength range of 210-400 nm.

Example 14

3, 4-[[4-(1-methyl-1H-imidazol-4-yl)phenylamino]-(2-oxo-1-phenyl-1,2-dihydroindol-3-ylidene)methyl]benzamide

[0301]



[0302] Starting from 4-cyanobenzoic acid synthesis was carried out analogously to Example 40. Only the last step differed as follows. 0.6 g of 3-[(4-cyanophenyl)-[4-(1-methyl-1H-imidazol-4-yl)phenylamino]methylene]-1-phenyl-1,3-dihydroindol-2-one was dissolved in 20 mL of dichloromethane and 30 mL of 40% ethanolic hydrochloric acid and stored in the refrigerator for 12 hours. The solution was evaporated down to the residue, combined with 50 mL of 6N ethanolic ammonia solution, and refluxed for 3 hours. The solution was evaporated down with silica gel and purified by chromatography with dichloromethane/methanol (8:2) on silica gel 70 (flow rate 20 mL/min). The clean fractions were evaporated down and the crystals with acetone, suction filtered, and washed. Yield: 0.2 g of yellow crystals; melting point 241° C. (decomp.).


[0303] Example 25 was prepared analogously (melting point 230° C.-232° C. as hydrochloride), as was Example 38 (melting point>275° C. with decomp. as hydrochloride).

Example 56

4. 3-[[4-(1-amino-1-methylethyl)phenyl]-[4-(1-methyl-1H-imidazol-4-yl)phenylamino]-methylene]-3H-benzofuran-2-one

4.1 1-[4-(4,4-dimethyl-4,5-dihydroxazol-2-yl)phenyl]-1-methylethylamine


[0304]



[0305] 130 g of anhydrous cerium (III) chloride was suspended in 1 L of THF and stirred for 1 hour, cooled to -60°C . with thorough mechanical stirring and combined dropwise with 330 mL of a 1.5M methyl lithium solution in THF/cumene, while the internal temperature did not rise above -50°C . The yellow suspension was stirred for 30 minutes and at -50°C . combined with a solution of 34.4 g of 2-(4-cyanophenyl)-4,4-dimethyl-4,5-dihydroxazole in 50 mL of THF. After another 12 hours, 330 mL of concentrated ammonia solution was added dropwise at -55°C ., whereupon a violet precipitate was formed. It was allowed to come up to RT, filtered, and the residue was washed with 1 L of THF. The organic phase was evaporated down. The oily yellow residue was purified by chromatography (dichloromethane/methanol (9:1), silica gel 70, flow rate 50 mL/min). The clean fractions yielded 33.3 g of yellow oil with an R_f value 0.20 (dichloromethane/methanol (7:3)).

4.2 tert-butyl {1-[4-(4,4-dimethyl-4,5-dihydroxazol-2-yl)phenyl]-1-methylethyl} carbamate


[0306]



[0307] A solution of 0.4 g of sodium carbonate in 4 mL of water was added to a solution of 0.8 g of 1-[4-(4,4-dimethyl-4,5-dihydroxazol-2-yl)phenyl]-1-methylethylamine in 15 mL of dioxane. Then within 10 minutes, a solution of 0.83 g of Boc-anhydride in dioxane was added dropwise. The yellow solution was stirred for 12 hours at RT. The dioxane was distilled off and the aqueous residue was extracted 2x with ethyl acetate. The organic phases were combined, dried over MgSO_4 , filtered, and evaporated down to the residue. After crystallization, 0.87 g of yellowish-white product was obtained.

4.3 4-(1-tert-butoxycarbonylamino-1-methylethyl)benzoic acid

[0308]




[0309] A solution of 0.83 g of tert-butyl {1-[4-(4,4-dimethyl-4,5-dihydroxazol-2-yl)phenyl]-1-methylethyl} carbamate and 3.1 mL of methyl iodide in 2 mL of DMF was stirred for 12 hours at 60°C . in a pressure tube reactor and, after cooling, evaporated down to the residue. The oil was stirred with 15 mL of 1M NaOH for 3 hours at RT. The suspension was extracted with ether. The aqueous phase was acidified to pH 2 with 1M HCl and the precipitate formed was taken up in ether. After the aqueous phase had been extracted twice more with ether, the organic phases were combined, dried over MgSO_4 , filtered, and the filtrate was evaporated down to the residue. Yield: 0.4 g of crystals; R_f value 0.41 (dichloromethane/methanol (9:1)).

[0310] The process used in Example 40 hereinbefore was used to cleave the Boc protecting group.

4.4 3-[[4-(1-amino-1-methylethyl)phenyl]-[4-(1-methyl-1H-imidazol-4-yl)phenylamino]methylene]-3H-benzofuran-2-one

[0311]




[0312] 2.5 g of tert-butyl(1-methyl-1-[4-[[4-(1-methyl-1H-imidazol-4-yl)phenylamino]-(2-oxobenzofuran-3-ylidene)methyl]phenyl]ethyl)carbamate (N-Boc derivative of Example 56) was dissolved in 50 mL of dioxane and combined with 50 mL of a 4N solution of hydrogen chloride in dioxane. The solution was stirred for 1 hour at RT and evaporated down. The residue was dissolved in 50 mL of methanol and evaporated down with another 150 mL of isopropanol almost to the residue and then crystallized. After


diluting with acetone, the product was suction filtered and washed with acetone. Yield: 1.9 g of yellow crystals of melting point $>280^{\circ}\text{C}$. as hydrochloride; R_f value: 0.18 (dichloromethane/methanol (8:2)).

[0313] The components of the synthesis described above are known in the literature, commercially available or prepared as follows:

Component 1




[0314] 26.3 g of 2-bromophenylacetic acid was dissolved in 200 mL of dichloromethane and combined with 15 mL of oxalyl chloride and one drop of DMF. After 1 hour at RT, the solution was evaporated down and the acid chloride was further used directly. The acid chloride was dissolved in 40 mL of dichloromethane and added dropwise to a solution of 9 g of cyclobutylamine and 27 mL of diisopropylethylamine in dichloromethane at 15°C .- 20°C . After 12 hours at RT, the dichloromethane was distilled off, the residue was dissolved in 600 mL of ethyl acetate and extracted 2x with 4N HCl, 2x with 4N NaOH, and 3x with water. The organic phase was dried over MgSO_4 , filtered, and evaporated down to the crystalline residue. After the addition of diisopropyl ether, the mixture was suction filtered and 24.1 g of 2-bromophenylacetic acid-cyclobutylamide was obtained. Melting point 159°C .- 161°C .; R_f value: 0.67 (dichloromethane/methanol (9:1)).




[0315] 400 mL of toluene was degassed with argon and then 13.4 g of 2-bromophenylacetic acid-cyclobutylamide, 9.7 g of powdered K_2CO_3 , 0.9 g of tris(dibenzylideneacetone)dipalladium (0) and 1.2 g of tri-*o*-tolylphosphine were added. After 72 hours stirring at 100°C . and cooling, the mixture was diluted to 2 L with diethyl ether, filtered to remove insoluble matter, and evaporated down to the residue. After crystallization in diisopropyl ether, the mixture was suction filtered and the filtrate was evaporated down to the residue with silica gel. The purification was carried out by chromatography on silica gel 60 (eluant: dichloromethane). Yield: 3.75 g of oil; R_f value: 0.18 (dichloromethane).


Component 2



[0316] 100 g of *p*-nitrophenacylbromide and 400 mL of formamide were stirred for 2 hours at 175°C . After cooling, the mixture was made alkaline with 20 mL of ammonia, stirred into 800 mL of water, and the precipitate was suction filtered. The crystals were recrystallized from methanol and suction filtered. Yield: 46.5 g; melting point 217°C .- 219°C .




[0317] 45 g of 1H-4-(4-nitrophenyl)imidazole was placed in 300 mL of DMSO and 30 g of potassium *tert*-butoxide was added batchwise while cooling with ice. The mixture was heated to RT and stirred for 1 hour. Then 16.5 mL of methyl iodide was added dropwise between 20°C .- 25°C . and stirred for a further 2 hours at RT. Then the mixture was poured onto 300 g of ice, stirred, the precipitate formed was suction filtered and washed thoroughly with water. Yield: 39.8 g.




[0318] 22 g of 1-methyl-4-(4-nitrophenyl)imidazole was dissolved in 1.6 L of methanol and combined with 4 g of Pd/C (10%). The mixture was hydrogenated at 50 psi and RT for approximately 5 hours. Then another gram of catalyst was added and the mixture was hydrogenated for 18 hours, then for a further 12 hours with the addition of another 2 g of catalyst. Finally, the catalyst was suction filtered and the filtrate was evaporated down. The residue was taken up in toluene and some methanol, and evaporated down somewhat until a dark-grey precipitate formed. This was suction filtered, taken up in methanol, combined with 4 g of activated

charcoal, and filtered off. The filtrate was combined with 100 mL of toluene, concentrated by rotary evaporation, and crystallized. Yield: 22 g; R_f value: 0.34 (dichloromethane/methanol (9:1)).


Component 3



[0319] 100 g of 3-nitroaniline was dissolved in 400 mL of pyridine and 57 mL of methanesulfonyl chloride was added dropwise while cooling gently with ice. After 12 hours at RT, the red reaction mixture was poured onto 1.2 L of ice, stirred, suction filtered, and the solid was suspended with copious amounts of water, washed, and dried. Yield: 148 g of solid with a melting point of 165° C.-166° C.; R_f value: 0.65 (dichloromethane/methanol (9:1)).




50 g of 3-methanesulfonylamino-4-nitrobenzene was placed in 200 mL of DMF, 31 g of potassium tert-butoxide was added and the reaction mixture was stirred for 1 hour at RT. After the addition of 28.5 g of 2-chloro-N,N-dimethylacetamide, the mixture was stirred for 12 hours at 60° C. After cooling to RT, water was added and the mixture was extracted 5x with ethyl acetate. The organic phases were washed with water, dried over $MgSO_4$, filtered, and the filtrate was evaporated down to the residue. The residue was crystallized from diisopropylether. Yield: 31.5 g.




31.5 g of 2-(N-methanesulfonyl-N-(3-nitrophenyl)amino)-N,N-dimethylacetamide was hydrogenated with 3 g of Pd/C at RT and 50 psi in ammoniacal methanol for 24 hours. Then the catalyst was suction filtered while warm and washed several times with ethyl acetate. The filtrate was evaporated down to the residue. Yield: 20 g of solid with a melting point of 143° C.-144° C.

Component 4




[0320] 9.25 g (47 mmol) of 4-aminophenylpyrrole hydrochloride is placed in 150 mL of methanol and hydrogenated with 1.00 g of Nishimura catalyst at ambient temperature under a pressure of 50 psi. Then the catalyst is filtered off and the filtrate is concentrated by evaporation. The residue is crystallized from acetonitrile and the enantiomers are obtained by chromatographic separation. Yield: 1.89 g (20%) of cis compound; NMR: LH201668; m.p.: 175° C.-177° C.

Component 5



[0321] 4.00 g (15 mmol) of Component 3 is placed in 60 mL of glacial acetic acid and hydrogenated with 0.600 g of Nishimura catalyst at ambient temperature under a pressure of 50 psi. Then the catalyst is filtered off and the filtrate is concentrated by evaporation. The residue is dissolved in a little water, treated with activated charcoal, and filtered. The filtrate is lyophilized. Yield: 4.90 g (84%); NMR: LH201565.

Component 6



2-methyl-2-(4-nitrophenyl)-[1,3]dioxolane: 33.03 g (200 mmol) of 4-nitroacetophenone, 12.30 mL (220 mmol) of ethylene glycol, and 1.00 g (5 mmol) of p-toluenesulfonic acid are placed in 250 mL of toluene, then refluxed for 16 hours using the water separator. Then the reaction mixture is cooled and extracted with water. The organic phase is dried and evaporated to dryness. The residue is extracted with diisopropyl ether and suction filtered. Yield: 34.45 g (82%).

[0322] 4-(2-methyl-[1,3]dioxolan-2-yl)cyclohexylamine: 8.37 g (40 mmol) of 2-methyl-2-(4-nitrophenyl)-[1,3]dioxolane is placed in 160 mL of methanol and hydrogenated with 1.00 g of Nishimura catalyst at ambient temperature under a pressure of 50 psi. Then the catalyst is filtered off and the filtrate is concentrated by evaporation. The residue is dissolved in cyclohexane, filtered to remove insoluble matter, and the filtrate is evaporated to dryness. Yield: 7.00 g (94%), cis/trans ratio 77:23; NMR: LG201616.

[0323] 9H-fluoren-9-ylmethyl[4-(2-methyl-[1,3]dioxolan-2-yl)cyclohexyl]carbamate: A solution of 12.72 g (120 mmol) of sodium carbonate is placed in 120 mL of water and combined with a solution of 7.00 g (38 mmol) of 4-(2-methyl-[1,3]dioxolan-2-yl)cyclohexylamine (cis/trans mixture) in 50 g of dioxane. After cooling to 0° C., a solution of 10.09 g (39 mmol) of Fmoc-chloride in 100 g of dioxane is added dropwise within 0.2 hours. The reaction mixture is stirred for 16 hours with the cooling removed. Then the mixture is poured onto water and extracted with ethyl acetate. The organic phase is washed with water, dried, and evaporated to dryness. The residue is purified by chromatography. Yield: 13.10 g (85%) cis/trans ratio 4:1; NMR: LH201618.

[0324] 9H-fluoren-9-ylmethyl(4-acetylcyclohexyl)carbamate: 13.10 g (32 mmol) of 9H-fluoren-9-ylmethyl[4-(2-methyl-[1,3]dioxolan-2-yl)cyclohexyl]carbamate (cis/trans mixture) and 1.30 g of p-toluenesulfonic acid are refluxed in 25 mL of water and 500 mL of acetone for 16 hours with stirring. Then the reaction mixture is concentrated by evaporation and the residue is dissolved in ethyl acetate and extracted with water. The organic phase is dried and evaporated to dryness. The residue is extracted with diisopropylether and suction filtered. Yield: 7.77 g (89%) of cis compound; NMR: LH201628.

[0325] 9H-fluoren-9-ylmethyl[4-(2-bromoacetyl)cyclohexyl]carbamate: 7.77 g (21 mmol) of 9H-fluoren-9-ylmethyl(4-acetylcyclohexyl)carbamate (cis compound) is dissolved in 100 mL of methanol at ambient temperature and combined with 1.09 mL (21 mmol) of bromine. The mixture is stirred for 16 hours at ambient temperature, cooled, and the crystals precipitated are suction filtered. Yield: 6.35 g (50%); NMR: LG201641.

[0326] 9H-fluoren-9-ylmethyl[4-(2-methylthiazol-4-yl)cyclohexyl]carbamate: 4.00 g (7 mmol) of 9H-fluoren-9-ylmethyl[4-(2-bromoacetyl)cyclohexyl]carbamate (cis compound) and 0.700 g (9 mmol) of thioacetamide are refluxed in 50 mL of acetonitrile for 72 hours with stirring. Then the reaction mixture is concentrated by evaporation, the residue is purified by chromatography. Corresponding fractions are combined and evaporated to dryness. The crystalline residue is extracted with diisopropyl ether and suction filtered. Yield: 1.10 g (39%) of cis compound; NMR: LH201648; m.p.: 140° C.-141° C.

[0327] 4-(2-methylthiazol-4-yl)cyclohexylamine (Component 6): 1.10 g (2 mmol) of 9H-fluoren-9-ylmethyl[4-(2-methylthiazol-4-yl)cyclohexyl]carbamate (cis compound) and 15 mL of diethylamine are stirred in 30 mL of THF for 16 hours at ambient temperature. Then the reaction mixture is concentrated by evaporation and the residue is combined with THF and concentrated again by evaporation. The residue is purified by chromatography. Yield: 0.13 g (36%).

Indications

[0328] As has been found, the compounds of formula 1 are characterized by their wide range of applications in the therapeutic field. Particular mention should be made of those applications for which the compounds according to the invention of formula 1 are preferably suited on account of their pharmaceutical efficacy as PDE4 inhibitors. Examples include respiratory or gastrointestinal diseases or complaints, inflammatory diseases of the joints, skin, or eyes, cancers, and also diseases of the peripheral or central nervous system.

[0329] Particular mention should be made of the prevention and treatment of diseases of the airways and of the lung which are accompanied by increased mucus production, inflammations, and/or obstructive diseases of the airways. Examples include acute, allergic, or chronic bronchitis, chronic obstructive bronchitis (COPD), coughing, pulmonary emphysema, allergic or non-allergic rhinitis or sinusitis, chronic rhinitis or sinusitis, asthma, alveolitis, Farmer's disease, hyperreactive airways, infectious bronchitis or pneumonitis, pediatric asthma, bronchiectases, pulmonary fibrosis, ARDS (acute adult respiratory distress syndrome), bronchial edema, pulmonary edema, bronchitis, pneumonia or interstitial pneumonia triggered by various causes, such as aspiration, inhalation of toxic gases, or bronchitis, pneumonia or interstitial pneumonia as a result of heart failure, irradiation, chemotherapy, cystic fibrosis, or mucoviscidosis, or alpha1-antitrypsin deficiency.

[0330] Also deserving special mention is the treatment of inflammatory diseases of the gastrointestinal tract. Examples include acute or chronic inflammatory changes in gall bladder inflammation, Crohn's disease, ulcerative colitis, inflammatory pseudopolyps, juvenile polyps, colitis cystica profunda, pneumatosis cystoides intestinales, diseases of the bile duct, and gall bladder, e.g., gallstones and conglomerates, for the treatment of inflammatory diseases of the joints such as rheumatoid arthritis or inflammatory diseases of the skin and eyes.

[0331] Preferential mention should also be made of the treatment of cancers. Examples include all forms of acute and chronic leukemias such as acute lymphatic and acute myeloid leukemia, chronic lymphatic and chronic myeloid leukemia, and bone tumors such as osteosarcoma and all types of glioma such as oligodendroglioma and glioblastoma.

[0332] Preferential mention should also be made of the prevention and treatment of diseases of the peripheral or central nervous system. Examples of these include depression, bipolar or manic depression, acute and chronic anxiety states, schizophrenia, Alzheimer's disease, Parkinson's disease, acute and chronic multiple sclerosis, or acute and chronic pain as well as injuries to the brain caused by stroke, hypoxia, or craniocerebral trauma.

[0333] Particularly preferably the present invention relates to the use of compounds of formula 1 for preparing a pharmaceutical composition for the treatment of inflammatory or obstructive diseases of the upper and lower respiratory tract including the lungs, such as, for example, allergic rhinitis, chronic rhinitis, bronchiectasis, cystic fibrosis, idiopathic pulmonary fibrosis, fibrosing alveolitis, COPD, chronic bronchitis, chronic sinusitis, asthma, Crohn's disease, ulcerative colitis, particularly COPD, chronic bronchitis, and asthma.

[0334] It is most preferable to use the compounds of formula 1 for the treatment of inflammatory and obstructive diseases such as COPD, chronic bronchitis, chronic sinusitis, asthma, Crohn's disease, ulcerative colitis, particularly COPD, chronic bronchitis, and asthma.

[0335] It is also preferable to use the compounds of formula 1 for the treatment of diseases of the peripheral or central nervous system such as depression, bipolar or manic depression, acute and chronic anxiety states, schizophrenia,

Alzheimer's disease, Parkinson's disease, acute and chronic multiple sclerosis, or acute and chronic pain as well as injuries to the brain caused by stroke, hypoxia, or cranio-cerebral trauma.

[0336] An outstanding aspect of the present invention is the reduced profile of side effects. This means, within the scope of the invention, being able to administer a dose of a pharmaceutical composition without inducing vomiting, preferably nausea and most preferably malaise in the patient. It is particularly preferable to be able to administer a therapeutically effective quantity of substance without inducing emesis or nausea, at every stage of the disease.

Combinations

[0337] The compounds of formula 1 may be used on their own or in conjunction with other active substances of formula 1 according to the invention. If desired the compounds of formula 1 may also be used in combination with other pharmacologically active substances. It is preferable to use for this purpose active substances selected, for example, from among betamimetics, anticholinergics, corticosteroids, other PDE4-inhibitors, LTD4-antagonists, EGFR-inhibitors, dopamine agonists, H1-antihistamines, PAF-antagonists, and PI3-kinase inhibitors or double or triple combinations thereof, such as, for example, combinations of

[0338] betamimetics with corticosteroids, PDE4-inhibitors, EGFR-inhibitors, or LTD4-antagonists,

[0339] anticholinergics with betamimetics, corticosteroids, PDE4-inhibitors, EGFR-inhibitors, or LTD4-antagonists,

[0340] corticosteroids with PDE4-inhibitors, EGFR-inhibitors, or LTD4-antagonists

[0341] PDE4-inhibitors with EGFR-inhibitors, or LTD4-antagonists

[0342] EGFR-inhibitors with LTD4-antagonists

[0343] MRP4-inhibitors.

[0344] The invention also encompasses combinations of three active substances, each selected from one of the abovementioned categories of compounds.

[0345] Suitable betamimetics used are preferably compounds selected from among albuterol, bambuterol, bitolterol, bixaterol, carbuterol, clenbuterol, fenoterol, formoterol, arformoterol, zinterol, hexoprenaline, ibuterol, isotharine, isoprenaline, levosalbutamol, mabuterol, meladrine, metaproterenol, orciprenaline, pirbuterol, procaterol, reproterol, rimiterol, ritodrine, salmeterol, salmefamol, soterol, sulfoneterol, tiaramide, terbutaline, tolubuterol, CHF-1035, HOKU-81, KUL-1248, 3-(4-{6-[2-hydroxy-2-(4-hydroxy-3-hydroxymethylphenyl)ethylamino]hexyloxy}butyl)benzylsulfonamide, 5-[2-(5.6-diethylindan-2-ylamino)-1-hydroxyethyl]-8-hydroxy-1H-quinolin-2-one, 4-hydroxy-7-[2-[[2-[[3-(2-phenylethoxy)propyl]sulfonyl]ethyl]amino]ethyl]-2(3H)benzothiazolone, 1-(2-fluoro-4-hydroxyphenyl)-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[3-(4-methoxybenzylamino)-4-hydroxyphenyl]-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-methoxyphenyl)-2-

methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-n-butylloxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-{4-[3-(4-methoxyphenyl)-1,2,4-triazol-3-yl]-2-methyl-2-butylamino}ethanol, 5-hydroxy-8-(1-hydroxy-2-isopropylaminobutyl)-2H-1,4-benzoxazin-3-(4H)-one, 1-(4-amino-3-chloro-5-trifluoromethylphenyl)-2-tert-butylamino]ethanol, 6-hydroxy-8-{1-hydroxy-2-[2-(4-methoxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-(ethyl 4-phenoxyacetate)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-phenoxyacetic acid)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 8-{2-[1,1-dimethyl-2-(2,4,6-trimethylphenyl)ethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-hydroxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-isopropylphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethylphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethoxyphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 4-(4-{2-[2-hydroxy-2-(6-hydroxy-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl)ethylamino]-2-methylpropyl}phenoxy)butyric acid, 8-{2-[2-(3,4-difluorophenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, and 1-(4-ethoxycarbonylamino-3-cyano-5-fluorophenyl)-2-(tert-butylamino)ethanol, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof.

[0346] Preferably the betamimetics are selected from among bambuterol, bitolterol, carbuterol, clenbuterol, fenoterol, formoterol, hexoprenaline, ibuterol, pirbuterol, procaterol, reproterol, salmeterol, sulfoneterol, terbutaline, tolubuterol, 3-(4-{6-[2-hydroxy-2-(4-hydroxy-3-hydroxymethylphenyl)ethylamino]hexyloxy}butyl)benzenesulfonamide, 5-[2-(5.6-diethylindan-2-ylamino)-1-hydroxyethyl]-8-hydroxy-1H-quinolin-2-one, 4-hydroxy-7-[2-[[2-[[3-(2-phenylethoxy)propyl]sulfonyl]ethyl]amino]ethyl]-2(3H)benzothiazolone, 1-(2-fluoro-4-hydroxyphenyl)-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[3-(4-methoxybenzylamino)-4-hydroxyphenyl]-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-N,N-dimethylaminophenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-methoxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-n-butylloxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[4-[3-(4-methoxyphenyl)-1,2,4-triazol-3-yl]-2-methyl-2-butylamino]ethanol, 5-hydroxy-8-(1-hydroxy-2-isopropylaminobutyl)-2H-1,4-benzoxazin-3-(4H)-one, 1-(4-amino-3-chloro-5-trifluoromethylphenyl)-2-tert-butylamino]ethanol, 6-hydroxy-8-{1-hydroxy-2-[2-(4-methoxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(ethyl 4-phenoxyacetate)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-phenoxyacetic acid)-1,1-dimethylethylamino]ethyl}-4H-

benzo[1,4]oxazin-3-one, 8-{2-[1,1-dimethyl-2-(2,4,6-trimethylphenyl)ethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-hydroxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-isopropylphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethylphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethoxyphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 4-(4-{2-[2-hydroxy-2-(6-hydroxy-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl)ethylamino]-2-methylpropyl}phenoxy)butyric acid, 8-{2-[2-(3,4-difluorophenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, and 1-(4-ethoxycarbonylamino-3-cyano-5-fluorophenyl)-2-(tert-butylamino)ethanol, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof.

[0347] Particularly preferred betamimetics are selected from among fenoterol, formoterol, salmeterol, 3-(4-{6-[2-hydroxy-2-(4-hydroxy-3-hydroxymethylphenyl)ethylamino]hexyloxy}butyl)benzenesulfonamide, 5-[2-(5,6-diethylindan-2-ylamino)-1-hydroxyethyl]-8-hydroxy-1H-quinoline-2-one, 1-[3-(4-methoxybenzylamino)-4-hydroxyphenyl]-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-N,N-dimethylaminophenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-methoxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-n-butylloxyphenyl)-2-methyl-2-propylamino]ethanol, 6-hydroxy-8-{1-hydroxy-2-[2-(4-methoxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(ethyl 4-phenoxyacetate)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-phenoxyacetic acid)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 8-{2-[1,1-dimethyl-2-(2,4,6-trimethylphenyl)ethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-hydroxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-isopropylphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethylphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethoxyphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 4-(4-{2-[2-hydroxy-2-(6-hydroxy-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl)ethylamino]-2-methylpropyl}phenoxy)butyric acid, 8-{2-[2-(3,4-difluorophenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, and 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[4-[3-(4-methoxyphenyl)-1,2,4-triazol-3-yl]-2-methyl-2-butylamino]ethanol, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof.

[0348] Of these betamimetics the particularly preferred ones according to the invention are formoterol, salmeterol, 3-(4-{6-[2-hydroxy-2-(4-hydroxy-3-hydroxymethylphenyl)ethylamino]hexyloxy}butyl)benzenesulfonamide, 6-hy-

droxy-8-{1-hydroxy-2-[2-(4-methoxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(ethyl 4-phenoxyacetate)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-phenoxyacetic acid)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 8-{2-[1,1-dimethyl-2-(2,4,6-trimethylphenyl)ethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-hydroxyphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 6-hydroxy-8-{1-hydroxy-2-[2-(4-isopropylphenyl)-1,1-dimethylethylamino]ethyl}-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethylphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 8-{2-[2-(4-ethoxyphenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, 4-(4-{2-[2-hydroxy-2-(6-hydroxy-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl)ethylamino]-2-methylpropyl}phenoxy)butyric acid, 8-{2-[2-(3,4-difluorophenyl)-1,1-dimethylethylamino]-1-hydroxyethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one, and 5-[2-(5,6-diethylindan-2-ylamino)-1-hydroxyethyl]-8-hydroxy-1H-quinoline-2-one, optionally in the form of the racemates, enantiomers, diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, or hydrates thereof.

[0349] According to the invention the acid addition salts of the betamimetics are preferably selected from among the hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethanesulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate, and hydro-p-toluenesulfonate, preferably the hydrochloride, hydrobromide, hydrosulfate, hydrophosphate, hydrofumarate, and hydromethanesulfonate. Of the abovementioned acid addition salts the salts of hydrochloric acid, methanesulfonic acid, benzoic acid, and acetic acid are particularly preferred according to the invention.

[0350] The anticholinergics used are preferably compounds selected from among the tiotropium salts, oxitropium salts, flutropium salts, ipratropium salts, glycopyrronium salts, trospium salts, tropenol 2,2-diphenylpropionate methobromide, scopine 2,2-diphenylpropionate methobromide, scopine 2-fluoro-2,2-diphenylacetate methobromide, tropenol 2-fluoro-2,2-diphenylacetate methobromide, tropenol 3,3',4,4'-tetrafluorobenzilate methobromide, scopine 3,3',4,4'-tetrafluorobenzilate methobromide, tropenol 4,4'-difluorobenzilate methobromide, scopine 4,4'-difluorobenzilate methobromide, tropenol 3,3'-difluorobenzilate methobromide, scopine 3,3'-difluorobenzilate methobromide, tropenol 9-hydroxyfluorene-9-carboxylate methobromide, tropenol 9-fluorofluorene-9-carboxylate methobromide, scopine 9-hydroxyfluorene-9-carboxylate methobromide, scopine 9-fluorofluorene-9-carboxylate methobromide, tropenol 9-methylfluorene-9-carboxylate methobromide, scopine 9-methylfluorene-9-carboxylate methobromide, cyclopropyltropine benzilate methobromide, cyclopropyltropine 2,2-diphenylpropionate methobromide, cyclopropyltropine 9-hydroxyxanthene-9-carboxylate methobromide, cyclopropyltropine 9-methylfluorene-9-carboxylate methobromide, cyclopropyltropine 9-methylxanthene-9-carboxylate methobromide, cyclopropyltropine 9-hydroxyfluorene-9-carboxylate methobromide, methylcyclopropyltropine 4,4'-difluorobenzilate methobromide, tropenol 9-hydroxyxanthene-9-carboxylate methobromide,

scopine 9-hydroxyxanthene-9-carboxylate methobromide, tropenol 9-methylxanthene-9-carboxylate methobromide, scopine 9-methylxanthene-9-carboxylate methobromide, tropenol 9-ethylxanthene-9-carboxylate methobromide, tropenol 9-difluoromethylxanthene-9-carboxylate methobromide, scopine 9-hydroxymethylxanthene-9-carboxylate methobromide, optionally in the form of the solvates, or hydrates thereof.

[0351] In the abovementioned salts the cations tiotropium, oxitropium, flutropium, ipratropium, glycopyrronium, and tropium are the pharmacologically active ingredients. As anions, the abovementioned salts may preferably contain chloride, bromide, iodide, sulfate, phosphate, methanesulfonate, nitrate, maleate, acetate, citrate, fumarate, tartrate, oxalate, succinate, benzoate, or p-toluenesulfonate, while chloride, bromide, iodide, sulfate, methanesulfonate, or p-toluenesulfonate are preferred as counter-ions. Of all the salts, the chlorides, bromides, iodides, and methanesulfonate are particularly preferred.

[0352] Of particular importance is tiotropium bromide. In the case of tiotropium bromide the pharmaceutical combinations according to the invention preferably contain it in the form of the crystalline tiotropium bromide monohydrate, which is known from WO 02/30928. If the tiotropium bromide is used in anhydrous form in the pharmaceutical combinations according to the invention, it is preferable to use anhydrous crystalline tiotropium bromide, which is known from WO 03/000265.

[0353] Corticosteroids used here are preferably compounds selected from among prednisolone, prednisone, butixocortpropionate, flunisolide, beclomethasone, triamcinolone, budesonide, fluticasone, mometasone, ciclesonide, rofleponide, dexamethasone, betamethasone, deflazacort, RPR-106541, NS-126, (S)-fluoromethyl 6,9-difluoro-17-[(2-furanylcarbonyloxy)-11-hydroxy-16-methyl-3-oxoandrosta-1,4-diene-17-carbothionate, and (S)-(2-oxotetrahydrofuran-3S-yl) 6,9-difluoro-11-hydroxy-16-methyl-3-oxo-17-propionyloxyandrosta-1,4-diene-17-carbothionate, optionally in the form of the racemates, enantiomers, or diastereomers thereof, and optionally in the form of the salts and derivatives, solvates, and/or hydrates thereof.

[0354] Particularly preferred is the steroid selected from among flunisolide, beclomethasone, triamcinolone, budesonide, fluticasone, mometasone, ciclesonide, rofleponide, dexamethasone, NS-126, (S)-fluoromethyl 6,9-difluoro-17-[(2-furanylcarbonyloxy)-11-hydroxy-16-methyl-3-oxoandrosta-1,4-diene-17-carbothionate, and (S)-(2-oxotetrahydrofuran-3S-yl) 6,9-difluoro-11-hydroxy-16-methyl-3-oxo-17-propionyloxyandrosta-1,4-diene-17-carbothionate, optionally in the form of the racemates, enantiomers, or diastereomers thereof, and optionally in the form of the salts and derivatives, solvates, and/or hydrates thereof.

[0355] Particularly preferred is the steroid selected from among budesonide, fluticasone, mometasone, ciclesonide, and (S)-fluoromethyl 6,9-difluoro-17-[(2-furanylcarbonyloxy)-11-hydroxy-16-methyl-3-oxoandrosta-1,4-diene-17-carbothionate, optionally in the form of the racemates, enantiomers, or diastereomers thereof, and optionally in the form of the salts and derivatives, solvates, and/or hydrates thereof.

[0356] Any reference to steroids includes a reference to any salts or derivatives, hydrates, or solvates thereof which

may exist. Examples of possible salts and derivatives of the steroids may be: alkali metal salts, such as, for example, sodium or potassium salts, sulfobenzoates, phosphates, isonicotinates, acetates, propionates, dihydrogen phosphates, palmitates, pivalates, or furoates thereof.

[0357] Other PDE4 inhibitors which may be used are preferably compounds selected from among enprofyllin, theophyllin, roflumilast, ariflo (cilomilast), tofomilast, pumafentrin, lirimilast, arofyllin, atizoram, D-4396 (Sch-351591), AWD-12-281 (GW-842470), NCS-613, CDP-840, D-4418, PD-168787, T-440, T-2585, V-11294A, C1-1018, CDC-801, CDC-3052, D-22888, YM-58997, Z-15370, N-(3,5-dichloro-1-oxopyridin-4-yl)-4-difluoromethoxy-3-cyclopropylmethoxybenzamide, (-)-p-[(4aR*, 10bS*)-9-ethoxy-1,2,3,4,4a,10b-hexahydro-8-methoxy-2-methylbenzo[s][1,6]naphthyridin-6-yl]-N,N-diisopropylbenzamide, (R)-(+)-1-(4-bromobenzyl)-4-[(3-cyclopentyl-4-methoxyphenyl)-2-pyrrolidone, 3-(cyclopentyl-4-methoxyphenyl)-1-(4-N'-[N-2-cyano-5-methylisothioureido]benzyl)-2-pyrrolidone, cis[4-cyano-4-(3-cyclopentyl-4-methoxyphenyl)cyclohexane-1-carboxylic acid], 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexane-1-one, cis[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol], (R)-(+)-ethyl[4-(3-cyclopentyl-4-methoxyphenyl)pyrrolidin-2-ylidene]acetate, (S)-(-)-ethyl[4-(3-cyclopentyl-4-methoxyphenyl)pyrrolidin-2-ylidene]acetate, 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(2-thienyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine, and 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(tert-butyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine, optionally in the form of the racemates, enantiomers, or diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, and/or hydrates thereof.

[0358] Particularly preferably the PDE4-inhibitor is selected from among enprofyllin, roflumilast, ariflo (cilomilast), arofyllin, atizoram, AWD-12-281 (GW-842470), T-440, T-2585, PD-168787, V-11294A, C₁₋₁₀₁₈, CDC-801, D-22888, YM-58997, Z-15370, N-(3,5-dichloro-1-oxopyridin-4-yl)-4-difluoromethoxy-3-cyclopropylmethoxybenzamide, cis[4-cyano-4-(3-cyclopentyl-4-methoxyphenyl)cyclohexane-1-carboxylic acid], 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-one, cis[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol], 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(2-thienyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine, and 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(tert-butyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine, optionally in the form of the racemates, enantiomers, or diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, and/or hydrates thereof.

[0359] Particularly preferably the PDE4-inhibitor is selected from among roflumilast, ariflo (cilomilast), arofyllin, AWD-12-281 (GW-842470), 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-one, cis[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol], atizoram, Z-15370, 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(2-thienyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine, and 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(tert-butyl)-9H-pyrazolo[3,

4-c]-1,2,4-triazolo[4,3-a]pyridine, optionally in the form of the racemates, enantiomers, or diastereomers, and optionally in the form of the pharmacologically acceptable acid addition salts, solvates, and/or hydrates thereof.

[0360] By acid addition salts with pharmacologically acceptable acids which the abovementioned PDE4-inhibitors might be in a position to form are meant, for example, salts selected from among the hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethanesulfonate, hydronitrate, hydromaleate, hydroacetate, hydrobenzoate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate, and hydro-p-toluenesulfonate, preferably hydrochloride, hydrobromide, hydrosulfate, hydrophosphate, hydrofumarate, and hydromethanesulfonate.

[0361] LTD4-antagonists which may be used are preferably compounds selected from among montelukast, pranlukast, zafirlukast, MCC-847 (ZD-3523), MN-001, MEN-91507 (LM-1507), VUF-5078, VUF-K-8707, L-733321, 1-(((R)-(3-(2-(6,7-difluoro-2-quinolinyl)ethenyl)phenyl)-3-(2-(2-hydroxy-2-propyl)phenyl)thio)methylcyclopropane acetic acid, 1-(((1(R)-3(3-(2-(2,3-dichlorothieno[3,2-b]pyridin-5-yl)-(E)-ethenyl)phenyl)-3-(2-(1-hydroxy-1-methyl-ethyl)phenyl)propyl)thio)methyl)cyclopropane acetic acid, and [2-[[2-(4-tert-butyl-2-thiazolyl)-5-benzofuranyl]oxymethyl]phenyl]acetic acid, optionally in the form of the racemates, enantiomers, or diastereomers, optionally in the form of the pharmacologically acceptable acid addition salts, and optionally in the form of the salts and derivatives, solvates, and/or hydrates thereof.

[0362] Preferably the LTD4-antagonist is selected from among montelukast, pranlukast, zafirlukast, MCC-847 (ZD-3523), MN-001, MEN-91507 (LM-1507), VUF-5078, VUF-K-8707, and L-733321, optionally in the form of the racemates, enantiomers, or diastereomers, optionally in the form of the pharmacologically acceptable acid addition salts, and optionally in the form of the salts and derivatives, solvates, and/or hydrates thereof.

[0363] Particularly preferably the LTD4-antagonist is selected from among montelukast, pranlukast, zafirlukast, MCC-847 (ZD-3523), MN-001, and MEN-91507 (LM-1507), optionally in the form of the racemates, enantiomers, or diastereomers, optionally in the form of the pharmacologically acceptable acid addition salts, and optionally in the form of the salts and derivatives, solvates, and/or hydrates thereof.

[0364] By acid addition salts with pharmacologically acceptable acids which the LTD4-antagonists may be capable of forming are meant, for example, salts selected from among the hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethanesulfonate, hydronitrate, hydromaleate, hydroacetate, hydrobenzoate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate, and hydro-p-toluenesulfonate, preferably hydrochloride, hydrobromide, hydrosulfate, hydrophosphate, hydrofumarate, and hydromethanesulfonate. By salts or derivatives which the LTD4-antagonists may be capable of forming are meant, for example: alkali metal salts, such as, for example, sodium or potassium salts, alkaline earth metal salts, sulfobenzoates, phosphates, isonicotinates, acetates, propionates, dihydrogen phosphates, palmitates, pivalates, or furoates.

[0365] The EGFR-inhibitors used are preferably compounds selected from among 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-s ((R)-6-methyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-((R)-6-methyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-[(S)-(tetrahydrofuran-3-yl)oxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-((R)-2-methoxymethyl-6-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[2-((S)-6-methyl-2-oxomorpholin-4-yl)ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(N,N-bis-(2-methoxyethyl)amino)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-[N-(2-methoxyethyl)-N-ethylamino]-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-((R)-tetrahydrofuran-3-yloxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-((S)-tetrahydrofuran-3-yloxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N-cyclopropyl-N-methylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-[(R)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-ethynylphenyl)amino]-6,7-bis-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-7-[3-(morpholin-4-yl)propyloxy]-6-[(vinylcarbonyl)amino]quinazoline, 4-[(R)-(1-phenylethyl)amino]-6-(4-hydroxyphenyl)-7H-pyrrolo[2,3-d]pyrimidine, 3-cyano-4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-ethoxyquinoline, 4-[[3-chloro-4-(3-fluorobenzoyloxy)phenyl]amino]-6-(5-[[2-(methanesulfonyl)ethyl]amino]methyl)furan-2-

yl)quinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-((R)-6-methyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-[(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(4-(5,5-dimethyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl)amino]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-[(R)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-7-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-6-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-[4-(2-oxomorpholin-4-yl)piperidin-1-yl]ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-(tert-butyloxycarbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-aminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-methanesulfonylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-3-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(morpholin-4-yl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(methoxymethyl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(piperidin-3-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-(2-acetylaminoethyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-ethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(S)-tetrahydrofuran-3-yloxy)-7-hydroxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{trans-4-[(dimethylamino)sulfonylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{trans-4-[(morpholin-4-yl)carbonylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{trans-4-[(morpholin-4-yl)sulfonylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-acetylaminoethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methanesulfonylamino)ethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(piperidin-1-yl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-aminocarbonylmethylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-(tetrahydropyran-4-yl)carbonyl]-N-methylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-(morpholin-4-yl)carbonyl]-N-methylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-(morpholin-4-yl)sulfonyl]-N-methylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-ethansulfonylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-ethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-(2-methoxyacetyl)piperidin-4-yloxy]-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-acetylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-[1-(tert-butyloxycarbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-(piperidin-1-yl)carbonyl]-N-methylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-(4-methylpiperazin-1-yl)carbonyl]-N-methylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-[(morpholin-4-yl)carbonylamino]cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-[2-(2-oxopyrrolidin-1-yl)ethyl]piperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(morpholin-4-yl)carbonyl)piperidin-4-yloxy]-7-(2-methoxyethoxy)quinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-acetylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-isopropylloxycarbonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-methylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-[N-(2-methoxyacetyl)-N-methylamino]cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(piperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-[1-(2-methoxyacetyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-[(1-(morpholin-4-yl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(cis-2,6-dimethylmorpholin-4-yl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(2-methylmorpholin-4-yl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(S,S)-(2-oxa-5-azabicyclo[2,2,1]hept-5-yl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(N-methyl-N-2-methoxyethylamino)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-ethylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(2-methoxyethyl)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(3-methoxypropylamino)carbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-methanesulfonyl)-N-methylamino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-acetyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline,

4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-methylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[trans-4-(N-methanesulfonyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-dimethylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-[N-[(morpholin-4-yl)carbonyl]-N-methylamino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-cyanopiperidin-4-yloxy)-7-methoxyquinazoline, cetuximab, trastuzumab, ABX-EGF, and Mab ICR-62, optionally in the form of the racemates, enantiomers, or diastereomers thereof, optionally in the form of the pharmacologically acceptable acid addition salts thereof, the solvates, and/or hydrates thereof.

[0366] Preferred EGFR-inhibitors are selected from among 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-diethylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(R)-6-methyl-2-oxomorpholin-4-yl]-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(R)-6-methyl-2-oxomorpholin-4-yl]-1-oxo-2-buten-1-yl]amino]-7-[(S)-(tetrahydrofuran-3-yl)oxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(R)-2-methoxymethyl-6-oxomorpholin-4-yl]-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-((S)-6-methyl-2-oxomorpholin-4-yl)ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-((4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl)amino)-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(N,N-bis-(2-methoxyethyl)amino)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-((4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl)amino)-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(N-(2-methoxyethyl)-N-methylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(N-(2-methoxyethyl)-N-methylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-((R)-tetrahydrofuran-3-yloxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-(((S)-tetrahydrofuran-3-yloxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl]amino]-7-

cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N-cyclopropyl-N-methylamino)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-[(R)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-ethynylphenyl)amino]-6,7-bis-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-7-[3-(morpholin-4-yl)propyloxy]-6-[(vinyl-carbonyl)amino]quinazoline, 4-[(R)-(1-phenylethyl)amino]-6-(4-hydroxyphenyl)-7H-pyrrolo[2,3-d]pyrimidine, 3-cyano-4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-ethoxyquinoline, 4-[[3-chloro-4-(3-fluorobenzoyloxy)phenyl]amino]-6-(5-[[2-(methanesulfonyl)ethyl]amino]methyl)furanyl)quinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-((R)-6-methyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-[(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-bis-(2-methoxyethyl)amino)-1-oxo-2-buten-1-yl]amino]-7-[(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-ethynylphenyl)amino]-6-[[4-(5,5-dimethyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-[(R)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-7-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-6-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2-oxomorpholin-4-yl)piperidin-1-yl]ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-(tert-butylloxycarbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-aminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-methanesulfonylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-3-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-((4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl)amino)-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(morpholin-4-yl)carbonyl]piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(methoxymethyl)carbonyl]piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(piperidin-3-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[(1-(2-acetylaminoethyl)piperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-ethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-((S)-tetrahydrofuran-3-yloxy)-7-hydroxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[trans-4-[(dimethylamino)sulfonylamino]cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[trans-4-[(morpholin-4-yl)carbonylamino]cyclohexan-1-

yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{trans-4-[(morpholin-4-yl)sulfonylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-acetylaminoethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methanesulfonylaminoethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(piperidin-1-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-aminocarbonylmethylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-[(tetrahydropyran-4-yl)carbonyl]-N-methylamino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-[(morpholin-4-yl)carbonyl]-N-methylamino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-[(morpholin-4-yl)sulfonyl]-N-methylamino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-ethansulfonylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-ethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-(2-methoxyacetyl)piperidin-4-yloxy]-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-acetylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-[1-(tert-butylloxycarbonyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-[(piperidin-1-yl)carbonyl]-N-methyl amino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-[N-[(4-methylpiperazin-1-yl)carbonyl]-N-methylamino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{cis-4-[(morpholin-4-yl)carbonylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[2-(2-oxopyrrolidin-1-yl)ethyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]piperidin-4-yloxy}-7-(2-methoxyethoxy)quinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-acetylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-isopropylloxycarbonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-methylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{cis-4-[N-(2-methoxyacetyl)-N-methylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(piperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-[1-(2-methoxyacetyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(cis-2,6-dimethylmorpholin-4-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline,

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(2-methylmorpholin-4-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(S,S)-(2-oxa-5-azabicyclo[2,2,1]hept-5-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-2-methoxyethylamino)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-ethylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(2-methoxyethyl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(3-methoxypropylamino)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-methanesulfonyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-acetyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-methylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[trans-4-(N-methanesulfonyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-dimethylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-[N-[(morpholin-4-yl)carbonyl]-N-methylamino]cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-cyanopiperidin-4-yloxy)-7-methoxyquinazoline, and cetuximab, optionally in the form of the racemates, enantiomers, or diastereomers thereof, optionally in the form of the pharmacologically acceptable acid addition salts thereof, the solvates, and/or hydrates thereof.

[0367] It is particularly preferable within the scope of the present invention to use those EGFR-inhibitors which are selected from among 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-((R)-6-methyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-[(S)-(tetrahydrofuran-3-yl)oxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-((S)-6-methyl-2-oxomorpholin-4-yl)ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-([4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl]amino)-7-cyclopropylmethoxyquinazoline, 4-[(R)-(1-phenylethyl)amino]-6-([4-[N-(tetrahydropyran-4-yl)-N-methylamino]-1-oxo-2-buten-1-yl]amino)-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-([4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl]amino)-7-cyclopentylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-([4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino)-7-[(R)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-ethynylphenyl)amino]-6,7-bis-(2-methoxyethoxy)quinazoline, 4-[(R)-(1-phenylethyl)amino]-6-(4-hydroxyphenyl)-7H-pyrrolo[2,3-d]pyrimidine, 3-cyano-4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-ethoxyquinoline, 4-[(R)-(1-phenylethyl)amino]-6-[[4-((R)-

6-methyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl] amino}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl] amino]-7-[(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-ethynylphenyl)amino]-6-[[4-(5,5-dimethyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-[4-(2-oxomorpholin-4-yl)piperidin-1-yl]ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-aminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-methanesulfonylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-3-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(piperidin-3-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-(2-acetylaminoethyl)piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-ethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{trans-4-[(morpholin-4-yl)carbonylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(piperidin-1-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-{N-[(morpholin-4-yl)carbonyl]-N-methylamino}cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-ethansulfonylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-(2-methoxyacetyl)piperidin-4-yloxy]-7-(2-methoxyethoxy)quinazoline, 4-[(3-ethynylphenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(cis-4-{N-[(piperidin-1-yl)carbonyl]-N-methylamino}cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(morpholin-4-yl)carbonylamino]cyclohexan-1-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[2-(2-oxopyrrolidin-1-yl)ethyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-acetyl)piperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-ethynylphenyl)amino]-6-[1-(morpholin-4-yl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-2-methoxyethylamino)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-ethylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-methanesulfonyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-acetyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-methylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[trans-4-(N-methanesulfonyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline,

4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-dimethylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-{N-[(morpholin-4-yl)carbonyl]-N-methylamino}cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-cyanopiperidin-4-yloxy)-7-methoxyquinazoline, and 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(2-methoxyethyl)carbonyl]piperidin-4-yloxy}-7-methoxyquinazoline, optionally in the form of the racemates, enantiomers, or diastereomers thereof, optionally in the form of the pharmacologically acceptable acid addition salts thereof, the solvates, and/or hydrates thereof.

[0368] Particularly preferred EGFR-inhibitors according to the invention are the compounds selected from among 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-cyclopropylmethoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-((R)-6-methyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-[(S)-(tetrahydrofuran-3-yl)oxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-((S)-6-methyl-2-oxomorpholin-4-yl)ethoxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-({4-[N-(2-methoxyethyl)-N-methylamino]-1-oxo-2-buten-1-yl}amino)-7-cyclopropylmethoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6,7-bis-(2-methoxyethoxy)quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino]-7-[(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-ethynylphenyl)amino]-6-[[4-(5,5-dimethyl-2-oxomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-methanesulfonylamino)cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(tetrahydropyran-3-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-[(morpholin-4-yl)carbonyl]piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-[2-(2-oxopyrrolidin-1-yl)ethyl]piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-acetyl)piperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, 4-[(3-ethynylphenyl)amino]-6-[(1-methylpiperidin-4-yloxy)-7-(2-methoxyethoxy)quinazoline, 4-[(3-ethynylphenyl)amino]-6-[1-(morpholin-4-yl)carbonyl]piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[1-[(2-methoxyethyl)carbonyl]piperidin-4-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-methanesulfonyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[cis-4-(N-acetyl-N-methylamino)cyclohexan-1-yloxy]-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-dimethylaminocyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(trans-4-{N-[(morpholin-4-yl)carbonyl]-N-

methylamino}cyclohexan-1-yloxy)-7-methoxyquinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxomorpholin-4-yl)ethoxy]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]quinazoline, 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-methanesulfonylpiperidin-4-yloxy)-7-methoxyquinazoline, and 4-[(3-chloro-4-fluorophenyl)amino]-6-(1-cyanopiperidin-4-yloxy)-7-methoxyquinazoline, optionally in the form of the racemates, enantiomers, or diastereomers thereof, optionally in the form of the pharmacologically acceptable acid addition salts thereof, the solvates, and/or hydrates thereof.

[0369] By acid addition salts with pharmacologically acceptable acids which the EGFR-inhibitors may be capable of forming are meant, for example, salts selected from among the hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethanesulfonate, hydronitrate, hydromaleate, hydroacetate, hydrobenzoate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate, and hydro-p-toluene-sulfonate, preferably hydrochloride, hydrobromide, hydrosulfate, hydrophosphate, hydrofumarate, and hydromethanesulfonate.

[0370] Examples of dopamine agonists which may be used preferably include compounds selected from among bromocriptine, cabergoline, alpha-dihydroergocryptine, lisuride, pergolide, pramipexole, roxindole, ropinirole, talipexole, terguride, and viozan. Any reference to the above-mentioned dopamine agonists within the scope of the present invention includes a reference to any pharmacologically acceptable acid addition salts and optionally hydrates thereof which may exist. By the physiologically acceptable acid addition salts which may be formed by the above-mentioned dopamine agonists are meant, for example, pharmaceutically acceptable salts which are selected from the salts of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, acetic acid, fumaric acid, succinic acid, lactic acid, citric acid, tartaric acid, and maleic acid.

[0371] Examples of H1-antihistamines preferably include compounds selected from among epinastine, cetirizine, azelastine, fexofenadine, levocabastine, loratadine, mizolastine, ketotifen, emedastine, dimethindene, clemastine, bami-pine, dexchlorpheniramine, pheniramine, doxylamine, chlorphenoxamine, dimenhydrinate, diphenhydramine, promethazine, ebastine, desloratidine, and meclizine. Any reference to the above-mentioned H1-antihistamines within the scope of the present invention includes a reference to any pharmacologically acceptable acid addition salts which may exist.

[0372] Examples of PAF-antagonists preferably include compounds selected from 4-(2-chlorophenyl)-9-methyl-2-[3(4-morpholinyl)-3-propanon-1-yl]-6H-thieno-[3,2-f]-[1,2,4]triazolo[4,3-a][1,4]diazepines, and 6-(2-chlorophenyl)-8,9-dihydro-1-methyl-8-[(4-morpholinyl)carbonyl]-4H,7H-cyclopenta-[4,5]thieno-[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepines.

[0373] MRP4-inhibitors used are preferably compounds selected from among N-acetyldinitrophenylcysteine, cGMP, cholate, diclofenac, dehydroepiandrosterone 3-glucuronide, dehydroepiandrosterone 3-sulfate, dilazep, dinitrophenyl-S-glutathione, estradiol 17-β-glucuronide, estradiol 3,17-disulfate, estradiol 3-glucuronide, estradiol 3-sulfate, estrone

3-sulfate, flurbiprofen, folate, N⁵-formyltetrahydrofolate, glycocholate, glycolithocholic acid sulfate, ibuprofen, indomethacin, indoprofen, ketoprofen, lithocholic acid sulfate, methotrexate, MK571 ((E)-3-[[[3-[2-(7-chloro-2-quinolinyl)ethenyl]phenyl]-[[3-dimethylamino)-3-oxopropyl]thio]methyl]thio]propanoic acid), α-naphthyl-β-D-glucuronide, nitrobenzyl mercaptopurine riboside, probenecid, PSC833, sildenafil, sulfinpyrazone, taurochenodeoxycholate, taurocholate, taurodeoxycholate, tauroolithocholate, tauroolithocholic acid sulfate, topotecan, trequinsin, and zaprinast, dipyrindamole, optionally in the form of the racemates, enantiomers, diastereomers, and the pharmacologically acceptable acid addition salts and hydrates thereof.

[0374] Preferably the invention relates to the use of MRP4-inhibitors for preparing a pharmaceutical composition for the treatment of respiratory complaints, containing the PDE4B-inhibitors and MRP4-inhibitors, the MRP4-inhibitors preferably being selected from among N-acetyldinitrophenylcysteine, dehydroepiandrosterone 3-sulfate, dilazep, dinitrophenyl-S-glutathione, estradiol 3,17-disulfate, flurbiprofen, glycocholate, glycolithocholic acid sulfate, ibuprofen, indomethacin, indoprofen, lithocholic acid sulfate, MK571, PSC833, sildenafil, taurochenodeoxycholate, taurocholate, tauroolithocholate, tauroolithocholic acid sulfate, trequinsin, and zaprinast, dipyrindamole, optionally in the form of the racemates, enantiomers, diastereomers, and the pharmacologically acceptable acid addition salts and hydrates thereof.

[0375] The invention relates more preferably to the use of MRP4-inhibitors for preparing a pharmaceutical composition for treating respiratory complaints, containing the PDE4B-inhibitors and MRP4-inhibitors according to the invention, the MRP4-inhibitors preferably being selected from among dehydroepiandrosterone 3-sulfate, estradiol 3,17-disulfate, flurbiprofen, indomethacin, indoprofen, MK571, taurocholate, optionally in the form of the racemates, enantiomers, diastereomers, and the pharmacologically acceptable acid addition salts and hydrates thereof. The separation of enantiomers from the racemates can be carried out using methods known from the art (e.g., chromatography on chiral phases, etc.).

[0376] By acid addition salts with pharmacologically acceptable acids are meant, for example, salts selected from among the hydrochlorides, hydrobromides, hydroiodides, hydrosulfates, hydrophosphates, hydromethanesulfonates, hydronitrates, hydromaleates, hydroacetates, hydrobenzoates, hydrocitrate, hydroftimrates, hydrotartrates, hydroxalates, hydrosuccinates, hydrobenzoates, and hydro-p-toluenesulfonates, preferably the hydrochlorides, hydrobromides, hydrosulfates, hydrophosphates, hydrofumarates, and hydromethanesulfonates.

[0377] The invention further relates to pharmaceutical preparations which contain a triple combination of the PDE4B-inhibitors, MRP4-inhibitors, and another active substance according to the invention, such as, for example, an anticholinergic, a steroid, an LTD4-antagonist, or a betamimetic, and the preparation thereof, and the use thereof for treating respiratory complaints.

Formulations

[0378] Suitable forms for administration are, for example, tablets, capsules, solutions, syrups, emulsions, or inhalable

powders or aerosols. The content of the pharmaceutically effective compound(s) in each case should be in the range from 0.1 to 90 wt. %, preferably 0.5 to 50 wt. % of the total composition, i.e., in amounts which are sufficient to achieve the dosage range specified hereinafter.

[0379] The preparations may be administered orally in the form of a tablet, as a powder, as a powder in a capsule (e.g., a hard gelatine capsule), as a solution or suspension. When administered by inhalation the active substance combination may be given as a powder, as an aqueous or aqueous-ethanolic solution or using a propellant gas formulation.

[0380] Preferably, therefore, pharmaceutical formulations are characterized by the content of one or more compounds of formula 1 according to the preferred embodiments above.

[0381] It is particularly preferable if the compounds of formula 1 are administered orally, and it is also particularly preferable if they are administered once or twice a day. Suitable tablets may be obtained, for example, by mixing the active substance(s) with known excipients, for example, inert diluents such as calcium carbonate, calcium phosphate, or lactose, disintegrants such as corn starch or alginic acid, binders such as starch or gelatine, lubricants such as magnesium stearate or talc, and/or agents for delaying release, such as carboxymethyl cellulose, cellulose acetate phthalate, or polyvinyl acetate. The tablets may also comprise several layers.

[0382] Coated tablets may be prepared accordingly by coating cores produced analogously to the tablets with substances normally used for tablet coatings, for example, collidone or shellac, gum arabic, talc, titanium dioxide, or sugar. To achieve delayed release or prevent incompatibilities the core may also consist of a number of layers. Similarly the tablet coating may consist of a number of layers to achieve delayed release, possibly using the excipients mentioned above for the tablets.

[0383] Syrups containing the active substances or combinations thereof according to the invention may additionally contain a sweetener such as saccharine, cyclamate, glycerol, or sugar, and a flavor enhancer, e.g., a flavoring such as vanillin or orange extract. They may also contain suspension adjuvants or thickeners such as sodium carboxymethyl cellulose, wetting agents such as, for example, condensation products of fatty alcohols with ethylene oxide, or preservatives such as *asp*-hydroxybenzoates.

[0384] Capsules containing one or more active substances or combinations of active substances may, for example, be prepared by mixing the active substances with inert carriers such as lactose or sorbitol and packing them into gelatine capsules.

[0385] Suitable suppositories may be made, for example, by mixing with carriers provided for this purpose, such as neutral fats or polyethyleneglycol or the derivatives thereof.

[0386] Excipients which may be used include, for example, water, pharmaceutically acceptable organic solvents such as paraffins (e.g., petroleum fractions), vegetable oils (e.g., groundnut or sesame oil), mono- or polyfunctional alcohols (e.g., ethanol or glycerol), carriers such as, e.g., natural mineral powders (e.g., kaolins, clays, talc, chalk), synthetic mineral powders (e.g., highly dispersed silicic acid and silicates), sugars (e.g., cane sugar, lactose, and glucose),

emulsifiers (e.g., lignin, spent sulfite liquors, methylcellulose, starch, and polyvinylpyrrolidone), and lubricants (e.g., magnesium stearate, talc, stearic acid, and sodium lauryl sulfate).

[0387] For oral administration the tablets may, of course, contain, apart from the abovementioned carriers, additives such as sodium citrate, calcium carbonate, and dicalcium phosphate together with various additives such as starch, preferably potato starch, gelatine, and the like. Moreover, lubricants such as magnesium stearate, sodium lauryl sulfate, and talc may be used at the same time for the tableting process. In the case of aqueous suspensions the active substances may be combined with various flavor enhancers or colorings in addition to the excipients mentioned above.

[0388] It is also preferred if the compounds of formula 1 are administered by inhalation, particularly preferably if they are administered once or twice a day. For this purpose, the compounds of formula 1 have to be made available in forms suitable for inhalation. Inhalable preparations include inhalable powders, propellant-containing metered-dose aerosols or propellant-free inhalable solutions, which are optionally present in admixture with conventional physiologically acceptable excipients.

[0389] Within the scope of the present invention, the term propellant-free inhalable solutions also includes concentrates or sterile ready-to-use inhalable solutions. The preparations which may be used according to the invention are described in more detail in the next part of the specification.

Inhalable Powders

[0390] If the active substances of formula 1 are present in admixture with physiologically acceptable excipients, the following physiologically acceptable excipients may be used to prepare the inhalable powders according to the invention: monosaccharides (e.g., glucose or arabinose), disaccharides (e.g., lactose, saccharose, maltose), oligo- and polysaccharides (e.g., dextran), polyalcohols (e.g., sorbitol, mannitol, xylitol), salts (e.g., sodium chloride, calcium carbonate) or mixtures of these excipients with one another. Preferably, mono- or disaccharides are used, while the use of lactose or glucose is preferred, particularly, but not exclusively, in the form of their hydrates. For the purposes of the invention, lactose is the particularly preferred excipient, while lactose monohydrate is most particularly preferred. Methods of preparing the inhalable powders according to the invention by grinding and micronizing and by finally mixing the components together are known from the prior art.

Propellant-Containing Inhalable Aerosols

[0391] The propellant-containing inhalable aerosols which may be used according to the invention may contain 1 dissolved in the propellant gas or in dispersed form. The propellant gases which may be used to prepare the inhalation aerosols according to the invention are known from the prior art. Suitable propellant gases are selected from among hydrocarbons such as *n*-propane, *n*-butane, or isobutane, and haloalkanes such as preferably fluorinated derivatives of methane, ethane, propane, butane, cyclopropane, or cyclobutane. The propellant gases mentioned above may be used on their own or in mixtures thereof. Particularly preferred propellant gases are fluorinated alkane derivatives selected from TG134a (1,1,1,2-tetrafluoroethane), TG227 (1,1,1,2,3,3,3-heptafluoropropane), and mixtures thereof.

The propellant-driven inhalation aerosols used within the scope of the use according to the invention may also contain other ingredients such as co-solvents, stabilizers, surfactants, antioxidants, lubricants, and pH adjusters. All these ingredients are known in the art.

Propellant-Free Inhalable Solutions

[0392] The compounds of formula 1 according to the invention are preferably used to prepare propellant-free inhalable solutions and inhalable suspensions. Solvents used for this purpose include aqueous or alcoholic, preferably ethanolic solutions. The solvent may be water on its own or a mixture of water and ethanol. The solutions or suspensions are adjusted to a pH of 2 to 7, preferably 2 to 5, using suitable acids. The pH may be adjusted using acids selected from inorganic or organic acids. Examples of particularly suitable inorganic acids include hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, and/or phosphoric acid. Examples of particularly suitable organic acids include ascorbic acid, citric acid, malic acid, tartaric acid, maleic acid, succinic acid, fumaric acid, acetic acid, formic acid, and/or propionic acid, etc. Preferred inorganic acids are hydrochloric and sulfuric acids. It is also possible to use the acids which have already formed an acid addition salt with one of the active substances. Of the organic acids, ascorbic acid, fumaric acid, and citric acid are preferred. If desired, mixtures of the above acids may also be used, particularly in the case of acids which have other properties in addition to their acidifying qualities, e.g., as flavorings, antioxidants, or complexing agents, such as citric acid or ascorbic acid, for example. According to the invention, it is particularly preferred to use hydrochloric acid to adjust the pH.


[0393] Co-solvents and/or other excipients may be added to the propellant-free inhalable solutions used for the purpose according to the invention. Preferred co-solvents are those which contain hydroxyl groups or other polar groups, e.g., alcohols—particularly isopropyl alcohol, glycols—particularly propyleneglycol, polyethyleneglycol, polypropylene glycol, glycol ether, glycerol, polyoxyethylene alcohols, and polyoxyethylene fatty acid esters. The terms excipients and additives in this context denote any pharmacologically acceptable substance which is not an active substance but which can be formulated with the active substance or substances in the pharmacologically suitable solvent in order to improve the qualitative properties of the active substance formulation. Preferably, these substances have no pharmacological effect or, in connection with the desired therapy, no appreciable or at least no undesirable pharmacological effect. The excipients and additives include, for example, surfactants such as soya lecithin, oleic acid, sorbitan esters, such as polysorbates, polyvinylpyrrolidone, other stabilizers, complexing agents, antioxidants, and/or preservatives which guarantee or prolong the shelf life of the finished pharmaceutical formulation, flavorings, vitamins, and/or other additives known in the art. The additives also include pharmacologically acceptable salts such as sodium chloride as isotonic agents. The preferred excipients include antioxidants such as ascorbic acid, for example, provided that it has not already been used to adjust the pH, vitamin A, vitamin E, tocopherols, and similar vitamins or provitamins occurring in the human body. Preservatives may be used to protect the formulation from contamination with pathogens. Suitable preservatives are those which are known in the art, particularly cetyl pyri-

dinium chloride, benzalkonium chloride, or benzoic acid or benzoates such as sodium benzoate in the concentration known from the prior art.

[0394] For the treatment forms described above, ready-to-use packs of a medicament for the treatment of respiratory complaints are provided, containing an enclosed description including, for example, the words respiratory disease, COPD, or asthma, a pteridine and one or more combination partners selected from those described above.

We claim:

1. A compound of formula 1



wherein:

A is CO, C=NH, C₁₋₆-alkylene, or C₃₋₈-cycloalkylene;

B¹ is phenyl or an aromatic or non-aromatic ring optionally containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen and which are optionally mono- or polysubstituted by one or more groups selected, from OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, C₁₋₆-alkyl, and C₁₋₆-haloalkyl;

B² is phenyl or a heteroaryl containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen;

X is O, S, NR⁵, or CR⁶R⁷;


n is 0, 1, 2, or 3;

R¹ is H, C₁₋₆-alkyl, C₁₋₆-haloalkyl, COR^{1.1}, COOR^{1.1}, or CH₂COOR^{1.1}, wherein R^{1.1} is H or C₁₋₆-alkyl;

R² is H, C₁₋₆-alkyl, or C₁₋₆-haloalkyl; or

R¹ and R² together with the nitrogen form a non-aromatic heterocycle optionally containing one, two, or three heteroatoms selected from oxygen and nitrogen; or

R², N, A, and B¹ together form a bicyclic group of formula (i)



wherein: A is CO, C=NH, or C₁₋₃-alkyl, and m is 1, 2, or 3;

R^3 is H or a group selected from OH, C_{1-6} -haloalkyl, a C_{6-10} -aryl, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, wherein the C_{3-10} -heterocycle and the C_{5-10} -heteroaryl contain one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen, optionally substituted by one or more groups selected from C_{1-6} -alkyl, C_{6-10} -aryl, optionally bridged C_{3-8} -cycloalkyl and C_{1-6} -haloalkyl, each optionally substituted by a group selected from C_{1-6} -alkyl, C_{1-4} -haloalkyl, OH, halogen, and C_{6-10} -aryl,


R^3 is C_{1-6} -alkyl optionally substituted by one or more groups selected from halogen, OH, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, and $N(C_{1-6}$ -alkyl) $_2$,

R^3 is a group selected from C_{3-8} -cycloalkyl, C_{5-8} -cycloalkenyl, C_{1-6} -alkyl, and C_{1-6} -alkanol, each optionally substituted by one or more groups selected from C_{6-10} -aryl, C_{3-8} -cycloalkyl, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, each optionally substituted by one or more groups selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, CO -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, $N(C_{1-6}$ -alkyl) $_2$, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, wherein the C_{5-10} -heteroaryl and the C_{3-10} -heterocycle are each optionally substituted by a group selected from oxo, hydroxyl, halogen, C_{1-6} -alkyl, and C_{1-6} -haloalkyl,

R^3 is a group selected from C_{6-10} -aryl, a C_{6-10} -heteroaryl, and a C_{3-10} -heterocycle, each optionally substituted by one or more groups selected from aryl, C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, $N(C_{1-6}$ -alkyl) $_2$, and $N-(SO_2-C_{1-4}$ -alkyl)($R^{3,4}$),

R^3 is a group selected from C_{1-6} -alkyl, C_{6-10} -aryl- C_{1-6} -alkylene, C_{5-10} -heteroaryl- C_{1-6} -alkylene, C_{3-7} -cycloalkyl, C_{6-10} -aryl, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, each optionally substituted by one or more groups selected from B, halogen, OH, C_{1-6} -alkyl, and oxo, wherein B is a compound of formula 2

wherein:



Z^1 is H, OH, halogen, C_{1-6} -alkyl, C_{1-6} -alkanol, $O(C_{1-6}$ -alkyl), C_{6-10} -aryl, $O-C_{6-10}$ -aryl, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, or C_{3-7} -cycloalkyl, and

Z^2 is OH, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, $O(C_{1-6}$ -alkyl), mono- or bicyclic C_{3-7} -cycloalkyl, mono- or bicyclic C_{3-10} -heterocycle, mono- or bicyclic C_{5-10} -heteroaryl, or C_{6-10} -aryl,

R^3 is a group selected from C_{6-10} -aryl and a C_{5-10} -heteroaryl, each optionally substituted by C_{1-6} -alkyl optionally substituted by a group selected from $COOR^{3,3}$, $NR^{3,3}R^{3,4}$, $NHCOR^{3,3}$, $NHCOOR^{3,3}$, and phenyl optionally substituted by a group selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, CN, C_{1-6} -haloalkyl, $CONR_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, and $N(C_{1-6}$ -alkyl) $_2$, a C_{3-10} -heterocycle, and a C_{5-10} -heteroaryl, wherein the C_{3-10} -heterocycle and the C_{5-10} -heteroaryl are each optionally substituted by an oxo group or a methyl group, wherein:

$R^{3,3}$ is H or C_{1-6} -alkyl, and

$R^{3,4}$ is H, C_{1-6} -alkyl, or C_{7-11} -aralkyl, C_{1-6} -alkylene, or C_{5-10} -heteroaryl- C_{1-6} -alkylene,

R^3 is a group selected from C_{6-10} -aryl and a C_{5-10} -heteroaryl, each optionally substituted by $NR^{3,1}R^{3,2}$, wherein:

$R^{3,1}$ is H, C_{1-6} -alkyl, C_{3-6} -cycloalkyl, C_{1-6} -haloalkyl, $COR^{3,1,1}$, $COOR^{3,1,1}$, $CONR^{3,1,1}R^{3,1,2}$, or $SO_2-R^{3,1,1}$, wherein $R^{3,1,1}$ is H, C_{1-6} -alkyl, C_{1-6} -haloalkyl, C_{3-6} -cycloalkyl, or C_{6-10} -aryl, and $R^{3,1,2}$ is H, C_{1-6} -alkyl, C_{1-6} -haloalkyl, C_{3-6} -cycloalkyl, or C_{6-10} -aryl, and $R^{3,2}$ is H or a group selected from C_{1-6} -alkyl, C_{3-6} -cycloalkyl, and C_{1-6} -haloalkyl, each optionally substituted by one or more groups selected from NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, oxo, and a non-aromatic C_{3-10} -heterocycle containing one or two heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the non-aromatic C_{3-10} -heterocycle is optionally substituted by C_{1-4} -alkyl, or

R³ is a group selected from C₆₋₁₀-aryl and a C₅₋₁₀-heteroaryl, each optionally substituted by a C₃₋₁₀-heterocycle optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, CN, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, oxo, OH, O—C₁₋₆-alkyl, halogen, SH, S—C₁₋₆-alkyl, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂, or

R³ is benzimidazolyl optionally substituted by a group selected from C₁₋₆-alkyl, C₁₋₆-haloalkyl, and C₃₋₆-cycloalkyl; and

R⁴ is C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-haloalkyl, OR^{4.1}, NR^{4.1}R^{4.2}, CN, or halogen, wherein: R^{4.1} is H, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, or C₁₋₆-haloalkyl, and R^{4.2} is H, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, or C₁₋₆-haloalkyl;

R⁵ is C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-haloalkyl, COR^{5.1}, CONHR^{5.1}, CON(R^{5.1})₂, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-haloalkyl, SO₂-aryl, or a group selected from R^{5.2}, SO₂—C₁₋₆-alkyl-R^{5.2}, and C₁₋₆-alkyl-R^{5.2}, each optionally substituted by one or more groups selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, CN, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, O—C₃₋₆-cycloalkyl, O—C₁₋₆-haloalkyl, O—C₁₋₆-alkyl, halogen, SO₂—C₁₋₄-alkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂, wherein: R^{5.1} is C₁₋₆-alkyl, C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, or C₆₋₁₀-aryl-C₁₋₆-alkylene, and R^{5.2} is C₆₋₁₀-aryl or a C₅₋₁₀-heteroaryl;

R⁶ is H, C₁₋₆-alkyl, or C₁₋₆-haloalkyl; and

R⁷ is H, C₁₋₆-alkyl, or C₁₋₆-haloalkyl; or

R⁶ and R⁷ together form a 3-6 membered carbocycle,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

2. The compound of formula 1 according to claim 1, wherein:

B¹ is phenyl or an aromatic or non-aromatic ring optionally containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen and optionally mono- or polysubstituted by one or more groups selected from OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, C₁₋₆-alkyl, and C₁₋₆-haloalkyl;

B² is phenyl or pyridinyl;

X is O, S, NR⁵, or CR⁶R⁷;

n is 0, 1, 2, or 3;

R¹ is H, C₁₋₄-alkyl, or C₁₋₄-haloalkyl;

R² is H, C₁₋₄-alkyl, or C₁₋₄-haloalkyl; or

R¹ and R² together with the nitrogen form a non-aromatic heterocycle containing one or two nitrogen atoms;


R³ is H, OH, C₁₋₆-haloalkyl, or a group selected from C₆₋₁₀-aryl, a C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, each optionally substituted by a methyl group, oxo, or OH,

R³ is C₁₋₆-alkyl optionally substituted by one or more groups selected from halogen, OH, CN, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, COaryl, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, SH, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, SO₂—C₁₋₆-alkanol, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₆-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂,

R³ is a group selected from C₃₋₈-cycloalkyl, a C₃₋₈-cycloalkyl bridged by C₁₋₃-alkylene, C₅₋₈-cycloalkenyl, C₁₋₆-alkyl, and C₁₋₆-alkanol, each optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₃₋₈-cycloalkyl, C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, each optionally substituted by one or more groups selected from C₁₋₆-alkyl, C₁₋₆-haloalkyl, CN, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, NO₂, NH₂, NH—C₁₋₆-alkyl, N(C₁₋₆-alkyl)₂, a C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, wherein the C₅₋₁₀-heteroaryl and the C₃₋₁₀-heterocycle are each optionally substituted by a group selected from oxo, hydroxyl, halogen, C₁₋₆-alkyl, and C₁₋₆-haloalkyl,

R³ is a group selected from C₁₋₆-alkyl, C₆₋₁₀-aryl-C₁₋₆-alkylene, C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, C₃₋₇-cycloalkyl, C₆₋₁₀-aryl, C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, each substituted one or more groups selected from B, halogen, OH, C₁₋₆-alkyl, oxo, wherein B is a compound of formula 2

wherein:



Z¹ is H, OH, halogen, C₁₋₆-alkyl, C₁₋₆-alkanol, O(C₁₋₆-alkyl), C₆₋₁₀-aryl, O—C₆₋₁₀-aryl, NH₂, NH(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂, or C₃₋₇-cycloalkyl, and

Z² is OH, NH₂, NH(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂, O(C₁₋₆-alkyl), mono- or bicyclic C₃₋₇-cycloalkyl, a mono- or bicyclic C₃₋₁₀-heterocycle, a mono- or bicyclic C₅₋₁₀-heteroaryl, or C₆₋₁₀-aryl,

R³ is phenyl optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, CONH₂, CONH—C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO—C₁₋₆-alkyl, COH, CO—C₁₋₆-alkyl, COaryl, OH, O—C₁₋₆-alkyl, O—C₁₋₄-haloalkyl, halogen, SH, S—C₁₋₆-alkyl, S—C₁₋₄-haloalkyl, SO₂—C₁₋₆-alkyl, SO₂—C₁₋₄-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₆-alkyl, SO₂—N(C₁₋₆-alkyl)₂, NO₂, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂,

R³ is phenyl substituted by C₁₋₄-alkyl optionally substituted by a group selected from COOR^{3.3}, NR^{3.3}R^{3.4},

NHCOR^{3.3}, NHCOOR^{3.3}, and phenyl, each optionally substituted by one or more groups selected from methyl, tert-butyl, F, Cl, Br, CN, OH, and a heterocycle containing one, two, or three heteroatoms selected from oxygen and nitrogen, wherein the heterocycle is optionally substituted by an oxo group or a methyl group, wherein:

R^{3.3} is H or C₁₋₆-alkyl, and

R^{3.4} is H, C₁₋₆-alkyl, C₆₋₁₀-aryl-C₁₋₆-alkylene, or C₅₋₁₀-heteroaryl-C₁₋₆-alkylene,

R³ is phenyl substituted with NR^{3.1}R^{3.2}, wherein:

R^{3.1} is H, C₁₋₄-alkyl, C₁₋₄-haloalkyl, COR^{3.1.1}, COOR^{3.1.1}, CONR^{3.1.1}R^{3.1.2}, or SO₂-R^{3.1.1}, wherein R^{3.1.1} is H, C₁₋₄-alkyl, C₁₋₄-haloalkyl, C₃₋₆-cycloalkyl, or C₆₋₁₀-aryl, and R^{3.1.2} is H, C₁₋₄-alkyl, C₁₋₄-haloalkyl, C₃₋₆-cycloalkyl, or C₆₋₁₀-aryl, and

R^{3.2} is H or C₁₋₄-alkyl optionally substituted by one or more groups selected from NH₂, NH(C₁₋₄-alkyl), N(C₁₋₄-alkyl)₂, oxo, or a non-aromatic C₃₋₁₀-heterocycle optionally containing one or two nitrogen atoms and optionally substituted by a methyl group,

R³ is C₆₋₁₀-aryl optionally substituted by a C₅₋₁₀-heteroaryl containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen, and optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₁₋₄-alkyl, C₃₋₆-cycloalkyl, CN, C₁₋₄-haloalkyl, CONH₂, CONH-C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO-C₁₋₄-alkyl, COH, CO-C₁₋₄-alkyl, OH, O-C₁₋₄-alkyl, halogen, SH, S-C₁₋₄-alkyl, NH₂, NH-C₁₋₄-alkyl, and N(C₁₋₄-alkyl)₂,

R³ is C₆₋₁₀-aryl optionally substituted by a non-aromatic C₃₋₁₀-heterocycle optionally containing one or two heteroatoms selected from oxygen, sulfur, and nitrogen, wherein the C₃₋₁₀-heterocycle is optionally substituted by one or more groups selected from C₁₋₄-alkyl and oxo, or

R³ is benzimidazolyl optionally substituted by one or more groups selected from C₁₋₄-alkyl, C₁₋₄-haloalkyl, and C₃₋₆-cycloalkyl;

R⁴ is C₁₋₄-alkyl, C₃₋₆-cycloalkyl, C₁₋₄-haloalkyl, OR^{4.1}, NR^{4.1}R^{4.2}, CN, or halogen, wherein:

R^{4.1} is H, C₁₋₄-alkyl, C₃₋₆-cycloalkyl, or C₁₋₄-haloalkyl, and

R^{4.2} is H, C₁₋₄-alkyl, C₃₋₆-cycloalkyl, or C₁₋₄-haloalkyl;

R⁵ is C₁₋₄-alkyl, C₃₋₈-cycloalkyl, C₁₋₄-haloalkyl, COR^{5.1}, CONHR^{5.1}, CON(R^{5.1})₂, SO₂-C₁₋₄-alkyl, SO₂-C₁₋₄-haloalkyl, SO₂-aryl, or a group selected from R^{5.2}, SO₂-C₁₋₄-alkyl-R^{5.2}, and C₁₋₄-alkyl-R^{5.2}, each optionally substituted by C₁₋₄-alkyl, C₂₋₄-alkenyl, C₂₋₄-alkynyl, CN, C₁₋₄-haloalkyl, CONH₂, CONH-C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO-C₁₋₄-alkyl, COH, CO-C₁₋₄-alkyl, O-C₁₋₄-cycloalkyl, O-C₁₋₄-haloalkyl, O-C₁₋₄-alkyl, halogen, SO₂-C₁₋₄-alkyl, SO₂-NH₂, SO₂-NH-C₁₋₄-alkyl, SO₂-N(C₁₋₄-alkyl)₂, NO₂, NH₂, NH-C₁₋₄-alkyl, and N(C₁₋₄-alkyl)₂, wherein:

R^{5.1} is C₁₋₆-alkyl, C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, or C₆₋₁₀-aryl-C₁₋₆-alkylene, and R^{5.2} is C₆₋₁₀-aryl or a C₅₋₁₀-heteroaryl;

R⁶ is H, C₁₋₆-alkyl, or C₁₋₆-haloalkyl; and

R⁷ is H, C₁₋₆-alkyl, or C₁₋₆-haloalkyl; or

R⁶ and R⁷ together form a 3-6 membered carbocycle,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

3. The compound of formula 1 according to claim 1, wherein:


R³ is C₁₋₆-alkyl optionally substituted by one or more groups selected from halogen, OH, CN, CONH₂, CONH-C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO-C₁₋₆-alkyl, COH, CO-C₁₋₆-alkyl, COaryl, OH, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen, SH, S-C₁₋₆-alkyl, S-C₁₋₆-haloalkyl, SO₂-C₁₋₆-alkanol, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-haloalkyl, SO₂-NH₂, SO₂-NH-C₁₋₆-alkyl, SO₂-N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH-C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂,

R³ is a group selected from C₃₋₈-cycloalkyl, a C₃₋₈-cycloalkyl bridged with C₁₋₃-alkylene, a C₅₋₈-cycloalkenyl, C₁₋₆-alkyl, and C₁₋₆-alkanol, each optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₃₋₈-cycloalkyl, a C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, each optionally substituted by one or more groups selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, CN, CONH₂, CONH-C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO-C₁₋₆-alkyl, COH, CO-C₁₋₆-alkyl, CO-C₆₋₁₀-aryl, OH, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen, SH, S-C₁₋₆-alkyl, S-C₁₋₆-haloalkyl, SO₂-C₁₋₆-alkanol, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-haloalkyl, SO₂-NH₂, SO₂-NH-C₁₋₆-alkyl, SO₂-N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH-C₁₋₆-alkyl, N(C₁₋₆-alkyl)₂, C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, each optionally substituted by one or more groups selected from oxo, hydroxyl, halogen, or C₁₋₆-alkyl, and C₁₋₆-haloalkyl,

R³ is a group selected from C₆₋₁₀-aryl, C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, each optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, CONH₂, CONH-C₁₋₆-alkyl, CON(C₁₋₆-alkyl)₂, COOH, COO-C₁₋₆-alkyl, COH, CO-C₁₋₆-alkyl, CO-C₆₋₁₀-aryl, OH, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen, SH, S-C₁₋₆-alkyl, S-C₁₋₆-haloalkyl, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-alkanol, SO₂-C₁₋₆-haloalkyl, SO₂-NH₂, SO₂-NH-C₁₋₆-alkyl, SO₂-N(C₁₋₆-alkyl)₂, NO₂, NH₂, NH-C₁₋₆-alkyl, N(C₁₋₆-alkyl)₂, and N-(SO₂-C₁₋₄-alkyl)(R^{3.4}), or

R³ is a group selected from C₁₋₆-alkyl, C₆₋₁₀-aryl-C₁₋₆-alkylene, C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, C₃₋₇-cycloalkyl, C₆₋₁₀-aryl, C₅₋₁₀-heteroaryl, and a C₃₋₁₀-heterocycle, each optionally substituted by one or more groups selected from B, halogen, OH, C₁₋₆-alkyl, and oxo, wherein B is a compound of formula 2

wherein:



Z^1 is H, OH, halogen, C_{1-6} -alkyl, C_{1-6} -alkanol, $O(C_{1-6}$ -alkyl), C_{6-10} -aryl, $O-C_{6-10}$ -aryl, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, or C_{3-7} -cycloalkyl, and

Z^2 is OH, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, $O(C_{1-6}$ -alkyl), mono- or bicyclic C_{3-7} -cycloalkyl, mono- or bicyclic C_{5-10} -heteroaryl, mono- or bicyclic C_{3-10} -heterocycle, or C_{6-10} -aryl,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

4. The compound of formula 1 according to claim 3, wherein:

A is CH_2 , CD_2 , $C=NH$, $CHMe$, CMe_2 , 1,1'-cyclopropylene, or 1,1'-cyclobutylidene;

B^1 and B^2 are each phenyl;

X is O or NR^5 , wherein R^5 is methyl, ethyl, cyclopropyl, cyclobutyl, $CONHCH_2$ -phenyl, CH_2CF_3 , or benzyl, each optionally substituted by F, and n is 0 or 1; and

R^1 and R^2 are each H,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

5. The compound of formula 1 according to claim 3, wherein:

R^3 is C_{1-6} -alkyl optionally substituted by one or more groups selected from halogen, OH, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, and $N(C_{1-6}$ -alkyl) $_2$,


R^3 is a group selected from C_{3-8} -cycloalkyl, a C_{3-8} -cycloalkyl bridged by C_{1-3} -alkylene, and C_{1-6} -alkyl, each optionally substituted by one or more groups selected from C_{6-10} -aryl, C_{3-8} -cycloalkyl, a C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, each optionally substituted by one or more groups selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, CN, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SH, $S-C_{1-6}$ -alkyl, $S-C_{1-6}$ -haloalkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl,

$N(C_{1-6}$ -alkyl) $_2$, C_{5-10} -heteroaryl, and a C_{3-10} -heterocycle, each optionally substituted by one or more groups selected from oxo, hydroxyl, halogen, or C_{1-6} -alkyl, and C_{1-6} -haloalkyl,

R^3 is a group selected from C_{6-10} -aryl, a C_{3-8} -heterocycle with 1 to 4 heteroatoms selected from N, O, and S, and a C_{5-10} -heteroaryl with 1 to 2 heteroatoms selected from N, O, and S, each optionally substituted by one or more groups selected from C_{6-10} -aryl, C_{1-6} -alkyl, C_{1-6} -haloalkyl, $CONH_2$, $CONH-C_{1-6}$ -alkyl, $CON(C_{1-6}$ -alkyl) $_2$, $COOH$, $COO-C_{1-6}$ -alkyl, COH , $CO-C_{1-6}$ -alkyl, $CO-C_{6-10}$ -aryl, OH, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -alkanol, SO_2-C_{1-6} -haloalkyl, SO_2-NH_2 , SO_2-NH-C_{1-6} -alkyl, $SO_2-N(C_{1-6}$ -alkyl) $_2$, NO_2 , NH_2 , $NH-C_{1-6}$ -alkyl, $N(C_{1-6}$ -alkyl) $_2$, and $N-(SO_2-C_{1-4}$ -alkyl)($R^{3,4}$), wherein $R^{3,4}$ is a C_{5-10} -heteroaryl- C_{1-6} -alkylene, or

R^3 is a group selected from C_{1-6} -alkyl, C_{6-10} -aryl- C_{1-6} -alkylene, C_{5-10} -heteroaryl- C_{1-6} -alkylene, C_{3-7} -cycloalkyl, C_{6-10} -aryl, C_{3-8} -heterocycle with 1 to 4 heteroatoms selected from N, O, and S, and a C_{5-10} -heteroaryl with 1 to 2 heteroatoms selected from N, O, and S, each optionally substituted by one or more groups selected from B, halogen, OH, C_{1-6} -alkyl, and oxo, wherein B is a compound of formula 2

wherein:



Z^1 is H, OH, halogen, C_{1-6} -alkyl, C_{1-6} -alkanol, or $O(C_{1-6}$ -alkyl), and

Z^2 is OH, NH_2 , $NH(C_{1-6}$ -alkyl), $N(C_{1-6}$ -alkyl) $_2$, $O(C_{1-6}$ -alkyl), mono- or bicyclic C_{3-7} -cycloalkyl, mono- or bicyclic C_{5-10} -heteroaryl, mono- or bicyclic C_{3-10} -heterocycle, or C_{6-10} -aryl; and

R^4 is H, F, or Cl,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

6. The compound of formula 1 according to claim 1, wherein:

A is CO, $C=NH$, C_{1-4} -alkylene, C_{3-6} -cycloalkylene, C_{1-6} -alkylene, or C_{3-8} -cycloalkylene;

B^1 is phenyl or pyridinyl;

B^2 is phenyl or pyridinyl;

X is O or NR^5 ;

n is 0, 1, 2, or 3;

R^1 is H, methyl, ethyl, or propyl;

R² is H, methyl, ethyl, or propyl;

R³ is H, OH, C₁₋₆-haloalkyl, or C₆₋₁₀-aryl, or a group selected from a C₅₋₁₀-heteroaryl and a C₃₋₁₀-cycloalkyl, each containing one, two, or three nitrogen atoms and optionally substituted by a methyl group,

R³ is a group selected from cyclopentyl, cyclohexyl, cyclopentenyl, cyclohexenyl, methyl, ethyl, propyl, and butyl, each optionally substituted by one or more groups selected from C₆₋₁₀-aryl and a C₅₋₁₀-heterocycle, each optionally substituted by one or more groups selected from C₁₋₆-alkyl, C₁₋₆-haloalkyl, CN, OH, O—C₁₋₆-alkyl, O—C₁₋₆-haloalkyl, halogen, S—C₁₋₆-alkyl, S—C₁₋₆-haloalkyl, NO₂, NH₂, NH—C₁₋₆-alkyl, and N(C₁₋₆-alkyl)₂,

R³ is phenyl optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₂₋₄-alkenyl, C₂₋₄-alkynyl, C₁₋₄-haloalkyl, CONH₂, CONH—C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO—C₁₋₄-alkyl, COH, CO—C₁₋₄-alkyl, CO—C₆₋₁₀-aryl, OH, O—C₁₋₄-alkyl, O—C₁₋₄-haloalkyl, halogen, SH, S—C₁₋₄-alkyl, S—C₁₋₄-haloalkyl, SO₂—C₁₋₄-alkyl, SO₂—C₁₋₄-haloalkyl, SO₂—NH₂, SO₂—NH—C₁₋₄-alkyl, SO₂—N(C₁₋₄-alkyl)₂, NO₂, NO₂, NH₂, NH—C₁₋₄-alkyl, and N(C₁₋₄-alkyl)₂,

R³ is phenyl optionally substituted by C₁₋₄-alkyl optionally substituted by a group selected from COOR^{3.3}, NR^{3.3}R^{3.4}, NHCOR^{3.3}, NHCOOR^{3.3}, p-fluorophenyl, and a heterocycle containing one, two, or three heteroatoms selected from oxygen and nitrogen and optionally substituted by an oxo group, wherein:

R^{3.3} is H or C₁₋₄-alkyl, and

R^{3.4} is H, C₁₋₄-alkyl, C₆₋₁₀-aryl-C₁₋₆-alkylene, or C₅₋₁₀-heteroaryl-C₁₋₆-alkylene,

R³ is phenyl optionally substituted by NR^{3.1}R^{3.2}, wherein:

R^{3.1} is H, C₁₋₄-alkyl, COR^{3.1.1}, COOR^{3.1.1}, CONR^{3.1.1}R^{3.1.2}, or SO₂—R^{3.1.1}, wherein R^{3.1.1} is H, C₁₋₄-alkyl, or C₆₋₁₀-aryl, and R^{3.1.2} is H, C₁₋₄-alkyl, or C₆₋₁₀-aryl, and

R^{3.2} is H or C₁₋₄-alkyl optionally substituted by one or more groups selected from NH₂, NH(C₁₋₄-alkyl), N(C₁₋₄-alkyl)₂, oxo, or a C₃₋₁₀-heterocycle containing one or two nitrogen atoms and optionally substituted by a methyl group,

R³ is phenyl optionally substituted by a C₅₋₁₀-heteroaryl containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen, wherein the C₅₋₁₀-heteroaryl is optionally substituted by one or more groups selected from C₆₋₁₀-aryl, C₁₋₄-alkyl, C₁₋₄-haloalkyl, C₃₋₆-cycloalkyl, CN, CONH₂, CONH—C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO—C₁₋₄-alkyl, COH, CO—C₁₋₄-alkyl, OH, O—C₁₋₄-alkyl, halogen, NH₂, and N(C₁₋₄-alkyl)₂,

R³ is phenyl optionally substituted by a C₅₋₁₀-heteroaryl containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen, wherein the C₅₋₁₀-heteroaryl is optionally substituted by one or more groups selected from C₁₋₄-alkyl and oxo, or


R³ is benzimidazolyl optionally substituted by one or more groups selected from methyl, ethyl, propyl, CF₃, CH₂CF₃, cyclopropyl, cyclopentyl, and cyclohexyl;

R⁴ is C₁₋₄-alkyl, C₁₋₄-haloalkyl, or halogen;

R⁵ is a group selected from C₁₋₄-alkyl, C₃₋₆-cycloalkyl, COR^{5.1}, CONHR^{5.1}, C₆₋₁₀-aryl, SO₂—C₆₋₁₀-aryl-C₁₋₆-alkylene, SO₂—C₆₋₁₀-aryl, C₆₋₁₀-aryl-C₁₋₆-alkylene, and C₅₋₁₀-heteroaryl-C₁₋₆-alkylene, each optionally substituted by one or more groups selected from C₁₋₄-alkyl, C₂₋₄-alkenyl, C₂₋₄-alkynyl, CN, C₁₋₄-haloalkyl, CONH₂, CONH—C₁₋₄-alkyl, CON(C₁₋₄-alkyl)₂, COOH, COO—C₁₋₄-alkyl, COH, CO—C₁₋₄-alkyl, OH, O—C₁₋₄-alkyl, halogen, SO₂—C₁₋₄-alkyl, SO₂—NH₂, SO₂—NH—C₁₋₄-alkyl, SO₂—N(C₁₋₄-alkyl)₂, NO₂, NH₂, NH—C₁₋₄-alkyl, and N(C₁₋₄-alkyl)₂, wherein R^{5.1} is C₁₋₄-alkyl or C₆₋₁₀-aryl-C₁₋₆-alkylene,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

7. The compound according to claim 1, having formula 1a



wherein A, X, n, R¹, R², R³, and R⁴ are as defined in claim 1, or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

8. The compound of formula 1 according to claim 1, wherein X is O, or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

9. The compound of formula 1a according to claim 7, wherein X is O, or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

10. The compound of formula 1 according to claim 1, wherein X is NR⁵, or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

11. The compound of formula 1a according to claim 7, wherein X is NR⁵, or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

12. The compound of formula 1 according to claim 1, wherein X is CR⁶R⁷, or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

13. The compound of formula 1a according to claim 7, wherein X is CR⁶R⁷, or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

14. The compound of formula 1 according to claim 1, wherein:

A is CH₂, CHMe, CMe₂, C=NH, 1,1'-cyclopropylene, or 1,1'-cyclobutylidene;

B¹ and B² are each phenyl;

X is O or NR⁵;

n is 0, 1, or 2;

R¹ is H, methyl, or ethyl;

R² is H, methyl, or ethyl;

R³ is H, cyclopropyl, cyclobutyl, N-methylpiperidinyl, pyridinyl, phenyl, or 4-phenylcyclohexane, R³ is phenyl optionally substituted by one or more groups selected from phenyl, methyl, ethyl, propyl, butyl, CF₃, CONH₂, CONHMe, CONMe₂, COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, SH, SO₂Me, SONH₂, SONMe₂, NO₂, NH₂, NHMe, and NMe₂, R³ is phenyl optionally substituted by a group selected from methyl and ethyl, each optionally substituted by one or more groups selected from COOH, COOMe, NH₂, NMe₂, NHCOMe, NHCOO-tert-butyl, NMe(benzyl), p-fluorophenyl, pyrrolidinyl, piperidinyl, morpholinyl, pyrrolidin-2-onyl, imidazolyl, and triazolyl,

R³ is phenyl substituted by NR^{3.1}R^{3.2}, wherein R^{3.1} is H, methyl, COH, COMe, COOMe, CONH₂, CONMe₂, SO₂Me, SO₂CF₃, or SO₂-phenyl, and R^{3.2} is H or a group selected from methyl and ethyl, each optionally substituted by one or more groups selected from NH₂, NHMe, NMe₂, N-piperidinyl, N-morpholinyl, and N-methylpiperazinyl, wherein the N-piperidinyl, N-morpholinyl, and the N-methylpiperazinyl are each optionally substituted by a further oxo,

R³ is phenyl optionally substituted by a C₅₋₁₀-heteroaryl containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen, wherein the C₅₋₁₀-heteroaryl is optionally substituted by one or more groups selected from phenyl, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, CF₃, CN, CONH₂, CONMe₂, CONEt₂, COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, NH₂, NMe₂, NEt₂, and NPr₂,

R³ is phenyl substituted by a C₃₋₁₀-heterocycle containing one or two heteroatoms selected from oxygen and nitrogen, wherein the C₃₋₁₀-heterocycle is optionally substituted by one or more groups selected from C₁₋₄-alkyl and oxo, or

R³ is benzimidazolyl optionally substituted by one or more groups selected from methyl, propyl, CF₃, CH₂CF₃, cyclopropyl, and cyclohexyl;

R⁴ is methyl, ethyl, propyl, butyl, CF₃, CH₂CF₃, F, Cl, or Br; and

R⁵ is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, CF₃, CH₂CF₃, COR^{5.1}, CONHR^{5.1}, phenyl, phe-

nylsulfonyl, or benzyl optionally substituted by one or more groups selected from methyl, ethyl, propyl, butyl, CF₃, CN, CONH₂, CONMe₂, CONEt₂, COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, SO₂Me, SONH₂, SONMe₂, NO₂, NH₂, NMe₂, NEt₂, and NPr₂, wherein R^{5.1} is methyl, ethyl, propyl, butyl, or benzyl,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

15. The compound of formula 1 according to claim 1, wherein:

A is CH₂, CHMe, CMe₂, 1,1'-cyclopropylene, or 1,1'-cyclobutylidene;

B¹ and B² are each phenyl;

X is O or NR⁵;

n is 0 or 1;

R¹ and R² are each H;

R³ is H or 4-phenylcyclohexanyl,

R³ is phenyl optionally substituted by NR^{3.1}R^{3.2}, wherein R^{3.1} is H, methyl, SO₂Me, SO₂CF₃, or SO₂-phenyl, and R^{3.2} is H or a group selected from methyl and ethyl, each optionally substituted by one or more groups selected from NH₂, NHMe, NMe₂, oxo, N-piperidinyl, N-morpholinyl, and N-methylpiperazinyl,

R³ is phenyl substituted by a C₅₋₁₀-heteroaryl containing one, two, or three heteroatoms selected from oxygen, sulfur, and nitrogen, wherein the C₅₋₁₀-heteroaryl is optionally substituted by one or more groups selected from phenyl, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, CF₃, CN, CONH₂, CONMe₂, CONEt₂, COOH, COOMe, COOEt, COH, COMe, OH, OMe, OEt, F, Cl, Br, SO₂Me, SONH₂, SONMe₂, NO₂, NH₂, NMe₂, NEt₂, and NPr₂, or

R³ is phenyl optionally substituted by a C₃₋₁₀-heterocycle containing one or two heteroatoms selected from oxygen and nitrogen, wherein the C₃₋₁₀-heterocycle is optionally substituted by one or more groups selected from C₁₋₄-alkyl and oxo;

R⁴ is H, F, or Cl; and

R⁵ is methyl, ethyl, cyclopropyl, cyclobutyl, CH₂CF₃, or benzyl optionally substituted by F,

or a pharmacologically acceptable salt, enantiomer, hydrate, or solvate thereof, or deuterated form thereof.

16. A pharmaceutical formulation comprising a compound according to one of claims 1 to 15 and a pharmaceutically acceptable excipient.

* * * * *