

FACILE SYNTHESIS OF AZIDES: CONVERSION OF HYDRAZINES
USING DINITROGEN TETROXIDE


Yong Hae Kim, Kweon Kim, and Sung Bo Shim









Department of Chemistry, The Korea Advanced Institute of
Science and Technology, P.O. Box 150, Chong-Yang, Seoul, Korea

Abstract : Various hydrazines such as aryl-, carbonyl-, and sulfonyl-hydrazine were reacted with dinitrogen tetroxide to give the corresponding azides in excellent yields under mild conditions at low temperature (-20~-40 °C) in acetonitrile.

Intensive studies on organic syntheses¹ using organic azides have been reported for the important preparation of amines^{1,2}, azo compounds^{1,3}, azomethines^{1,3}, iminophosphoranes⁴, iminosulphuranes⁵, isocyanates⁶, and aziridines⁷. Azides are essential intermediates in organic synthesis and have been prepared by substitution on carbon of sodium azide⁸, trimethyl silyl azide⁹ or by nitrosation of hydrazine derivatives¹⁰ using nitrous acid¹¹, nitrosyl chloride¹², organic nitrites¹³ and ferric nitrate¹⁴. Previously we reported that dinitrogen tetroxide is a good nitrosation reagent for activating thiols to convert into the thionitrites¹⁵. We have now found that various hydrazines reacted rapidly with dinitrogen tetroxide¹⁶ to give the corresponding azides in mostly quantitative yields under mild conditions at low temperature (-20~-40 °C) in acetonitrile. In a typical procedure, dinitrogen tetroxide solution (2 mmole, CCl₄: 300 μ l) was slowly added to the benzoyl hydrazide (0.138g, 1 mmole, anhydrous CH₃CN : 5ml) at -10 °C with stirring. The reaction was monitored by t.l.c. (Merck; Kiesegel 60F₂₅₄, CHCl₃). After being stirred for 10 minutes, 25 ml of ether was added to the reaction mixture and this ether layer was twice washed with 10 ml of saturated sodium bicarbonate solution and dried over anhydrous magnesium sulfate. Evaporation of the solvent gave the highly pure azide (0.141g, 96%), which were further purified by column chromatography (Silicagel; Wako gel 300, ether:n-hexane=1:1, 0.133g, 91%). All the products thus obtained were identified by comparing their ir, ¹H nmr spectra and melting points with those from the authentic samples. The results obtained are summarized in Table 1.

Table 1. Conversion of Hydrazines into Azides




Run	R	Reactn. Temp.(°C)	Reactn. time(min.)	Yield(%) ^a	Ref. No ^b
1		-30	10	91 ^c	14
2		-20	10	95 ^c	14
3		-20	20	87	14
4		-25	20	86	20
5		-10	10	92	21
6		-30	10	89	22
7		-40	20	84	22
8		-30	10	90	23
9		-30	10	95	24

a : Isolated and purified yield.

b : References for the known products of azides are cited in the reference section.

c : When the reaction was carried out at ca. 0 °C, the yield of azide was about the same.

N_2O_4 is known to be in equilibrium with NO^+ and NO_3^- in solution by its self ionization¹⁸. Probably the electrophilic attack of NO^+ on the nitrogen atom of the amine forms the corresponding β -nitroso hydrazine intermediate (2) which may finally convert into the products, azides(3). Nitrosation of secondary amines with dinitrogen tetroxide are well known by isolation of the stable secondary nitroso compounds¹⁹.



The yields of the azides are markedly high when the reaction temperatures are controlled at lower than $-20^\circ C$. While, p-nitrophenyl- and phenyl azides appear to rearrange to the heterocycles¹⁴ due to their instability at high temperature. Our method can be carried out in the wide range of low reaction temperature ($-10\sim-40^\circ C$) for short reaction time (10-20 min) and is simple for work up after the completion of the reaction. Thus, this facile method seems to be widely available for the preparation of various azides from hydrazines.

Acknowledgment

This work was supported by KAIST and KOSEF grant.

References

- 1) T. Sheradsky, Ch 6 in *The Chemistry of the Azido Group*, S. Patai ed., Interscience-Wiley, London, 331-395, 1971.
- 2) E.J. Corey, K.C. Nicolaou, R.D. Balanson, and Y. Machida, *Synthesis*, 590, 1975.
- 3) H. Bayley, D.N. Standring, and J.R. Knowles, *Tetrahedron Lett.*, **19**, 3633, 1978.
- 4) H. Staudinger, and J. Meyer, *Helv. Chim. Acta.*, **2**, 635, 1979.
- 5) L. Horner, and A. Christmann. *Chem. Ber.*, **96**, 388, 1963.
- 6) P.A.S. Smith, *Org. Reactions*, **3**, 337-449, 1946.
- 7) Y. Ittah, Y. Sasson, I Shahak, S. Tsaroom, and T. Blum, *J. Org. Chem.*, **43**, 4271, 1978.
- 8) H. Kuzuhara, and S. Emoto, *Tetrahedron Lett.*, 5051, 1973.
- 9) J.H. MacMillan, and S.S. Washburne, *J. Org. Chem.*, **38**, 2982, 1973; H.R. Kricheldorf, *Chem. Ber.*, 105, 3958, 1972.
- 10) J. Honzl, and R. Rudinger, *Coll. Czech. Chem. Comm.*, **26**, 2333, 1961.

- 11) H. Roesky, and D. Glemser, *Chem. Ber.*, **97**, 1710, 1964.
- 12) H. Neunhoffer, G. Cunny, and W.K. Franke, *Ann. Chem.*, **96**, 713, 1968.
- 13) N.A. Smart, G.T. Young, and M.W. Williams, *J. Chem. Soc.*, 3902, 1960.
- 14) P. Laszlo, and E. Polla, *Tetrahedron Lett.*, **25**, 3701, 1984.
- 15) S. Oae, D. Fukushima, and Y.H. Kim, *J. Chem. Soc.*, 407, 1977; Y.H. Kim, K. Shinhama, D. Fukushima, and S. Oae, *Tetrahedron Lett.*, 1212, 1978.
- 16) N_2O_4 is commercially available or can be easily prepared¹⁷. N_2O_4 gas was trapped in a cold flask containing P_2O_5 at -20°C . The liquid N_2O_4 was trapped in cold CCl_4 , and this solution titrated was used for the reaction.
- 17) A. Pedler, and F.H. Pollard, "Inorganic Synthesis", Vol. V, 87, McGraw-Hill, New York, 1957.
- 18) W.R. Angus, R.W. Jones, and G.O. Fiips, *Nature*, **164**, 433, 1947; K. Clusius, M. Vecchi, *Helv. Chim. Acta.*, **36**, 930, 1953.
- 19) F. Wudl, and T.B.K. Lee, *J. Amer. Chem. Soc.*, **93**, 271, 1971; D.H.R. Barton, and S.C. Narang, *J. Chem. Soc., Perkin I.*, 1114, 1977.
- 20) R. Kuhn, and G. Kruger, *Chem. Ber.*, **89**, 473, 1956.
- 21) S. Munekata, and S. Kikuchi, *Seisan-Kenkyu*, **20**(10), 519, 1968.
- 22) Y. Yukawa and Y. Tsuno, *J. Amer. Chem. Soc.*, **79**, 5530, 1957.
- 23) A.S. Bailey and J.R. Case, *Tetrahedron*, **3**, 113, 1958.
- 24) E. Lieber, T.S. Cao, and C.N.R. Rao, *J. Org. Chem.*, **22** 654, 1957.

(Received in Japan 19 June 1986)