

Oxone/Sodium Chloride: A Simple and Efficient Catalytic System for the Oxidation of Alcohols to Symmetric Esters and Ketones

Agnes Schulze, Georgia Pagona, and Athanassios Giannis
Institut für Organische Chemie, Universität Leipzig, Leipzig, Germany

Abstract: An efficient method for the oxidation of alcohols is presented. The use of catalytic amounts of sodium chloride in combination with oxone allows the conversion primary aliphatic alcohols to symmetric esters. Secondary alcohols can be easily oxidized to ketones, and benzylic alcohols are converted to the corresponding aldehydes. The method is cost effective and environmentally benign.

Keywords: Alcohols, aldehydes, catalytic oxidation, esterification, ketones

Oxidation of primary and secondary alcohols to aldehydes or carboxylic acids and ketones, respectively, is a fundamental process in organic chemistry, and several reagents have been developed for this purpose.^[1] However, many of the oxidants were subsequently abolished for environmental reasons. Instead, enzymatic^[2] and particularly metal-free^[3] oxidations are now of great interest, and a variety of valuable methods have been developed. To the last category belong, for example, oxidations mediated by hypervalent iodine reagents such as IBX^[4,5] or TEMPO^[1] in combination with NaClO,^[6] Br₂,^[7] I₂,^[8] or Cl₂.^[9] TEMPO has been also used in combination with oxone and *n*-Bu₄NBr as a phase-transfer catalyst.^[10]

Received in Poland July 28, 2005

Address correspondence to Athanassios Giannis, Institut für Organische Chemie, Universität Leipzig, Johannisallee 29, 04103 Leipzig, Germany. Fax: +49/341 97 36 599; E-mail: giannis@chemie.uni-leipzig.de

A shortcoming of the TEMPO-mediated oxidations is the fact that one equivalent sodium chloride is generated per molecule of alcohol oxidized. Other shortcomings are the use of sodium bromide as a cocatalyst and chlorinated solvents. Finally, although TEMPO is used catalytically, it is expensive, and efficient recycling is an important issue. Several approaches have been developed^[1d] to circumvent these problems.

Because of their importance, several methods for the conversion of primary alcohols to symmetric esters have been developed.^[11] Symmetric esters are of considerable interest as ingredients for cosmetic, dermatological, and pharmaceutical preparations. A search in SciFinder revealed more than 300 patents for this class of compounds. They are furthermore utilized for the production of denture adhesives and stable aqueous surfactant compositions, in transdermal drug-delivery systems, as lubricants, and as mold-release agents for polyurethanes. Esters of branched-chain carboxylic acids with branched-chain alcohols have been described as antibacterial, antimycotic, and antiparasitic substances and are used in skin preparations.^[12]

Herein, we present a catalytic method for the oxidation of alcohols with NaCl/oxone. Using this system, primary aliphatic alcohols are converted to symmetric esters, and secondary alcohols can be easily oxidized to ketones. Furthermore, benzylic alcohols are converted to the corresponding aldehydes.

We found that the combination of oxone (1 equiv.) and *n*-Bu₄NBr (0.1 equiv.) in a biphasic system (EtOAc/H₂O) effectively converts 1-octanol (1 equiv.) to the corresponding symmetric ester. Importantly, employing the same method and KBr (0.1 equiv.) instead of *n*-Bu₄NBr afforded also octyl octanoate in comparable yield. The later experiment was a clear indication that 1) tetra-*n*-butylammonium oxone^[13] was not the actual oxidant and 2) the use of a phase-transfer catalyst was not necessary.

In fact, the use of *n*-Bu₄NHSO₄ and oxone or oxone alone did not induce conversion of 1-octanol to octyl octanoate (or to octanal or octanoic acid) even after prolonged reaction time.













Encouraged by these results, we next investigated the oxone/NaCl (0.05–0.1 equiv.) system and were pleased to realize that it was more effective in terms of yield and reaction time than the combination of oxone with KBr or with *n*-Bu₄NBr. With this reagent combination, a variety of aliphatic alcohols could be converted to the corresponding symmetric esters fast and at room temperature. In several cases formation of the corresponding carboxylic acid was also observed (see Table 1 and Scheme 1). The use of KI does not lead to any oxidation products.

In contrast to primary aliphatic alcohols, benzylic alcohols were oxidized to the corresponding aldehydes (Scheme 2) in moderate yields. In these cases we observed that the use of 0.5–1 equiv. of NaCl significantly increased the yield.

Finally, secondary alcohols were oxidized to the corresponding ketones (Scheme 3) in high yields.









However, our method is not compatible with double and triple bonds.











Table 1. Catalytic oxidation of primary and secondary alcohols with NaCl/oxone

Entry	Substrate	Reaction time (h)	Product	Ratio E:A ^a	Total yield (%) ^b
1		5		6:1	98
2		5.5		4:1	95
3		4.5		3.2:1	79
4		4.5		1:0	85
5		3.5		1:0	90
6		5		2.2:1	93

(continued)

Table 1. Continued

Entry	Substrate	Reaction time (h)	Product	Ratio E:A ^a	Total yield (%) ^b
7		5		1.1:1	68
8		5		—	60 ^c
9		7		—	30 ^c
10		2		—	85 ^c

11		3.5		—	85
12		4		—	80
13		1.5		—	90 ^c
14		1		—	94
15		4		—	80 ^c


^aE: ester, A: acid.

^bYield of isolated product.

^cThe yield was determined by GC/MS.

Oxone/Sodium Chloride

1151




Scheme 1. Oxidation of primary aliphatic alcohols.

Oxone has been applied in the past for several oxidations. For example, at elevated temperature and in combination with concentrated sulfuric acid or with acetic acid/ethanol, it has been used for the conversion of alkenes to diols, phenols to quinones, pyridine to pyridin-1-oxide, sulfides to sulfoxides, and ketones to esters or to lactones.^[14] Recently, oxone has also been utilized for the oxidation of aromatic and aliphatic aldehydes to the corresponding carboxylic acids or esters.^[15] Furthermore, oxone is the reagent of choice for the synthesis of dimethyldioxirane^[16] and IBX.^[14a]


It is known that treatment of NaCl or NaBr with oxone generates chlorine and bromine respectively.^[17] This system has been used for chlorinations and brominations of α,β -unsaturated carbonyl compounds or for the conversion of oximes to gem-chloro-nitro derivatives.^[18] The combination of NaBr/oxone was also reported to oxidize benzylic alcohols to aldehydes, secondary alcohols to ketones, and butanol to the symmetric ester. An important shortcoming of this method is the bromination of the aromatic system and the need for an excess of NaBr.^[19]

The oxidation of primary alcohols to symmetric esters is described in Scheme 4: First, the aldehyde is formed, which is then attacked by the parent alcohol to form the corresponding hemiacetal. Oxidation of the latter affords the symmetric ester. The isolation of aromatic aldehydes (see Table 1, entries 8–10) supports this suggestion. They do not form symmetric esters probably because their carbonyl group is not sufficiently electrophilic^[20] to allow a nucleophilic attack by a second molecule of the aromatic alcohol present in the reaction mixture. Attempts to convert primary alcohols to tert. butylesters by performing the oxidations in the presence of tert. butanol have been unsuccessful.

In conclusion we presented a new, efficient, and fast method for the conversion of primary aliphatic alcohols to symmetric esters as well for the oxidation of secondary alcohols to ketones. Our method is cost-effective and environmentally benign. In fact, only catalytic amounts of sodium chloride are used, whereas oxone is inexpensive, stable, easy to transport, and nontoxic. The resulting inorganic by-products are also nonpolluting.



Scheme 2. Oxidation of benzylic alcohols.



Scheme 3. Oxidation of secondary alcohols.

EXPERIMENTAL


Reagents and solvents were commercially available and were used without further purification. ^1H and ^{13}C NMR spectra were recorded on a Varian Gemini 200 (200 MHz for ^1H NMR; 50 MHz for ^{13}C NMR) and Varian Gemini 2000 (200 MHz for ^1H NMR; 50 MHz for ^{13}C NMR). Chemical shifts were reported in the scale relative to the solvent used as an internal reference. HRMS was obtained on a Bruker Daltonics APEX II (for ESI). The reaction progress was monitored with a Varian Saturn 2200 GC/MS. Flash column chromatography was performed on Merck silica gel 60 (0.040–0.063 mm).

Typical Experimental Procedure

Oxone (1 mmol), NaCl (0.1 mmol), and finally H_2O (0.5 ml) were added to a solution of the alcohol (1 mmol) in EtOAc (2 ml). The reaction mixture was stirred at room temperature for 1–7 h depending on the alcohol (see Table 1). After completion of the reaction, the mixture was diluted with H_2O (5 ml) and EtOAc (3 ml), and the layers were separated. The aqueous layer was extracted with EtOAc (3×5 ml), and the combined organic layers were dried over Na_2SO_4 . The solvent was removed under reduced pressure to give the desired product. Analytical pure products were obtained by column chromatography.

6-Chlorohexyl 6-chlorohexanoate (3): oil; ^1H NMR (CDCl_3 , 200 MHz): $\delta = 1.41\text{--}1.82$ (m, 14H), 2.28–2.37 (m, 2H), 3.54 (t, 4H, $J = 6.6$ Hz), 4.07 (t, $J = 6.6$ Hz, 2H); ^{13}C NMR (CDCl_3 , 50 MHz): $\delta = 24.4, 25.4, 26.5, 26.6, 28.6, 32.4, 32.6, 34.2, 44.8, 44.9, 64.4, 173.7$; HR ESI-MS: calcd. for $(\text{M} + \text{Na})^+$: 291.08891; found: 291.08935.

2-Ethylbutyl 2-ethylbutanoate (4): oil; ^1H NMR (CDCl_3 , 200 MHz): $\delta = 0.89$ (t, 12H, $J = 7.4$ Hz), 1.29–1.66 (m, 9H), 2.13–2.28 (m, 1H), 4.01 (d, 2H, $J = 5.5$ Hz); analytical data were consistent with those reported in the literature.^[21]



Scheme 4. Oxidation of primary alcohols to symmetric esters by NaCl/oxone.

2-Ethylhexyl 2-ethylhexanoate (5): oil; ^1H NMR (CDCl_3 , 200 MHz): $\delta = 0.85\text{--}0.93$ (m, 12H), 1.19–1.66 (m, 17H), 2.19–2.32 (m, 1H), 4.07 (d, 2H, $J = 5.6$ Hz); analytical data were consistent with those reported in the literature.^[11c]

Compound (6): oil; ^1H NMR (CDCl_3 , 200 MHz): $\delta = 0.88\text{--}1.00$ (m, 18H), 1.11–1.72 (m, 17H), 2.08–2.41 (m, 3H), 4.11–4.16 (m, 2H); analytical data were consistent with those reported in the literature.^[22]

Cyclohexylmethyl cyclohexylcarboxylate (7): oil; ^1H NMR (CDCl_3 , 200 MHz): $\delta = 0.98\text{--}1.93$ (m, 21H), 2.21–2.40 (m, 1H), 3.87 (d, 2H, $J = 6.4$ Hz); analytical data were consistent with those reported in the literature.^[23]

Compounds **1**, **2**, **3**, **8**, **9**, **10**, **11**, **12**, **13**, **14**, and **15** are commercially available.

ACKNOWLEDGMENTS

A. Schulze is grateful for grants from the Graduiertenkolleg Mechanistische und Anwendungsaspekte nicht-konventioneller Oxidationsreaktionen. G. Pagona is grateful for a grant from European Union under the auspices of the Socrates Exchange Program.

REFERENCES

1. (a) Sheldon, R. A.; Koshi, J. K. *Metal-Catalyzed Oxidations of Organic Compounds*; Academic Press: New York, 1984; (b) Hudlicky, M. *Oxidations in Organic Chemistry*; ACS Monograph: Washington, 1990; p. 186; (c) Armstrong, A. Synthetic methods. Part 2, oxidation and reduction methods, annual reports on the progress of chemistry, section B. *Org. Chem.* **2003**, *99*, 21–48; (d) Sheldon, R. A.; Arends, I. W. C. E.; ten Brink, G. J.; Dijkstra, A. Green, catalytic oxidation of alcohols. *Acc. Chem. Res.* **2002**, *35*, 774–781; (e) de Nooy, A. E. J.; Besemer, A. C.; Bekkum, H. V. On the use of stable organic nitroxyl radicals for the oxidation of primary and secondary alcohols. *Synthesis* **1996**, 1153–1174.
2. Kroutil, W.; Mang, H.; Edegger, K.; Faber, K. Biocatalytic oxidation of primary and secondary alcohols. *Adv. Synth. Catal.* **2004**, *346*, 125–142.
3. Adam, W.; Saha-Möllner, C. R.; Ganeshpure, P. A. Synthetic applications of nonmetal catalysts for homogenous oxidations. *Chem. Rev.* **2001**, *101*, 3499–3548.
4. (a) Frigerio, M.; Santagostino, M.; Sputore, S. A user-friendly entry to 2-iodoxybenzoic acid (IBX). *J. Org. Chem.* **1999**, *64*, 4537–4538; (b) Frigerio, M.; Santagostino, M.; Sputore, S.; Palmisano, G. Oxidation of alcohols with o-iodoxybenzoic acid in DMSO: A new insight into an old hypervalent iodine reagent. *J. Org. Chem.* **1995**, *60*, 7272–7276.
5. (a) Tohma, H.; Kita, Y. Hypervalent iodine reagents for the oxidation of alcohols and their application to complex molecule synthesis. *Adv. Synth. Catal.* **2004**, *346*,

- 111–124; (b) Zhdankin, V. V.; Stang, P. J. Recent developments in the chemistry of polyvalent iodine compounds. *Chem. Rev.* **2002**, *102*, 2523–2584; (c) Varvoglis, A. *Hypervalent Iodine in Organic Synthesis*; Academic Press: San Diego, 1997.
- (a) Zhao, M.; Li, J.; Mano, E.; Song, Z.; Tschaen, D. M.; Grabowski, E. J. J.; Reider, P. J. Oxidation of primary alcohols to carboxylic acids with sodium chlorite catalyzed by TEMPO and bleach. *J. Org. Chem.* **1999**, *64*, 2564–2566; (b) Anelli, P. L.; Biffi, C.; Montanari, F.; Quici, S. Fast and selective oxidation of primary alcohols to aldehydes or to carboxylic acids and of secondary alcohols to ketones mediated by oxoammonium salts under two-phase conditions. *J. Org. Chem.* **1987**, *52*, 2559–2562.
 - Liu, R.; Liang, X.; Dong, C.; Hu, X. Transition-metal-free: A highly efficient catalytic aerobic alcohol oxidation process. *J. Am. Chem. Soc.* **2004**, *126*, 4112–4113.
 - Miller, R. A.; Hoerner, R. S. Iodine as a chemoselective reoxidant of TEMPO: Application to the oxidation of alcohols to aldehydes and ketones. *Org. Lett.* **2003**, *5*, 285–287.
 - Bjorsvik, H. R.; Liguori, L.; Constantino, F.; Minisci, F. A new modified “Montanari oxidation process” by means of chlorine dissolved in the reaction solvent as oxidant and TEMPO as catalyst: Oxidation of 3-*S*-quinuclidinol to 3-quinuclidinone. *Org. Process Res. Dev.* **2002**, *6*, 197–200.
 - Bolm, C.; Magnus, A. S.; Hildebrand, J. P. Catalytic synthesis of aldehydes and ketones under mild conditions using TEMPO/oxone. *Org. Lett.* **2000**, *2*, 1173–1175.
 - (a) Amati, A.; Dosualdo, G.; Zhao, L.; Bravo, A.; Fontana, F.; Minisci, F.; Bjorsvik, H. R. Catalytic processes of oxidation by hydrogen peroxide in the presence of Br₂ or HBr. Mechanism and synthetic applications. *Org. Process Res. Dev.* **1998**, *2*, 261–269; (b) Jenzer, G.; Schneider, M. S.; Wandeler, R.; Mallat, T.; Baiker, A. Palladium-catalyzed oxidation of octyl alcohols in “supercritical” carbon dioxide. *J. Catal.* **2001**, *199*, 141–148; (c) Simpura, I.; Nevainen, V. Symmetric esters by Tschtschenko reaction of aldehydes catalyzed by bi- and tridentate catalysts derived from catechol or gallol, trimethylaluminum and isopropanol. *Tetrahedron* **2001**, *57*, 9867–9872; (d) Sonnet, P. E.; Moore, G. G. Esterifications of 1- and rac-2-octanols with selected acids and acid derivatives using lipases. *Lipids* **1988**, *23*, 955–960; (e) Tamaru, Y.; Yamada, Y.; Inoue, K.; Yamamoto, Y.; Yoshida, Z. Oxidation of primary and secondary alcohols by the catalysis of palladium. *J. Org. Chem.* **1983**, *48*, 1286–1292.
 - Schreiber, J.; Juestel, C.; Wolf, F. Ger. Offen. (1999), DE 19742275; CAN 130:242372.
 - Trost, B. M.; Braslau, R. Tetra-*n*-butylammonium oxone. Oxidations under anhydrous conditions. *J. Org. Chem.* **1988**, *53*, 532–537.
 - Kennedy, R. J.; Stock, A. M. The oxidation of organic substances by potassium peroxymonosulfate. *J. Org. Chem.* **1960**, *25*, 1901–1906.
 - (a) Travis, B.; Sivakumar, M.; Hollist, G. O.; Borhan, B. Facile oxidation of aldehydes to acids and esters with oxone. *Org. Lett.* **2003**, *5*, 1031–1034; (b) Webb, K. S.; Ruskay, S. J. Oxidation of aldehydes with oxone in aqueous acetone. *Tetrahedron* **1998**, *54*, 401–410.
 - Murray, R. W.; Jeyaraman, R. Dioxiranes: Synthesis and reactions of methyldioxiranes. *J. Org. Chem.* **1985**, *50*, 2847–2853.

17. Dieter, R. K.; Nice, L. E.; Velu, S. E. Oxidation of α,β -enones and alkenes with oxone and sodium halides: A convenient laboratory preparation of chlorine and bromine. *Tetrahedron Lett.* **1996**, *37*, 2377–2380.
18. Ceccherelli, P.; Curini, M.; Epifano, F.; Marcotullio, M. C.; Rosati, O. One-step conversion of oximes to gem-chloro-nitro derivatives. *Tetrahedron Lett.* **1998**, *39*, 4385–4386.
19. Koo, B.-S.; Lee, C. K.; Lee, K.-J. Oxidation of benzyl alcohols with oxone and sodium bromide. *Synth. Commun.* **2002**, *32*, 2115–2123.
20. Mazitschek, R.; Mülbaier, M.; Giannis, A. IBX-vermittelte oxidation von primären alkoholen und aldehyden zu carbonsäuren. *Angew. Chem.* **2002**, *114*, 4216–4218; IBX-mediated oxidation of primary alcohols and aldehydes to form carboxylic acids. *Angew. Chem., Int. Ed.* **2002**, *41*, 4059–4061.
21. Kajigaeshi, S.; Kawamukai, H.; Fujisaki, S. Oxidation using quaternary ammonium polyhalides. III. An effective oxidation of alcohols and ethers by the use of benzyltrimethylammonium tribromide. *Bull. Chem. Soci. Jpn.* **1989**, *62*, 2585–2588.
22. Trost, B. M.; Masuyama, Y. Molybdenum catalyzed reactions. Selectivity in oxidations with hydrogen peroxide and ammonium molybdate. *Isr. J. Chem.* **1984**, *24*, 134–143.
23. Ooi, T.; Miura, T.; Itagaki, Y.; Ichikawa, H.; Maruoka, K. Catalytic Meerwein–Ponndorf–Verley (MPV) and Oppenauer (OPP) reactions: Remarkable acceleration of the hydride transfer by powerful bidentate aluminium alkoxides. *Synthesis* **2002**, *2*, 279–291.