

REACTIONS OF NITRILES WITH HYDROGEN HALIDES AND NUCLEOPHILIC REAGENTS

E. N. Zil'berman

CONTENTS

I. Introduction	615
II. Reactions of nitriles with hydrogen halides	615
A. Nitrile hydrohalides of composition $\text{RCN} \cdot n\text{HX}$	615
B. Some properties of solutions of hydrogen halides in nitriles	617
C. Nitrile hydrohalides of composition $2\text{RCN} \cdot n\text{HX}$	618
D. Preparation of 1,3,5-triazines	618
E. Hydrogen cyanide hydrohalides	619
III. Reactions of nitriles with hydrogen halides and nucleophilic reagents	619
A. Reactions with water	619
B. Reactions with alcohols	621
C. Reactions with organic acids	624
D. Reactions with carbonyl compounds	625
E. Structure and chemical properties of amide hydrohalides	625
F. Selective reactivity of one cyano-group in adiponitrile	626
G. Reactions with amines	627
H. The Gattermann reaction	627
J. The Hoesch reaction	628
K. Reaction of dimeric nitrile salts ($2\text{RCN} \cdot n\text{HX}$) with hydroxy-compounds	628
IV. Reduction of nitriles in the presence of hydrogen chloride	629
A. Preparation of aldehydes by the Stephen method	629
B. Preparation of bisamides in the reduction of nitriles by the Stephen method	629
C. Preparation of amines	630

I. INTRODUCTION

The ability of nitriles to form addition compounds with nucleophilic and electrophilic reagents derives from the electronic structure of the triple carbon-nitrogen bond¹. We have recently reviewed the reactions of nitriles with electrophilic reagents¹, and these will be considered only to a limited extent. Because of the reduced electron density at the carbon atom of the cyano-group, nitriles should react very readily with nucleophilic reagents. Compounds such as hydrogen sulphide and hydroxylamine do in fact add to nitriles to form thioamides and amidoximes, but in general nitriles are very stable towards many other nucleophilic reagents (water, alcohols, amines, etc.). The reactivity of nitriles is increased and the range of their reactions with nucleophilic reagents broadened in the presence of acidic or alkaline compounds, the action of which is

evidently due to the ready polarisability of the cyano-group; also the presence of acids creates suitable conditions for the appearance of electrophilic activity at the nitrile carbon atom and the formation of new bonds between this atom and atoms of oxygen, sulphur, nitrogen, carbon, etc.

Hydrogen halides are particularly important as acid catalysts for reactions of the cyano-group, and in many cases, especially with hydrogen chloride, they are specific catalysts. The Pinner reaction (preparation of imidate salts), Hoesch reaction (preparation of ketimine salts), Gattermann reaction, Stephen reaction (preparation of aldimine salts), and several other acid-catalysed reactions of nitriles are carried out in the presence of hydrogen halides, and do not occur with other protogenic or Lewis acids in the absence of hydrogen halides. In the literature these reactions are very often discussed without consideration of the nature of the products of the reaction of nitriles with hydrogen halides, which may be formed under the experimental conditions. These products are often formulated as imidoyl halides, although these have never been isolated and there is no proof of their existence in the reaction medium. There have been no attempts in the review literature to generalise the existing data on the reactions of nitriles with hydrogen halides in the absence of other reagents.

This review is intended to fill the existing gaps. We shall consider the structure of the hydrohalogenation products of nitriles and hydrogen cyanide, and describe the reactions of nitriles with hydrogen halides and other reagents: hydroxylic compounds (water, alcohols, and organic acids), aldehydes and ketones, amines and aromatic compounds (the Gattermann and Hoesch reactions). The reduction of nitriles in the presence of hydrogen halides and other matters related to these reactions of nitriles will also be considered.


II. REACTIONS OF NITRILES WITH HYDROGEN HALIDES

A. Nitrile Hydrohalides of Composition $\text{RCN} \cdot n\text{HX}$


Nitrile hydrochlorination products were first isolated in 1866 by Gautier²⁻⁴, and the hydrohalogenation of nitriles was much studied in the last century⁵⁻¹⁰. Several nitrile hydrohalides were apparently obtained in admixture with other compounds in the bromination of aliphatic nitriles^{11,12}. Until 1931 it was thought that one or two hydrogen halide molecules could add to nitriles in the cold, to form imidoyl halides and amino-halides respectively, but, after more careful study of the composition of the very unstable reaction products of nitriles and hydrogen halides, Hantzsch concluded¹³ that the monohydrochlorides were not obtained, and that only the saltlike dihydrochlorides exist. By hydrobromination, however, Hantzsch obtained mono- and di-hydrobromides. Hantzsch's experiments on hydrobromination were later repeated by Klages and Grill¹⁴, who showed that Hantzsch's results on nitrile monohydrobromides were in error. In 1957 it was reported¹⁵ that treatment of *N*-acylaminoacetonitriles with hydrogen chloride in the cold produces monohydrochlorides, and these were stated to be the first isolated imidoyl chlorides or nitrilium salts; however, hydrogen chloride can add not only to the cyano-group, but also to the amide group, and this possibility was not considered. Moreover, the hydrochlorides obtained melted above 100°, and it is improbable that imidoyl halides would melt at such high temperatures. Up

to the present, therefore, only nitrile dihydrohalides have been prepared (textbooks¹⁶ and monographs¹⁷ still erroneously refer to the existence of solid monohydrohalogenated nitriles).

The 1:2 reaction products of nitriles and hydrogen halides are also obtained by treating carboxylic acid amides with PCl_5 ;¹⁸ these products have been shown to be identical with those obtained by direct hydrochlorination of nitriles¹³. Kirsanov has studied the reaction of carboxylic acid amides with PCl_5 ,¹⁹⁻²¹ and his data confirm the earlier suggestion²² that the primary reaction products are trichlorophosphazacyl compounds:




These can be obtained in high yields if the hydrogen chloride is removed from the reaction zone. In the presence of the hydrogen chloride they decompose into nitriles and POCl_3 :



The resulting nitriles react further with the hydrogen chloride in the normal fashion, *i.e.* to form the 1:2 addition products.

The 1:2 addition products of nitriles with hydrogen halides have been given various structural formulae. In the older literature they were considered to be amino-halides (A).²³ In 1931 Hantzsch¹³ suggested the formula (B), according to which they are hydrochlorides of nitrilium salts, and this formula is met with today¹⁶. Klages and Grill found¹⁴ that the properties of the nitrile dihydrohalides correspond best to those of imidoil halide salts²⁴, *i.e.* halogenoiminium halides (C). Finally, Janz has suggested²⁵ that the nitrile hydrochlorides may have formula (D):




The nitrile dihydrohalides show a high conductivity in liquid sulphur dioxide¹⁴ and are therefore salts and cannot be amino-halides (A). The nature of the bond between the nitrilium salt and the hydrogen halide molecule in formula (B) is not clear. There can be no suggestion of the formation of a hydrogen bond between the hydrogen halide and the cation, since the nitrilium ion has no free electron pair to act as a base¹⁴. Formula (B) cannot therefore represent the true structure of the nitrile dihydrohalides. Unlike formula (B), the other nitrilium salt structure (D) is quite possible, since the existence of the dihalide anion HX_2^- has been proved²⁶⁻³¹. It has been suggested²⁶ that the stability of the anion HCl_2^- , by analogy with HF_2^- , is due to the formation of a hydrogen bridge between the hydrogen atom in the HCl and the chloride ion ($\text{Cl}-\text{H}\cdots\text{Cl}^-$). There are also no particular objections to formula (C), representing a halogenoiminium structure for the salts $\text{RCN}\cdot 2\text{HX}$.

The infra-red spectra of acetonitrile dihydrobromide have shown it to be acetobromoiminium bromide²⁵; an attempt was also made to obtain the infra-red spectra of solid acetonitrile dihydrochloride, but was unsuccessful because of its instability. The infra-red spectra of very dilute solutions (up to 1.15 M) of hydrogen chloride in acetonitrile revealed no band due to a C-Cl bond (750 cm^{-1})

but there was a band at 1180 cm^{-1} characteristic of the HCl_2^- anion²⁷. It was therefore concluded that for the dihydrochloride, unlike the dihydrobromide or dihydroiodide, the most probable formula is (D). Janz did not consider this structure as proved, because (a) the structure of nitrile dihydrochlorides may be different in dilute solution than in the solid state, and (b) the measurements were made²⁵ at room temperature, while even at $+6^\circ$ acetonitrile dihydrochloride dissociates into the nitrile and hydrogen chloride³².


The existing data are therefore insufficient to ascribe to the dihydrochlorides a structure different from the structure of the other dihydrohalides. In order to establish finally the structure of nitrile dihydrochlorides, their spectra must be investigated under conditions such that their conversion into other nitrile-hydrogen chloride reaction products is excluded.

It is of interest that no difference has been found in the structures of solid cyanamide dihydrochloride and dihydrobromide³³ and both salts have been given the same halogenoformaminidinium halide structure:



The dihydrohalides described in the literature, which are in all probability of structure (C), include the dihydrochlorides of acetonitrile³², bromo- and fluoro-acetonitrile³⁴, benzonitrile⁸, *p*-tolunitrile¹³, and *p*-methoxybenzonitrile³⁵, the dihydrobromides of acetonitrile²⁵, bromoacetonitrile³⁴, phenylacetonitrile and benzonitrile¹⁴, and *p*-tolunitrile¹³, and the dihydroiodides of acetonitrile²⁵, phenylacetonitrile, *o*-, *m*-, and *p*-tolunitrile, *o*-, *m*-, and *p*-nitrobenzonitrile, succinonitrile, terephthalonitrile, etc.¹⁰

Of the acetonitrile dihydrohalides, the hydriodide melts at the highest temperature (above 100°); the hydrobromide and hydrochloride melt at $75^\circ-80^\circ$ and about 20° respectively^{25,36}. The thermal stability of the salts (ease of formation) also increases in this order:



The stability of the halogenoiminium halides is related to the polarisability of the hydrogen halides³⁶: the least polarisable hydrogen chloride forms the least stable salts, and the most polarisable hydrogen iodide the most stable salts.


On heating, the halogenoiminium halides are usually decomposed into the nitriles and the hydrogen halides. The vapour pressure of certain bromoiminium bromides has been shown to depend on the temperature¹⁴. The heat of dissociation into the initial components was calculated; for benzobromoiminium bromide $\Delta H = 30 \text{ kcal per mole}$.

Nitrile dihydrochlorides enter into exchange reactions with other anions, and form stable salts with complex anions¹⁴. The following have been described: benzochloroiminium and acetochloroiminium hexachlorostannates, and benzochloroiminium, *p*-nitrobenzochloroiminium, and acetochloroiminium hexachloroantimonates³⁷.

Certain nitrogen-unsubstituted nitrilium salts with complex anions have been prepared where the corresponding halides are unknown. Benzo- and aceto-nitrilium hexachlorostannates and *p*-nitrobenzo- and aceto-nitrilium hexachloroantimonates have been prepared³⁷. The reaction between benzonitrile, hydrogen chloride, and zinc chloride gave benzonitrilium trichlorozincate³⁸. Their conductivity in liquid sulphur dioxide confirms their salt structure³⁷.

According to Klages³⁷, which complex is formed depends on the reaction temperature. There is no reaction between a nitrile, the complex-forming reagent, and hydrogen chloride at a very low temperature; in the intermediate region nitrilium salts are formed, and at higher temperatures two hydrogen chloride molecules react, with the formation of imonium salts. In some cases there is no intermediate region, *i.e.* nitrilium salts could not be obtained.

On heating, the complex nitrilium salts disproportionate to form the more stable chloroimonium salts.



Study of the fusion diagrams of binary mixtures of hydrogen chloride with acetonitrile, propionitrile, butyronitrile, and benzonitrile has shown that these nitriles can form low-melting non-saltlike molecular compounds of formula $\text{RCN}\cdot\text{HCl}$.³⁹ Thus $\text{CH}_3\text{CN}\cdot\text{HCl}$ melts at -63.2° and $\text{C}_2\text{H}_5\text{CN}\cdot\text{HCl}$ at -97.2° . The molecular compounds $\text{RCN}\cdot\text{HCl}$ may be formed as a result of reaction between the unshared electron pair of the nitrile nitrogen and a molecule of hydrogen chloride. The above nitriles also form complexes of composition $\text{RCN}\cdot n\text{HCl}$ (where $n = 5$ or 7) and $2\text{RCN}\cdot 3\text{HCl}$, which have even lower melting points than the compounds $\text{RCN}\cdot\text{HCl}$. It is suggested³⁹ that in these compounds the unshared electron pair of the nitrogen and also π electrons act as donors, and the compounds $\text{RCN}\cdot n\text{HCl}$ are apparently therefore examples of the so-called π -complexes.

It must be emphasised that the compounds described by Murray and Schneider³⁹ are fundamentally different from all the other hydrohalogenation products of nitriles, which are salts. All the described molecular compounds retain the covalent bond between the hydrogen and chlorine in the hydrogen chloride molecule, and are apparently intermediates in the formation of the saltlike nitrile hydrohalides (see below).

B. Some Properties of Solutions of Hydrogen Halides in Nitriles

It has been known for a long time that hydrogen halides ionise comparatively readily in nitriles⁴⁰, and the conductance has been the most studied physical property of the solutions^{25,37,41-44}. The polarographic behaviour of solutions of hydrogen chloride in acetonitrile has been investigated^{45,46}, and cryoscopic measurements have been made on dilute solutions of hydrogen chloride in benzonitrile³⁷.

The conductimetric study of the reaction between hydrogen chloride and acetonitrile⁴³, benzonitrile, β -chloropropionitrile, $\alpha\beta$ -dichloropropionitrile, β -ethoxypropionitrile, and adiponitrile⁴⁴ shows that the conductance decreases as the HCl concentration is increased. In some cases the conductance curves pass through a minimum. Plots of λ (equivalent conductance) against \sqrt{c} for solutions of HCl in adiponitrile and β -chloropropionitrile show two intersecting straight lines, *i.e.* within certain concentration limits the conductivity of these dilute solutions is described by Kohlrausch's formula for low concentrations of strong electrolytes. Plots of λ against $1/\sqrt{c}$, which are used

for systems with chemical reaction⁴³, are linear for acetonitrile, adiponitrile, benzonitrile, and β -chloropropionitrile, the last two giving two intersecting lines⁴⁴.

The complexity of the conductance-concentration relationship, and in particular the presence of intersecting curves on the diagrams, can be explained by a given nitrile forming with the hydrogen halide, depending on the reaction conditions, not one but several compounds, some which ionise and some which do not⁴⁴. The nitrile solutions may contain, for instance, the molecular compounds $\text{RCN}\cdot n\text{HCl}$. Conductimetric and cryoscopic study of dilute solutions of HCl in benzonitrile has shown³⁷ that the compound $\text{C}_6\text{H}_5\text{CN}\cdot\text{HCl}$ is formed. Of the known types of saltlike compounds, the nitrile solutions may contain halogenoimonium halides^{14,47}, compounds with the HX_2^- anion²⁵, and possibly also nitrilium salts and salts of composition $2\text{RCN}\cdot n\text{HX}$ (see below). It must be assumed⁴⁴ that in general the nature of the reaction products between nitriles and hydrogen halides changes continuously in the qualitative and quantitative sense as the relative proportions of the reactants are varied.


The formation of compounds between nitriles and hydrogen halides is sometimes quite slow; for example, at 0° in ether (twice the reactant volume) the reaction between adiponitrile and hydrogen chloride in the molar ratio 1:2 is practically complete only after 24 h.⁴⁷ The conductance of freshly-prepared solutions of hydrogen halides in nitriles increases with time and reaches a maximum^{42,44}. The slowness of the reaction enables the conductimetric method to be used to study the reaction mechanism^{36,42}.

The increase in conductance of solutions of hydrogen halides in nitriles with time means that the ionising compounds are not formed immediately. A similar effect was found with the systems acetonitrile-picric acid⁴⁸ and acetonitrile-iodine monohalides⁴⁹.

The donor-acceptor properties of the solvent and solute influence the rate of equilibration in the reaction of nitriles with hydrogen halides. It is assumed that the slow increase in conductance is due to the high ionisation potential of the nitrile, so that the latter is slowly protonated^{42,48}. The polarisability of solvent and solute must also influence the rate of ionisation to a similar extent; the maximum conductance is reached earlier in the acetonitrile-HI and acetonitrile-HBr systems than in the acetonitrile-HCl system⁴², the equilibrium conductances in these systems being in the ratio 60:20:1.³⁶


From the conductivity data, Janz and Danyluk suggested⁴² two mechanisms for the reaction between nitriles and hydrogen halides. The primary reaction products are unionised molecular compounds ("outer complexes") which are then converted into saltlike compounds ("inner complexes")^{49,50}. Mulliken and Reid associate the slow salt-formation with a considerable energy barrier for this reaction⁵¹.

The first of Janz' and Danyluk's mechanisms⁴² for the reaction of nitriles with hydrogen halides comprises the following stages:




In the compound $\text{RCN}\cdot\text{HX}$, which is an "outer complex", the hydrogen halide molecule still retains its σ bond, but this is then broken to form the "inner complex", which ionises in solution to form the solvated anion X_{solv}^- ; this then reacts with a second hydrogen halide molecule to form the anion $\text{HX}_{2\text{solv}}^-$.

The second mechanism⁴² involves a nucleophilic attack by hydrogen halide on the "inner complex" to form the halogenoimmonium halide, the anion of which is then solvated:



The solvated halide ion can obviously again react with a second hydrogen halide molecule to form the anion $\text{HX}_{2\text{solv}}^-$, so that the overall reaction in the solution is




The compound $[\text{C}_6\text{H}_5\text{CCl}=\text{NHC}_6\text{H}_5]^+\text{HCl}_2^-$ has recently been prepared⁵² by reacting *N*-phenylimidoyl chloride with hydrogen chloride.

The second mechanism was essentially that put forward¹⁴ in 1955 as the only possible one for the reaction between nitriles and hydrogen halides. Attention was directed to the exceptional energetic instability of nitrogen-unsubstituted nitrilium salts, and to their strong tendency to add halide ions to yield undissociated imidoyl halides which, being bases, then react further with the hydrogen halide¹⁴. The addition of hydrogen halides to *N*-substituted imidoyl halides has been described^{53,54}.


C. Nitrile Hydrohalides of Composition $2\text{RCN}\cdot n\text{HX}$

Certain halogenonitriles react with hydrogen halides under normal conditions to give saltlike products which have different chemical properties from the nitrile hydrohalides described above⁵⁵⁻⁵⁹. Although these salts were known to give secondary amides with water, the attempt to assign to them a more or less acceptable structure was for long unsuccessful, until Grundmann and coworkers convincingly demonstrated⁶⁰ that they are dimer salts of composition $2\text{RCN}\cdot\text{HX}$ and have the *N*-alkylideneamidine structure (E) or (F):



Alkylideneamidine chlorides of this type were prepared⁶⁰ from mono-, di-, and tri-chloroacetonitrile and α -dichloropropionitrile; the last reacts similarly with hydrogen bromide.


The *N*-alkylideneamidine structure has been assigned to the reaction product of hydrogen chloride with α -cyanoethyl acetate⁶¹. It has been reported that unsubstituted acetonitrile forms a dimer salt³²; the reaction product was characterised as α -chloroethylideneamidine. The treatment of chloro-, fluoro- and bromo-acetonitriles with hydrogen chloride or bromide at about room temperature gives compounds of composition $2\text{RCN}\cdot 3\text{HCl}$ and $2\text{ClCH}_2\text{CN}\cdot 3\text{HBr}$ ^{34,62} having the same chemical properties as the salts $2\text{RCN}\cdot\text{HX}$; the products may therefore be formulated as, for instance,




The reaction products of chloro- and bromo-acetonitrile with hydrogen chloride, which were formulated as imidoyl chloride hydrochlorides⁶³, are in fact dimer salts³⁸.

Lazaris and coworkers³⁴, using the reactions of fluoro- and bromo-acetonitrile with hydrogen chloride, were the

first to show that, depending on the reaction conditions, the same nitriles may produce either halogenoimmonium halides or dimer salts: the former were obtained between -50° and -5° , and the latter between 0° and 20° and above. In each case unstable outer and inner complexes are apparently first formed; at low temperatures these compounds are stabilised by addition of a hydrogen halide molecule to form the halogenoimmonium halide. At a higher temperature the equilibrium in the formation of the unstable complexes is displaced towards the formation of the initial products, producing a higher concentration of the free nitrile. The cyano-group has both nucleophilic and electrophilic properties, and the unstable complexes are therefore stabilised by addition of the nitrile, giving the dimer salts:




Cyclic electron transfer is also a probable mechanism in the formation of the dimer salts:



D. Preparation of 1,3,5-Triazines

In the presence of hydrogen halides some nitriles, chiefly α -halogenonitriles, polymerise at room temperature and above to produce 1,3,5-triazines^{1,64}. Grundmann and coworkers⁶⁰ consider that the trimerisation of nitriles is preceded by the formation of the dimer salts (see above), which then react with the nitrile by the diene synthesis to form 1,3,5-triazines. However, this mechanism is inconsistent with their inability to prepare 1,3,5-triazines from the dimer salts and nitriles, and chloroacetonitrile, which forms the dimer salt more easily than any other nitrile, does not trimerise at all; on the other hand, trichloroacetonitrile, which gives the dimer salt with great difficulty⁶⁰, polymerises exceptionally readily. Hence the dimerisation of nitriles to dimer salts cannot be an intermediate stage in the production of 1,3,5-triazines, and these must be considered as competing reactions, as the early workers assumed.

The comparative ease of formation of 1,3,5-triazines may be due to cyclic electron transfer:



According to this mechanism, both the free nitrile and its hydrogen halide complex take part in the reaction. The required free nitrile content is met because the triazine synthesis is usually carried out at room temperature and above, when the nitrile hydrochlorides are already partially dissociated. Since the reaction proceeds less easily in the absence of the nitrile hydrohalide, an increase of reaction temperature above a certain limit should reduce the yield of triazine, and this has been confirmed experimentally⁶⁵. The experimentally observed evolution of hydrogen chloride in the formation of 1,3,5-triazines⁶⁰ also does not contradict the suggested reaction mechanism.


Dimer salts and triazines are obtained most easily from α -halogenonitriles: electronegative α -substituents produce a comparatively low electron density at the nitrile carbon atom, and hence increase the electrophilic reactivity of the nitriles.

Up to the present, therefore, the reaction of nitriles with hydrogen halides has given three main types of stable compound: the dihydrohalides (apparently halogenoimmonium halides), dimer salts (*N*-alkylideneamidines), and 1,3,5-triazines. The reactions commence with the formation of unstable complexes between the reactants, and these are then stabilised by one of the above mechanisms, depending on the properties of the initial components and the reaction conditions.

E. Hydrogen Cyanide Hydrohalides

Hydrogen cyanide reacts with hydrogen chloride to produce two compounds, of composition 2:3 and 2:1.

The sesquichloride $2\text{HCN}\cdot 3\text{HCl}$ ⁶⁶ has been assigned the dichloromethylformamidine hydrochloride structure ^{67,68}:




or, in more recent work, the following non-saltlike structure ⁶⁹⁻⁷⁴:



The sesquichloride readily loses two-thirds of its hydrogen chloride to form the compound $2\text{HCN}\cdot\text{HCl}$, which was earlier considered to be chloromethyleneformamidine $\text{ClCH}=\text{NCH}=\text{NH}$. ⁶⁹ Further dehydrochlorination, e.g. by means of quinoline, gives the so-called "dimeric hydrocyanic acid", ⁷⁵ which has been given the iminoformyl-carbylamine structure $\text{C}=\text{NCH}=\text{NH}$. ⁷⁶

The structure of these reaction products was considered afresh by Grundmann and coworkers ⁷⁷⁻⁸⁰, who put forward completely new formulae for them. According to Grundmann, the product initially formed is 2,4,6-trichlorohydro-1,3,5-triazine hydrochloride, the compound thought to be chloromethyleneformamidine is 1,3,5-triazine sesquichloride, and exhaustive dehydrochlorination produces 1,3,5-triazine, which is unstable:




At various times the reactions of the hydrogen cyanide hydrochlorination products with water, alkali, alcohols ⁶⁶, amines ⁶⁸, aromatic compounds (in the presence of aluminium chloride) ^{67,73,74}, and other compounds have been studied. The compositions of the molecular compounds formed between aluminium chloride and all three of the $\text{HCN}\text{-HCl}$ reaction products have also been investigated ⁸⁰.

Hydrogen cyanide reacts with hydrogen bromide to give the analogous bromine compounds $2\text{HCN}\cdot 3\text{HBr}$ and $2\text{HCN}\cdot\text{HBr}$. ^{71,81}


III. REACTIONS OF NITRILES WITH HYDROGEN HALIDES AND NUCLEOPHILIC REAGENTS

As stated above, in the absence of acids nitriles do not as a rule react with hydroxylic or other nucleophilic reagents. The addition of a proton to the nitrile nitrogen atom

"opens up" the triple bond, and hence considerably increases the ability of the cyano-group to react with electron-donating compounds; it must be assumed, therefore, that the reaction of nitriles with water, alcohols, organic acids, mercaptans, and other nucleophilic reagents is preceded by the formation of nitrile-hydrogen halide addition products. Possible intermediates in these reactions of nitriles are the unstable strongly polarised 1:1 nitrile-hydrogen halide complexes described in the previous section; these are then stabilised by addition of the nucleophilic reagent:



Since nitriles also form extremely stable reaction products with hydrogen halides, the halogenoimmonium halides, these must also be considered to take part in the reactions between nitriles, hydrogen halides, and nucleophilic reagents:



We shall consider below the reactions occurring, in the presence of hydrogen halides, between nitriles and hydroxylic compounds and amines, and also the Gattermann and Hoesch syntheses. In discussing these reactions, we shall show that in many cases there has been experimental confirmation of two main reaction stages: reaction between the nitrile and the hydrogen halide, and, subsequently, addition of the electron-donating compound. In some cases the experimental results have been sufficient to decide which of the two mechanisms is predominant in the first stage, i.e. whether only the unstable complexes are formed, or whether the saltlike nitrile dihydrohalides are also produced. Some of the results do not exclude the possibility, in individual cases, of synchronous reactions, i.e. reactions passing through a single transition state. Knowledge of the mechanism of the electrophilic reactions of nitriles has enabled known syntheses to be improved and new ones developed.

This section includes a subsection devoted to the reactions of the nitrile dimer salts with hydroxylic compounds. Since the previously little known amide hydrohalides are often the final products in the reaction of nitriles with hydrogen halides and hydroxylic compounds, a separate subsection is devoted to their properties. Finally, we shall consider the selective reactivity of one of the cyano-groups of adiponitrile, which is particularly evident in the presence of hydrogen chloride.

A. Reactions with Water


1. Preparation of amides and their hydrohalides. Amides are obtained when nitriles are treated with concentrated hydrochloric acid at room temperature ⁸²⁻⁹⁴, or, more rapidly, at 50°-60°. At higher temperatures carboxylic acids are formed ^{90,95}. Nitriles containing reactive groups are hydrated at room temperature or at 0°.

The hydrolysis of nitriles with hydrochloric acid of various concentrations has been investigated ⁹⁶; with concentrated acids hydrolysis stops at the amide stage, while in dilute acids carboxylic acids are obtained. Since the HCl content of concentrated hydrochloric acid at room temperature is at most 36-38%, the amides formed by the hydration of nitriles are sometimes partially or completely

hydrolysed to carboxylic acids even at low temperatures⁹⁷. To suppress hydrolysis of the amides the nitriles must be hydrated under conditions such that there is no large excess of water. In some cases the nitriles have been dissolved in concentrated hydrochloric acid, or mixed with a little water, and the solution cooled to between -5° and 0° and saturated with hydrogen chloride^{17,98-101}. Treatment of the reaction products of nitriles and dry hydrogen chloride with water usually gives the initial nitriles^{13,17,32}, but in certain cases nitrile-hydrogen chloride reaction products have been said to be intermediates in the production of amides from nitriles^{92,102,103}. The formation of amides from nitriles and water in the presence of hydrogen chloride is the basis of proposed methods for the dehydration of acids¹⁰⁴ and esters¹⁰⁵.

As early as 1877 Pinner and Klein⁸ passed hydrogen chloride into a mixture of equivalent quantities of water and acetonitrile or benzonitrile and obtained compounds which they described as amide hydrochlorides, but the reaction conditions and the yield and properties of the salts were not given. Later it was stated^{25,99,103} that amide hydrochlorides are intermediates in the preparation of amides from nitriles, water, and hydrogen chloride, but here also the amide hydrochlorides were not identified. Treatment of adiponitrile with water and hydrogen chloride¹⁰⁶ gave a chlorine-containing product which was wrongly described as an imidoyl chloride hydrate.

According to Zil'berman and Kulikova's results^{47,107-112}, earlier statements^{13,32} that the action of water on nitrile-hydrogen halide reaction products can only produce the initial reactants are only partly correct. With an excess of ice water acetochloroimonium chloride does in fact dissociate to give the initial nitrile, but the addition of small quantities of water to an ether solution of the chloride gives a precipitate of acetamide hydrochloride (acetimidic acid hydrochloride), which on hydrolysis produces the free amide:



Treatment of mixtures of equivalent quantities of nitriles and water with hydrogen chloride has given high yields of amide hydrochlorides derived from aliphatic acid amides^{47,110}, aliphatic acid diamides¹¹¹, and ring-substituted benzamides³⁵. In some cases the amide hydrochlorides are very unstable, readily losing hydrogen chloride, and even before hydrolysis are partially converted into the free amides. Some nitrile hydration products (*e.g.* from chloroacetonitrile¹¹⁰) contained so little hydrogen chloride that they should be classed as free amides rather than amide hydrochlorides. Hydration of trichloroacetonitrile¹¹⁰ and $\alpha\beta$ -dichloropropionitrile¹¹² gives the free amides. The considerable negative inductive effect of the chlorine atoms in these amides apparently so weakens the nucleophilic properties of the oxygen atom that it becomes unable to add a proton. Unlike these, however, capronitrile¹⁰⁹, methacrylonitrile, and lactonitrile¹¹² do apparently form amide hydrochlorides, but they could not be isolated from the reaction mixture. The corresponding amides were obtained immediately on hydrolysis of the reaction mixture.

As a rule, the order of addition of the nitrile, hydrogen chloride, and water to the reaction mixture plays substantially no part in the reaction, but in certain specific cases a particular order of addition is essential for obtaining optimum yields of hydrochlorides or free amides. In the


preparation of most amides equivalent quantities of nitrile and water are added to an organic solvent, and hydrogen chloride is passed into the solution; after a definite time the precipitate is filtered off. Because of the reactivity of the hydroxy-groups in lactonitrile and acetonecyano-hydrin, the corresponding amides can only be prepared by adding the nitrile to a solution of hydrogen chloride in aqueous ether¹¹². Adiponitrile was converted quantitatively into δ -cyanovaleramide hydrochloride by passing hydrogen chloride into a solution of adiponitrile in ether for at least four hours and then adding the calculated quantity of water⁴⁷. This procedure was also the most suitable for preparing diamides of dibasic acids¹¹¹.

The larger the excess of hydrogen chloride with respect to the nitrile, the more rapidly is the amide hydrochloride formed¹¹⁰; evidently excess of hydrogen chloride increases the concentration of primary nitrile-hydrogen chloride reaction products, which then react with the water. An excess of hydrogen chloride is not necessary in the hydration of nitriles of high electrophilic reactivity. In the preparation of lactamide, α -hydroxyisobutyramide hydrochloride¹¹², and δ -cyanovaleramide hydrochloride the molar ratio of nitrile to hydrogen chloride should be 1:2. The formation of most amide hydrochlorides is complete within 24 h, but in some cases (*e.g.* with stearamide, adipamic acid methyl ester¹¹⁰, *p*-toluamide³⁵, etc.) the reaction had to be continued for 5-10 days.

Amide hydrochlorides are usually prepared at $\sim 0^{\circ}$,¹¹⁰ but with some nitriles of low reactivity (*e.g.* stearonitrile, capronitrile, and benzonitrile) room temperature was preferred.


Amide hydrochlorides can be prepared from nitriles, water, and hydrogen chloride in the presence or absence of inert organic solvents, so long as the nitrile dissolves in the reaction mixture. The amide salts are often prepared in ether¹¹⁰. Substituted benzamides are prepared in dioxane, which unlike ether partially dissolves substituted benzonitriles³⁵. The comparatively slow-reacting capronitrile and stearonitrile have been hydrated in toluene, since ether, dioxane, and similar oxygen-containing compounds were not found to be suitable solvents. The reaction between hydrogen chloride and the ethers apparently displaces the equilibrium of the nitrile-hydrogen chloride reactions towards the initial reactants, and the reduced concentration of intermediate products in the reaction mixture lowers the rate of formation of the amides.

When $\alpha\beta$ -unsaturated nitriles (acrylonitrile, methacrylonitrile) are treated with hydrogen chloride and water, they are hydrated and also hydrochlorinated at the double bond¹¹²:



Some nitriles (β -ethoxypropionitrile, azodiisobutyronitrile¹¹⁰, *o*-nitrobenzonitrile³⁵, etc.) have proved impossible to hydrate to amides, presumably because of steric hindrance. *Ortho*-substituted benzonitriles are known^{113,114} to be inactive in other reactions also: *e.g.* they do not form imidate hydrochlorides with hydrogen chloride and alcohols. The stability of β -ethoxypropionitrile under normal conditions for the preparation of amide hydrochlorides may be because, as has been suggested¹¹⁵, β -substituents in the aliphatic series produce the same steric hindrance as is found in *ortho*-substituted aromatic compounds.


The reaction of nitriles with deuterium chloride and deuterium oxide is a convenient method for preparing

ON-deuterated imidic acid hydrochlorides¹¹⁶:

Treatment of these salts with heavy water causes deuterium exchange at the α -carbon atom, forming *NC*-deuterated amides with a high deuterium content. Trichloroacetone nitrile, which has no α -hydrogen and does not form the imidic acid hydrochloride, reacts with deuterium chloride and deuterium oxide to produce deuterotrichloroacetamide in high yield. Obviously, other isotopic varieties of water will react with nitriles in the same way as D_2O .

2. Mechanism of the reactions of nitriles with hydrogen halides and water. Two reaction mechanisms for the reaction between nitriles, hydrogen halides, and nucleophilic reagents were given at the commencement of Section III: (a) the formation of unstable complexes between nitriles and hydrogen halides, and (b) the formation as intermediates of saltlike nitrile hydrohalides (halogenoimmonium halides).

Investigation of the reaction of substituted benzonitriles³⁵ and aliphatic dinitriles¹¹¹ with hydrogen chloride and water has confirmed that the reaction proceeds in stages. The possibility of carrying out the reaction by way of chloroimmonium chloride was shown experimentally in the selective hydration of one nitrile group of adiponitrile⁴⁷. To obtain the theoretical yield of δ -cyanovalerimidic acid hydrochloride, the 1:2 nitrile-hydrogen chloride reaction product (δ -cyanovalerohydrochloroimmonium chloride) must first be prepared:



There are cases, *e.g.* in the hydration of chloroacetone nitrile⁶², when the chloroimmonium chloride formation is unnecessary or even impossible. In the treatment of certain reactive nitriles, particularly halogenoacetone nitriles, with water and hydrogen chloride, the unstable outer and inner complexes $\text{RCN} \cdot \text{HCl}$ may be intermediate products. It is also possible that in this case the addition of water and hydrogen chloride takes place in a single transition state.

It must be assumed that in most actual cases of the hydration of nitriles in the presence of hydrogen halides the reaction mechanism is more complex than those given above. The addition of water may proceed simultaneously, for example, by two or three mechanisms.


In each stepwise reaction mechanism there are at least two stages: (1) reaction of the nitrile with hydrogen chloride, and (2) reaction with the nucleophilic hydroxy-compound. To decide which is the rate-determining step, the influence of substituents in benzonitrile on the rate of formation of the corresponding benzimidic acid hydrochlorides has been studied³⁵. The activating effect of the substituents on the rate of reaction decreases in the series:

- (1) *p*-substituents: $\text{NO}_2 > \text{COOH} > \text{Br}, \text{Cl} > \text{CH}_3 > \text{H} \gg \gg \text{OCH}_3$;
- (2) *p*- $\text{NO}_2 > m$ - $\text{NO}_2 \gg o$ - NO_2 .

The more electronegative the *p*-substituent, the more readily, as a rule, is the nitrile hydrated in the presence of the hydrogen chloride. Since the inductive effect and the effect of conjugation of electronegative substituents in


benzonitrile reduce the electron density at the nitrile carbon atom, the rate-determining step in the reaction of nitriles with hydrogen chloride and hydroxy-compounds is the nucleophilic attack by water. In one case, however, this rule is not obeyed: *p*-tolunitrile has an electropositive substituent but stands above benzonitrile in the above series. The comparative ease with which *p*-tolunitrile is hydrated is apparently due to the activating effect of the *p*-methyl substituent on the first reaction step (reaction of the nitrile with hydrogen chloride). The presence of the electropositive substituent in the tolunitrile molecule causes additional displacement of electrons towards the nitrogen (the dipole moment of *p*-tolunitrile is 4.37 D, against 4.08 D for benzonitrile¹¹⁷) and so facilitates its nucleophilic reactions. Thus in the case of *p*-tolunitrile the rate of formation of the imidic acid hydrochloride is determined both by the reaction with the nucleophilic reagent and by the reaction of the nitrile with the hydrogen chloride. The reaction of *p*-tolunitrile with hydrogen chloride in the absence of water³⁵ supports this conclusion: it readily forms the chloroimmonium chloride, while under the same conditions benzonitrile does not. *p*-Methoxybenzonitrile forms the chloroimmonium chloride even more readily than *p*-tolunitrile, even in the presence of water, but because of the electropositive methoxy-group, the nucleophilic attack is completely suppressed, and *p*-methoxybenzimidic acid hydrochloride is not produced.

3. Reaction of alkyl thiocyanates with hydrogen chloride and water. Organic thiocyanates often react in the same way as nitriles. Treatment of methyl, ethyl, *n*-propyl, and *n*-butyl thiocyanates with hydrogen chloride and water produces the corresponding *S*-alkyl esters of monothiocarbamic acid in moderate yields¹¹⁸:




The comparatively slow hydration of alkyl thiocyanates is due to the presence in the SCN group of the sulphur atom, which has unshared electron pairs and by conjugation increases the electron density of the thiocyanate carbon atom, so hindering nucleophilic attack on this atom by a water molecule.


Heating methyl thiocyanacetate with hydrochloric acid produces 2,4-dioxothiazolidine¹¹⁹:

**B. Reactions with Alcohols**

1. Preparation of imidate hydrohalides. Boiling a mixture of a nitrile, alcohol, and water with strong acids produces esters¹²⁰, and these are also obtained by passing hydrogen chloride into hot mixtures of nitriles, alcohols, and water¹²¹. In an anhydrous medium the reaction between nitriles, alcohol, and hydrogen chloride (the so-called Pinner synthesis¹²²) produces imidate hydrochlorides (imino-ether hydrochlorides):^{8,123}




Under the same conditions nitriles, mercaptans, and hydrogen halides produce thioimide hydrohalides:




Imide hydrochlorides, unlike amide hydrochlorides, are well known in organic chemistry. The preparation and chemical properties of imidates have been described in recent detailed reviews^{17,122} covering the literature up to 1959, and we shall therefore consider here only matter outside the scope of these reviews and data which have appeared only after the date of submission of Roger and Neilson's review¹²².

Although three reactants participate in the formation of imide hydrohalides, it is usually presented as a single-stage reaction^{17,122}, but the mechanism of the reaction of nitriles and hydrogen halides with alcohols is close to the mechanism of their reaction with water. The imide hydrohalides are formed by way of the corresponding halogenoimmonium halides¹²⁴ and also directly from unstable reaction products of nitriles and hydrogen halides with alcohols. The halogenoimmonium halide mechanism is also confirmed by the treatment of acetochloroimmonium chloride³², bromoacetochloroimmonium chloride, and bromoacetobromoimmonium bromide³⁴ with alcohols, when the corresponding imide salts are obtained.


Imide hydrochlorides are formed not only from nitriles, but also from other compounds containing the cyano-group, when treated with alcohols and hydrogen chloride. Cyanamide, methanol, and hydrogen chloride give methylisourea hydrochloride¹²⁵:



Organic thiocyanates react similarly¹²⁶:



Various quantities of hydrogen halide have been used in the preparation of imidates. In his first study of imidates Pinner⁸ gave the following structure for the reaction products of nitriles, alcohols, and hydrogen chloride:



When dried, these split out one molecule of hydrogen chloride, leaving the imide hydrochloride. On this basis, in his early work^{8,127} on the preparation of imide salts Pinner used the nitrile, alcohol, and hydrogen chloride in the ratio 1:1:2, but later he apparently realised that the imide group usually combines with only one hydrogen chloride molecule, and in his monograph¹²³ he gave the optimum reactant ratio as 1:1:1. Up to the present this ratio of 1:1:1 has very frequently been used in the production of imide salts, or a slight excess (10–25%) of hydrogen chloride is used^{128–130}. However, as in the preparation of amide hydrochlorides (see above), an excess of hydrogen chloride accelerates the reaction, and therefore, when free hydrogen chloride does not cause decomposition of the resulting imide hydrochloride, it is useful to saturate the reaction mixture with hydrogen chloride^{131–136}. The use of a large excess of hydrogen halide is particularly effective when nitriles of low electrophilic reactivity are reacted with alcohols. There are particular cases¹²⁴ when the reaction product can only be obtained in a pure state by using a reactant ratio of 1:1:2; as will be shown

below, this ratio is not due to addition of two hydrogen chloride molecules to the imide, but is necessary because the intermediate reaction product is sometimes the corresponding chloroimmonium chloride.

The compound 6-methoxy-2-naphthimino-ether dihydrochloride was prepared in 1947.¹³⁷ The formation of this imide with two molecules of hydrogen chloride can be explained either by one of the HCl molecules adding to the oxygen of the methoxy-group, or by the imide salt having the HCl_2^- anion.


It has recently been suggested that hydrogen bromide or iodide be used to accelerate imide salt formation and increase the yield of product¹³⁸.

Ethyl ethoxycarbonylacetylacetimidate hydrochloride has been prepared from ethyl cyanoacetate and ethanol in the presence of hydrogen chloride formed *in situ* from equimolar quantities of thionyl chloride and water¹³⁹.

In addition to the solvents previously used¹²², CCl_4 has also been employed as reaction medium for the preparation of imide hydrohalides¹⁴⁰. When imide salts are to be prepared from nitriles which are insoluble in the usual organic solvents, the reaction can be carried out in suspension^{133,141}. In the preparation of the imide salt from β -cyanoethyl *p*-toluenesulphonate a saturated solution of hydrogen chloride in absolute alcohol was shaken at room temperature with the solid nitrile until a homogeneous solution was obtained¹⁴².

In the preparation of the overwhelming majority of imide salts, the order of addition of the nitrile, alcohol, and hydrogen halide to the reaction mixture is immaterial; usually the nitrile and alcohol are mixed with an inert organic solvent, into which the hydrogen halide is passed at $\sim 0^\circ$. In individual cases other procedures have proved convenient. To prevent side reactions in the preparation of the imidic acid hydrochloride derived from ethyl β -chlorolactate acid, hydrogen chloride is passed into the alcohol and then the nitrile is added while continuing the addition of hydrogen chloride¹⁴³.

Imide hydrochlorides are used in the preparation of certain polymerisable acrylates¹⁴⁴. When hydrogen chloride is passed through a mixture of alcohol and an $\alpha\beta$ -unsaturated nitrile, β -chloroimide hydrochlorides are obtained^{142,145}, for example¹⁴⁴:




Hydrolysis of the imide hydrochlorides produces esters of β -chlorocarboxylic acids, which in the presence of dehydrohalogenation agents are converted into esters of $\alpha\beta$ -unsaturated acids.

Imide hydrochlorides derived from dibasic aliphatic acids can be used as additives, *e.g.* for food products and synthetic polymers, to prevent the catalytic action of metals on the organic materials; they inhibit, for example, their autoxidation¹⁴⁶.


2. New syntheses from dinitriles. A large number of diimide salts obtained from dinitriles and mono-ethyl alcohols or mononitriles and glycols have been described¹²². Woodburn and coworkers¹⁴⁷ have studied the reaction of cyanogen with hydrogen chloride and certain

glycols; diimide hydrochlorides were prepared from propane-1, 2-diol, propane-1, 3-diol, and butane-2, 3-diol:




In most other reported reactions between dinitriles, glycols, and hydrogen chloride, however, mixtures of unidentified imide salts were obtained¹⁴⁸.


In 1959 Zil'berman and Kulikova showed^{124,149} that for successful preparation of individual compounds from certain dinitriles, glycols, and hydrogen chloride, the hydrochlorination of the dinitrile and the reaction of the product with the alcohol must be carried out separately, *i.e.* the dinitrile should first be treated with the calculated amount of hydrogen chloride, and then the alcohol added after a definite time. This procedure gave a number of interesting compounds from adiponitrile, which, because of the selective reactivity of one of its cyano-groups, gave the following series of reactions^{124,149-152}:



The yields of alkyl δ -cyanovalerates¹²⁴, ω -hydroxyalkyl δ -cyanovalerates, and alkylene (from alkanediols) di(δ -cyanovalerates)¹⁵² were close to theoretical. Earlier procedures for the preparation of alkyl δ -cyanovalerimides^{129,153,154} gave the desired products in low yields and in admixture with a molecular compound formed between the alkyl imide hydrochloride and adiponitrile¹²⁴. It is important to obtain alkyl δ -cyanovalerates in high yields because they are starting materials for the production of caprolactam¹⁵⁵:




The reaction between aliphatic dinitriles and hydrogen chloride in a molar ratio of at least 1 : 4 in the cold produces solid products, apparently di(chloroimonium) dichlorides, which precipitate if insufficient solvent is used. These were treated with twice the molar quantity of glycols without separation from the reaction mixture. The resulting diimide hydrochlorides when treated with an excess of water gave di(ω -hydroxyalkyl) esters of dibasic acids:




This reaction has been carried out with such glycols as butane-1, 4-diol, pentane-1, 5-diol, and hexane-1, 6-diol.^{151,152,156} Di(ω -hydroxyalkyl) esters were not obtained from ethylene-glycol or *p*-xylyleneglycol, since under the reaction conditions the hydroxy-groups of these glycols are partly replaced by chlorine^{151,157}.

The reaction between dinitriles, hydrogen chloride, and glycols in the ratio 1 : n : 1, where $n \geq 4$, has been used to prepare polyimide hydrochlorides, which on hydrolysis give polyesters^{151,157-160}:



Polyesters of molecular weight 650-1650 have been obtained from the dinitriles of adipic, azelaic, sebacic, and *p*-phenylenediacetic acids and ethyleneglycol, butane-1, 4-diol, diethyleneglycol, and *p*-xylyleneglycol. In addition to nitrile and hydroxy-groups, chlorine was also found as an end-group in some cases. Increasing the reaction temperature from 0 to 20° considerably increased the chlorine content of the polyester and lowered its molecular weight. Excess of glycol or dinitrile also reduces the molecular weight of the final polyester. Unlike the pasty polycondensation products obtained from dibasic acids and glycols, which are slightly coloured, the polyesters obtained from polyimide hydrochlorides are often white powders, because of the mild conditions under which they are prepared.

3. Preparation of amides and their hydrochlorides. The reaction products obtained from nitriles, hydrogen chloride, and tertiary alcohols remained unidentified for a long time^{161,162}. Zil'berman and Sladkov showed¹⁶³ that many nitriles react in the cold with hydrogen chloride and tertiary alcohols to yield not imide ester salts, but imidic acid hydrochlorides (amide hydrochlorides) and tertiary alkyl chlorides:




Nitriles with strongly electronegative substituents in the α -position give the free amides and not the amide hydrochlorides.

Several mechanisms are possible for the formation of amides and their hydrochlorides in this reaction¹⁶³; for example, tertiary alcohols react with hydrogen chloride to give tertiary alkyl chlorides and water, and the latter may be expected to react with the nitrile-HCl reaction products to give imidic acid hydrochlorides.

According to a second mechanism, the initial nitrile-hydrogen chloride reaction products, particularly the chloroimonium chlorides, which have a high electrophilic reactivity, dehydrate the tertiary alcohols, the liberated water reacting with the nitrile and HCl in the normal way.

In a third possible mechanism, the nitriles and hydrogen chloride react initially with the tertiary alcohols in the same way as with primary and secondary alcohols, *i.e.* to form imide hydrochlorides, which then undergo the Pinner rearrangement^{8,122,164}:



Normally this last reaction is carried out by heating the imide salts in the absence of free hydrogen chloride, and the products are then the free amides and not the amide hydrochlorides. The basic literature on this reaction has


been reviewed¹²², and we only mention here that this splitting of imidate hydrochlorides has been shown to involve a bimolecular nucleophilic substitution reaction (S_N2)^{165,166}, although in some cases the reaction follows an S_N1 mechanism^{167,168}. With increasing electron-donating properties of the alkyl radical, a carbon atom of which undergoes substitution in such reactions, the rate of reaction first decreases (S_N2 mechanism) and then increases (S_N1 mechanism)¹⁶⁹, and it may therefore be assumed that when the alkyl radical R' is the strongly electropositive tertiary alkyl group, the splitting will proceed by an S_N1 mechanism. This also explains the ease with which tertiary alkyl imidate hydrochlorides are split, even at the moment of their formation.

Experiments on the hydration of a number of substituted benzonitriles using hydrogen chloride and dimethylethyl-methanol aimed at establishing whether the reaction of the nitrile with the hydrogen chloride or the hydration is the rate-determining step gave the same results as in the study of the mechanism of the reaction between nitriles, HCl, and water³⁵ (see p. 621).

Amides or their hydrochlorides are also sometimes obtained in the reaction of nitriles and hydrogen chloride with primary or secondary alcohols; this was observed, for example, in attempts to prepare primary alkyl imidate hydrochlorides from trichloro-, dichloro-, tribromo-, nitro- and dichloronitro-acetonitrile^{162,170-172}. Chloro-acetonitrile, which has only one electronegative chlorine atom in the α -position, gives the imidate hydrochloride on reaction with hydrogen chloride and methanol¹⁷¹; the use of a secondary alcohol (cyclohexanol) in this reaction gives the imidate or amide hydrochloride, depending on the reaction conditions¹⁶³. α -Phenylaminoacetonitrile gives the amide instead of the imidate¹⁷³.


Unlike tertiary alkanols, tertiary alkyl mercaptans react normally with nitriles and HCl, to form the corresponding thioimidate hydrochlorides^{161,174}.

The scission of imidate salts into amides and alkyl halides may be used preparatively¹²²; it has been used for the production of oligomeric amides from adipic acid and hexamethylenediamine¹⁷⁵:



C. Reactions with Organic Acids

Treatment of mixtures of nitriles and carboxylic acids with hydrogen halides gives amide hydrochlorides and the acid anhydrides or chlorides¹⁷⁶⁻¹⁸⁰:



A similar reaction is obtained with hydrocyanic acid, hydrogen chloride, and carboxylic acids¹⁸¹.


An excess of hydrogen halide accelerates the formation of the amide salt¹⁷⁹. The reaction proceeds in any solvent, polar or non-polar. Under otherwise the same conditions, the basicity of the solvent greatly affects the yield of imidic acid hydrochloride. It must be supposed that the reaction of the HCl with solvents having nucleophilic properties competes with the main reaction and noticeably reduces its rate¹⁸⁰. However, as in reactions of nitriles and hydrogen chloride with other nucleophilic reagents, the determining factor in the choice of solvent is not its acid-base properties, but the solubility of the reaction components in it.

An excess of the hydrating acid may be used as solvent, and this is particularly useful when the nitrile is poorly soluble in other organic solvents, e.g. with *p*-nitrobenzonitrile and *p*-cyanobenzoic acid¹⁸⁰.

Hydration of the cyano-group under the influence of hydrogen bromide and acetic acid or acetic anhydride is evidently the basis of the production of caprolactam from ϵ -alkoxycapronitriles by way of ϵ -bromocapramide¹⁸².

Study of the liquid reaction products of acetonitrile, hydrogen chloride, and butyric acid has shown that the acid is converted partly into the anhydride and partly into the acid chloride.


These data and the study of the reaction of a number of substituted benzonitriles with hydrogen chloride and acetic acid³⁵ show the general relationship between the hydration of nitriles using carboxylic acids and the other reactions described above between nitriles, nucleophilic reagents, and hydrogen halides; we may assume that the initial stages in all these reactions are the same, the reaction products of nitriles and hydrogen halides with acids being converted into the unstable isoimide salts¹⁷⁹. Since these should possess considerable acylating activity, they react with the acids to form imidic acid hydrochlorides and the acid anhydrides or halides:




N-substituted imidoyl chlorides react with carboxylic acids by an analogous mechanism¹⁸³.

The reactive imidoyl derivatives (isoimides) of trichloroacetic acid have been widely used in recent years as reagents for the selective esterification of phosphoric acid derivatives¹⁸⁴⁻¹⁸⁸. The isoimides are also intermediates in other acylation reactions^{189,190}.

Nitriles react with thio-acids in the presence of hydrogen chloride by the mechanism¹⁹¹ given above for carboxylic acids:




In the presence of hydrogen chloride nitriles react with thio-amides in ether in the cold¹⁹² or in dimethylformamide on heating¹⁹³ in the same way as with acids:




This reaction has been proposed for the preparation of thio-amides¹⁹³.

When aniline^{228,229} or other aromatic amine²³⁰ was heated with acetimidic acid hydrochloride, δ -cyanovaleric acid hydrochloride, and benzimidic acid hydrochloride at 160°–170°, the *N*-acylated amines were sometimes obtained in high yield. Aliphatic amines are acylated with considerably more difficulty than aromatic amines. In the benzoylation of hexamethylenediamine with benzimidic acid hydrochloride, the diamine hydrochloride had to be used in place of the free amine²³⁰.

Heating imidic acid hydrochlorides with *o*-phenylenediamine at 140°–190° produces benzoimidazoles in high yields:




These imidic acid hydrochlorides react at 100°–170° with alcohols (*n*-butanol, ethylene glycol) and phenols (phenol, *o*- and *m*-cresol, resorcinol monomethyl ether, *p*-nitrophenol) to form the corresponding esters²³¹:



The aliphatic alcohol esters were obtained in 61–75% yields, and the phenol esters in somewhat lower yields. In some cases, *e.g.* in the preparation of ethylene and *p*-phenylene di(δ -cyanovalerates) (from ethylene glycol and hydroquinone), acylation with the imidic acid hydrochloride is more convenient than earlier methods for the preparation of esters.


Two products were obtained by heating *o*-aminophenol with imidic acid hydrochlorides²³⁰, the *o*-hydroxyanilides or the 2-substituted benzoxazoles; the former were the main product if the reaction were carried out under reflux, and the latter if the water of reaction were removed as it formed. Heating the *o*-hydroxyanilides with catalytic amounts of acids converted them into the benzoxazoles; the former are apparently intermediates in the preparation of benzoxazoles from *o*-aminophenol and imidic acid hydrochlorides:



where R = CH₃, (CH₂)₄CN or C₆H₅. In the single-stage production of benzoxazoles (without intermediate separation of the *o*-hydroxyanilides from the reaction mixture) the unreacted imidic acid hydrochloride or the ammonium chloride may act as the acid catalyst. The presence of the acid catalyst should promote lactimisation of the amide group of the *o*-hydroxyanilide; this increases the reactivity of the amide group and the intramolecular interaction of the functional groups thus becomes possible.

Thermal decomposition of imidic acid hydrochlorides gives the acid, acid halide, nitrile, and a secondary amide^{199,200}. The mechanism of the decomposition was investigated by studying the pyrolysis products (180°) from δ -cyanovaleric imidic acid hydrochloride²³²; hydrogen

chloride, ammonium chloride, di(δ -cyanovaleramide), δ -cyanovaleric acid, adiponitrile, adipamide, δ -cyanovaleramide and, apparently, δ -cyanovaleric anhydride were found. The following mechanism based on these results was suggested:



The pyrolysis of other imidic acid hydrochlorides shows the same features. This reaction may be used for the preparation of secondary amides²³². The reaction between formamide hydrochloride and *m*-nitrobenzaldehyde gives *m*-nitrobenzylidenebisformamide⁷².

The thermal decomposition of amide hydrochlorides thus gives products which are formed from the free amides only at considerably higher temperatures^{224,233,234}. This difference in the behaviour of amides and their hydrochlorides is apparently due to the latter existing in the more reactive imidic acid form.

F. Selective Reactivity of One Cyano-Group in Adiponitrile

Several instances of the selective reactivity of one of the two cyano-groups of adiponitrile with hydrogen chloride and nucleophilic reagents (water, primary alcohols, and diprimary glycols) have been described above. Aliphatic dinitriles with a hydrocarbon chain shorter or longer than adiponitrile do not exhibit this selective reactivity of one cyano-group¹¹¹. The different reactivities of the cyano-groups in adiponitrile and such compounds as, for example, δ -cyanovaleric imidic acid hydrochloride and alkyl δ -cyanovalerimidate hydrochlorides are apparently due to the particular molecular configuration of adipic acid derivatives with the formation of cyclic systems^{111,235}.


One cyano-group of $\alpha\alpha\alpha'\alpha'$ -tetrachloroadiponitrile could not be selectively hydrated in the presence of hydrogen chloride; the mutual repulsion between the chlorine atoms and the nitrile nitrogen apparently hinders the formation of cyclic systems during the hydrochlorination and hydration. However, treatment of β -methyladiponitrile with HCl and water in the ratio 1:2:1 with subsequent hydrolysis gave a mixture of nitrile-amides in high yield. The introduction of the β -methyl substituent into adiponitrile apparently restricts its internal rotation, so that "convoluted" configurations are more easily formed and the reacting groups approach each other more closely²³⁵.

With adiponitrile itself, the selective reactions of one cyano-group with hydrogen chloride take place under rigorously defined conditions. The interaction between the solvent, dinitrile, and hydrogen chloride has a large influence on the reactivity of adiponitrile. In the reaction


between adiponitrile, hydrogen chloride, and water (in the ratio 1:2:1) at 0° selective reactivity is found in isopropyl, ethyl or butyl ether as solvent, but not in solvents which are more basic (tetrahydrofuran, tetrahydropyran or dioxane) or less basic (benzyl ether, β -chloroethyl ether, carbon tetrachloride, toluene, benzene, dichloroethane, or chloroform) than the above ethers²³⁵. At room temperature there is no selective reaction of one cyano-group of adiponitrile with hydrogen chloride and water, even in ether²³⁵.

There is no selective reaction of one cyano-group of adiponitrile with hydrogen chloride and such nucleophilic reagents as acetic acid¹⁷⁹ and alkanethiols²³⁵, apparently because of the high reactivity of the hydrogen atom of mercapto- and carboxy-groups, which readily interact with the cyano-group and so destroy the unstable intramolecular ring of the reacting molecule; this straightens out, and both adiponitrile cyano-groups become energetically equivalent.

A non-equivalent reactivity of the two cyano-groups of adiponitrile is also sometimes found in the absence of hydrogen chloride. The formation of the unsymmetrical amidoxime has been described²³⁶:




Hydrogenation of adiponitrile can produce ϵ -aminocapronitrile in high yield²³⁷⁻²⁴⁰:



G. Reactions with Amines

The most general method for the preparation of amidines from nitriles is based on the reaction between imidate hydrohalides and amines²⁴¹. Many attempts have been made to obtain amidines directly from nitriles without the intermediate stage of imidate salt formation, but the direct addition of amines only proceeds readily with nitriles having two or three α -halogen atoms, e.g. trichloroacetonitrile and trifluoroacetonitrile²⁴²⁻²⁴⁸:




The amination of α -halogeno-nitriles is accelerated in the presence of hydrogen halides (in practice, amine hydrohalides)²⁴². Primary and secondary aliphatic amines add to hydrogen cyanide to form unstable formamidines, which with water are converted into *N*-substituted formamides^{249,250}.

When nitriles free of electronegative substituents react with ammonia or amines in the presence of the base hydrohalide, amidines are generally not formed or are formed in only moderate yields. The rather sparse literature on these reactions has been given in a review on the chemistry of amidines²⁴¹.

Amidine hydrohalides are usually prepared from nitriles and amine hydrohalides at 170°–200° or above^{241,251,252}. The thermally unstable nitrile-hydrogen halide compounds cannot participate in the reaction under these conditions, and compounds of the halogenoimmonium halide type easily lose hydrogen halide in the presence of amines. The activating effect of hydrogen halides in the preparation of amidines from nitriles may be explained by a mechanism similar to those already given (see p. 619).


One case has been described in which a chloroimmonium chloride reacts with an amine in the cold²⁵²: at -16° acetonitrile dihydrochloride and dimethylaniline yield α -tri(dimethylaminophenyl)ethane.

The amidine group is formed by intramolecular condensation when various γ -, δ -, and ϵ -amino-nitriles are heated with hydrogen chloride or hydrochloric acid²⁵³⁻²⁵⁷; for example, at 275° $\alpha\alpha$ -diphenyl- γ -diethylaminobutyronitrile gives 1-ethyl-2-imino-3,3-diphenylpyrrolidine hydrochloride:



Certain 2-iminopyridine and 2-iminohexamethyleneimine hydrochlorides have been prepared analogously. Hydrolysis of 2-iminopolymethyleneimines sometimes yields the corresponding lactams.


Since the 2-iminohexamethyleneimines are obtained at high temperatures, it is difficult to see how the unstable hydrochlorides of the initial nitriles can be intermediates²⁵⁷. A more probable interpretation of the intramolecular reaction between the functional groups is cyclic electron transfer:



H. The Gattermann Reaction

The Gattermann and Hoesch (Houben-Hoesch) reactions differ from the reactions between nitriles, hydrogen halides, and nucleophilic reagents described above, in that new carbon-carbon bonds are formed.

In the Gattermann reaction²⁵⁸ an aromatic compound reacts in an inert solvent in the cold with hydrogen chloride and hydrocyanic acid to yield an aldimine hydrochloride, which on hydrolysis is converted to the aldehyde:




According to Gattermann, hydrogen cyanide and hydrogen chloride react under the above conditions to produce formimidoyl chloride $\text{HC}(\text{:NH})\text{Cl}$, but as in the case of hydrochlorination products of nitriles, hydrogen cyanide monohydrochloride does not apparently exist (see p. 619).

Hinkel and coworkers²⁵⁹ suggested that in the Gattermann synthesis the hydrogen cyanide and hydrogen chloride react as chloromethyleneformamide ($2\text{HCN} \cdot \text{HCl}$), so that the reaction product is not an aldimine hydrochloride, but an arylmethyleneformamide hydrochloride $\text{RCH}=\text{NCH}=\text{NH} \cdot \text{HCl}$. Later⁷⁰ Hinkel showed that this view of the intermediates in the Gattermann reaction is incorrect, and he put forward²⁶⁰ a new mechanism, according to which the aromatic compound reacts with the sesquichloride $2\text{HCN} \cdot 3\text{HCl}$; however, this also is not supported by the experimental data on the Gattermann synthesis of aldehydes, and contradicts more recent views on the structure of hydrogen cyanide hydrochlorides⁷³.

Hydrogen cyanide dihydrochloride (a compound of the chloroimmonium chloride type) is unknown, or at least is not formed in the Gattermann reaction, and the only remaining possibility is that stable hydrogen cyanide hydrochlorides are not intermediates when the reaction is carried out in the normal manner (by passing hydrogen chloride into a mixture of the aromatic compound, hydrogen cyanide, and ZnCl_2 or AlCl_3 as catalyst). The reaction apparently involves the formation of unstable complexes between hydrogen chloride, hydrogen cyanide, and the catalyst, with a reduced electron density at the cyano-carbon atom. Analogous mechanisms for reactions of nitriles are given above. The Gattermann synthesis has been reviewed^{261,262}.

J. The Hoesch Reaction

The first studies describing the synthesis of ketones by the Hoesch reaction were developments of the Gattermann reaction. The Hoesch ketone synthesis²⁶³ is based on the reaction, in a solvent, between a nitrile, hydrogen chloride, and a phenol (usually a polyhydric phenol) to form a ketimine hydrochloride, followed by hydrolysis:



The suggestion²⁶⁴ that the primary reaction products are imidate hydrochlorides which then rearrange to ketimine hydrochlorides has not been confirmed²⁶⁵. Most workers consider that under the conditions of the Hoesch synthesis the nitrile and hydrogen chloride react to form the imidoyl chloride, which then condenses with the phenol^{264,266,267}. Zil'berman and Rybakova have put forward²⁶⁸ new mechanisms for the formation of ketimine hydrochlorides in the Hoesch synthesis, similar to those described above for the reaction of nitriles with hydrogen chloride and hydroxy-compounds.


The utility of the Hoesch synthesis is considerably increased when it is carried out in accordance with these mechanisms²⁶⁸. Unlike the usual procedures²⁶⁶ for preparing ketones by the Hoesch reaction, the reaction between the nitrile and hydrogen chloride and the subsequent electrophilic attack on the phenol are carried out separately. In many cases it has also been found preferable to carry out the reaction not at room temperature, but at 0° (decreasing the temperature increases the stability of the saltlike nitrile hydrohalides¹⁴). The use of the new procedure has shown that, contrary to published opinion²⁶⁶, ketimine hydrochlorides can be obtained from aromatic nitriles in yields as high as those obtained with aliphatic nitriles²⁶⁸. Diketones have been obtained in high yields from resorcinol and various dinitriles (adiponitrile, pimelonitrile, sebaco-nitrile, *p*-xylylene dicyanide, etc.)²⁶⁹, although it had previously been considered²⁶⁶ that the Hoesch synthesis is of little value for the production of diketones.

Of the more interesting studies of the Hoesch synthesis appearing after Spoerri and Du Bois' review²⁶⁶, mention may be made of the condensation of resorcinol or its derivatives with trichloroacetonitrile and hydrolysis of the reaction products to the corresponding benzoic acid derivatives²⁷⁰.

Under the conditions of the Hoesch synthesis, cyano-pyridines react with pyrroles and indoles to yield the corresponding ketimines and hence the corresponding ketones²⁷¹.

Dihydroisoquinolines have been obtained by condensation of the chlorination products of various amides with isosafrole and methylisoeugenol; the suggested mechanism for this reaction is close to that of the Hoesch synthesis²⁷².

The most frequently used catalyst in the Hoesch synthesis is zinc chloride²⁶⁶; in individual cases ferric chloride²⁷³ or aluminium chloride^{274,275} is a more suitable catalyst. The catalytic action of zinc chloride has recently been investigated in more detail³⁸. It might have been expected that the catalytic action of zinc chloride on the Hoesch synthesis is manifested through the complexes $\text{RCN}\cdot\text{ZnCl}_2$ and $(\text{RCN})_2\cdot\text{ZnCl}_2$ or the nitrilium salt $[\text{RCNH}]^+\text{ZnCl}_3^-$, but stable complexes of such structures were tested and found not to react with resorcinol in many cases, particularly at low temperature. It has been suggested that in the presence of zinc chloride the Hoesch synthesis involves the formation of unstable complexes with a covalently-bonded ZnCl_3 group, *e.g.*:




The part played by the zinc chloride in the production of ketimine hydrochlorides from nitriles which under the conditions of the Hoesch synthesis react with hydrogen chloride to form salts of the *N*-alkylideneamide type³⁸, is interesting; it was found that in the reaction between resorcinol, hydrogen chloride, and chloroacetonitrile the catalyst acts primarily to prevent dimerisation of the nitrile.

In the presence of SnCl_4 and TiCl_4 , which readily form stable complexes with nitriles, ketimine hydrochlorides are produced only in very low yields³⁸.

K. Reaction of Dimeric Nitrile Salts ($2\text{RCN}\cdot n\text{HX}$) with Hydroxy-Compounds

1. Reaction with water. The structures of most of the known dimer salts of composition $2\text{RCN}\cdot n\text{HX}$, where *n* is 1 or 3, have been proved by reaction with water:




This method for the production of secondary amides was known^{59,63} even before Gröndmann and coworkers suggested⁶⁰ this formula for the dimer salts. The reaction is usually carried out by dissolving the dimer salt in water at room temperature or at 0° . A modified procedure has been used to prepare hexachlorodiacyetamide without further hydrolysis of the product at the moment of its formation: a suspension of the trichloroacetonitrile dimer salt in chloroform was shaken with an excess of finely ground ice⁶⁰.


Hydrolysis of the dimer salts derived from chloroacetonitrile and hydrogen bromide yields $\alpha\alpha'$ -dichlorodiacyetamide having some of the chlorine atoms replaced by bromine. A similar exchange also took place in the reaction of bromoacetonitrile dimer salt with hydrogen chloride³⁴.

The $\alpha\alpha'$ -dichlorodiacyetamide obtained from chloroacetonitrile, deuterium chloride, and deuterium oxide contained 1.5 atoms of deuterium. The infra-red spectrum of the deuterated dichlorodiacyetamide indicated the presence of N-H , N-D , C-H , and C-D bonds¹¹⁶.

2. Reaction with alcohols (amidine hydrochloride synthesis). Treatment of dimer salts of composition $2RCN \cdot nHX$ derived from fluoro-, chloro-, and bromo-acetonitrile with primary and secondary alcohols at room temperature causes unsymmetrical scission, with the formation of halogenoacetamide hydrochlorides, ethers, and esters^{34,62,276}:



Methanol reacts anomalously; instead of the chloroacetamide salt, it gives a compound of empirical formula $2ClCH_2CN \cdot 2CH_3OH \cdot HCl$, which with water gave chloroacetamide hydrochloride. The primary reaction product probably has the constitution




The reaction of the dimer salts with other alcohols apparently also involves a first step in which a labile chlorine atom is replaced by an alkoxy-group, but the resulting intermediates are unstable and cannot therefore be isolated.

The fact that alcoholysis of the dimer salt produces (in addition to the amidine hydrochloride) a mixture of ester and ether, and not an ortho-ester, merits attention. Certain imidate salts also react with alcohols to yield a mixture of ester and ether instead of an ortho-ester²⁷⁷.

Cyclohexanol reacts with the di(chloroacetonitrile) salt to form a molecular compound of chloroacetamide hydrochloride with cyclohexanol³⁴.

3. Reaction with acetic acid. Treatment of the di(chloroacetonitrile) salt with acetic acid in the cold produces a compound of empirical formula $2ClCH_2CN \cdot CH_3COOH \cdot HCl$, most probably having the structure



With water it yields *sym*-dichlorodiacetamide³⁴.

IV. REDUCTION OF NITRILES IN THE PRESENCE OF HYDROGEN CHLORIDE

According to the data available, the reduction of nitriles in the presence of hydrogen chloride may produce three types of compound: (i) aldimines, which on hydrolysis yield aldehydes; (ii) acyclic trimerisation products, which give bisamides with water; and (iii) amines.

A. Preparation of Aldehydes by the Stephen Method

Nitriles can be reduced in the presence of hydrogen chloride by the Stephen method²⁷⁸ with stannous chloride; stannous bromide also gives good results. Chromous chloride, vanadium trichloride, and titanium trichloride could not be used as reducing agents in this reaction²⁷⁹. The reaction is usually carried out in ether, dioxane, chloroform, ethyl acetate, or other solvent at about room temperature. Aromatic nitriles, as a rule, give good yields of aldimines and hence of aldehydes. Aldehydes cannot always be obtained from aliphatic nitriles²⁸⁰.



The Stephen reaction has been reviewed^{281,282}. After the appearance of these reviews, it was shown that a reaction competing with the reduction of the nitrile to the aldimine is the conversion of the nitrile to the imidate, which takes place under the action of the alcohol formed as a result of scission of the ethyl ether by the hydrogen chloride. At elevated temperatures this process may completely suppress the reduction of the nitrile²⁸³.

The effect of steric hindrance in the reduction of substituted benzonitriles has been investigated²⁸⁴.

The order in which the various reactants enter into the reaction has not been studied. It must be assumed that, as in other reactions of nitriles in the presence of hydrogen halides, outer and inner complexes of nitriles with hydrogen halides and chloroimmonium chlorides are formed in the Stephen synthesis of aldehydes also; Stephen's last study²⁸⁵ supports this assumption.



Although earlier the reaction was usually carried out²⁸¹ by adding the nitrile to the solution of stannous chloride and hydrogen chloride, according to a modified procedure²⁸⁵ a solution of the nitrile in the inert solvent is saturated with hydrogen chloride at 0° , and then a solution of stannous chloride and hydrogen chloride in the same solvent is added. The modified Stephen method gives higher yields of aldehydes, presumably because of preliminary reaction between the nitrile and hydrogen chloride at the low temperature.

The mechanism of the Stephen reaction with the intermediate formation of nitrilium salts²⁸² may be represented as follows:



The aldimine hexachlorostannates very often separate from the reaction mixture as easily crystallising precipitates²⁸¹.

If the intermediates are chloroimmonium chlorides, the probable mechanism is




B. Preparation of Bisamides in the Reduction of Nitriles by the Stephen Method


Stephen showed²⁷⁸ that aliphatic nitriles are reduced to aldimines as readily as aromatic nitriles; however, more recent workers have found²⁸⁶ that the reduction of lauric acid nitrile by stannous chloride followed by hydrolysis produces not the aldehyde²⁸⁷, but the *NN'*-alkylidenebisacylamide RCONHCH(R)NHCOR .

Two mechanisms have been suggested for the formation of bisamides in the reduction of nitriles. The intermediate which is hydrolysed to the bisamide may be

formed²⁸⁶ by the stepwise addition of two nitrile molecules to the aldimine:



According to Meerwein²⁸², there is first formed a product from two nitrile molecules, similar to the dimer salts of α -halogenoacetonitriles (see p. 618), which is then reduced. The reduction product reacts with a third nitrile molecule to form a salt, which is hydrolysed to the bis-amide:



C. Preparation of Amines

In certain extremely rare cases the reduction of nitriles by the Stephen method does not stop at the aldimine stage, but proceeds further to form primary amines²⁸⁸. An analogous effect is observed in the stannous chloride reduction of various heterocyclic *N*-phenylimidoyl chlorides, where the reaction products were not the aldimines expected by Sonn and Müller²⁸⁹, but secondary amines²⁹⁰.

Hydrogen chloride also promotes the catalytic hydrogenation of nitriles²⁹¹⁻²⁹⁵; for example, mandelonitrile is reduced on a palladium catalyst to phenylacetone in the absence of hydrogen chloride and to phenylethylamine in its presence. The influence of hydrogen chloride on the catalytic reduction of nitriles is apparently due to the formation of reactive complexes between the nitriles and HCl.²⁹⁴

- E. N. Zil'berman, *Uspekhi Khim.*, **29**, 709 (1960) [*Russ. Chem. Rev.*, **331** (1960)].
- A. Gautier, *Compt. rend.*, **63**, 920 (1866).
- A. Gautier, *Annalen*, **142**, 289 (1867).
- A. Gautier, *Ann. Chim. Phys.* [4] **17**, 103 (1869).
- L. Henry, *Bull. Soc. chim. France*, [2] **7**, 85 (1867).
- C. Engler, *Z. phys. Chem.*, **3**, 506 (1867).
- C. Engler, *Annalen*, **149**, 297 (1869).
- A. Pinner and F. Klein, *Ber.*, **10**, 1889 (1877).
- A. Michael and J. F. Wing, *Amer. Chem. J.*, **7**, 72 (1885-1886).
- H. Biltz, *Ber.*, **25**, 2533 (1892).
- C. Engler, *Annalen*, **133**, 137 (1865).
- C. Engler, *Annalen*, **142**, 65 (1867).
- A. Hantzsch, *Ber.*, **64**, 667 (1931).
- F. Klages and W. Grill, *Annalen*, **594**, 21 (1955).
- E. Ronwin, *Canad. J. Chem.*, **35**, 1031 (1957).
- P. Karrer, "Organic Chemistry" (Translated into Russian), Goskhimizdat, 1960, p. 236.
- H. Henetzka and P. Kurtz, in Houben-Weyl, "Methoden der organischen Chemie" Vol. VIII, G, Theime Verlag, Stuttgart, 1952, p. 651.
- O. Wallach, *Annalen*, **184**, 1 (1877).
- A. V. Kirsanov, *Zhur. Obshch. Khim.*, **22**, 269 (1952).
- A. V. Kirsanov, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, **646** (1954).
- A. V. Kirsanov, Symposium, "Khimiya i Primenenie Fosfororganicheskikh Soedinenii" (Chemistry and Application of Organophosphorus Compounds), *Izd. Akad. Nauk SSSR*, 1957, p. 99.
- A. W. Titherley and E. Warral, *J. Chem. Soc.*, **95**, 1143 (1909).
- N. V. Sidgwick, "The Organic Chemistry of Nitrogen", Oxford, 1937, p. 153.
- J. Houben and E. Pfankuch, *Ber.*, **59**, 1594 (1926).
- G. J. Janz and S. S. Danyluk, *J. Amer. Chem. Soc.*, **81**, 3850 (1959).
- R. West, *J. Amer. Chem. Soc.*, **79**, 4568 (1957).
- T. C. Waddington, *J. Chem. Soc.*, 1708 (1958).
- D. W. A. Sharp, *J. Chem. Soc.*, 2558 (1958).
- T. C. Waddington and F. Klanberg, *J. Chem. Soc.*, 2329 (1960).
- T. C. Waddington and F. Klanberg, *J. Chem. Soc.*, 2332 (1960).
- M. E. Peach and T. C. Waddington, *J. Chem. Soc.*, 1238 (1961).
- L. E. Hinkel and G. J. Treharne, *J. Chem. Soc.*, 866 (1945).
- M. Kuhn and R. Mecke, *Angew. Chem.*, **72**, 462 (1960).
- A. Ya. Lazaris, E. N. Zil'berman, and O. D. Strizhakov, *Zhur. Obshch. Khim.*, **32**, 900 (1962).
- E. N. Zil'berman and A. Ya. Lazaris, *Zhur. Obshch. Khim.*, **31**, 980 (1961).
- G. J. Janz and S. S. Danyluk, *Chem. Rev.*, **60**, 209 (1960).
- F. Klages, R. Ruhna, and W. Hauser, *Annalen*, **626**, 60 (1959).
- E. N. Zil'berman and N. A. Rybakova, *Zhur. Obshch. Khim.*, **32**, 591 (1962).
- F. E. Murray and W. G. Schneider, *Canad. J. Chem.*, **33**, 795 (1955).
- A. Naumann, *Ber.*, **47**, 247 (1914).
- V. A. Pleskov, *Zhur. Fiz. Khim.*, **22**, 351 (1948).
- G. J. Janz and S. S. Danyluk, *J. Amer. Chem. Soc.*, **81**, 3846 (1959).
- G. J. Janz and S. S. Danyluk, *J. Amer. Chem. Soc.*, **81**, 3854 (1959).
- E. N. Zil'berman, T. S. Ivcher, and E. M. Perepletchikova, *Zhur. Obshch. Khim.*, **31**, 2037 (1961).
- A. A. Vlček, *Chem. Listy*, **48**, 1741 (1954).
- J. F. Coetzee, *Diss. Abs.*, **16**, 1071 (1956).
- E. N. Zil'berman and A. E. Kulikova, *Zhur. Obshch. Khim.*, **29**, 1694 (1959).
- C. M. French and I. G. Róe, *Trans. Faraday Soc.*, **43**, 791 (1953).
- A. I. Popov and W. A. Deskin, *J. Amer. Chem. Soc.*, **80**, 2976 (1958).
- R. S. Mulliken, *J. Phys. Chem.*, **56**, 801 (1952).
- R. S. Mulliken and C. Reid, *J. Amer. Chem. Soc.*, **76**, 3869 (1954).
- H. H. Bosshard and H. Zollinger, *Helv. Chim. Acta*, **42**, 1659 (1959).
- G. C. Lander and H. E. Laws, *J. Chem. Soc.*, **85**, 1695 (1904).
- H. Stephen and W. Bleloch, *J. Chem. Soc.*, 886 (1931).
- A. Weddige and M. Körner, *J. prakt. Chem.* [2] **31**, 148 (1885).
- N. Tschervén-Ivanoff, *J. prakt. Chem.* [2], **44**, 142 (1891).
- N. Tschervén-Ivanoff, *J. prakt. Chem.* [2], **46**, 142 (1892).
- J. Tröger, *J. prakt. Chem.* [2], **46**, 353 (1892).
- J. Tröger and O. Lüfning, *J. prakt. Chem.* [2], **69**, 347 (1909).
- Ch. Grundmann, G. Wiesse, and S. Seide, *Annalen*, **577**, 77 (1952).
- A. Colson, *Compt. rend.*, **121**, 825 (1895); *Chem. Zentr.*, **R32** (1897).
- E. N. Zil'berman and A. Ya. Lazaris, *Zhur. Obshch. Khim.*, **31**, 1321 (1961).

63. J. Houben and W. Fischer, *Ber.*, 60, 1759 (1927).
64. E. M. Smolin and L. Rapoport, "sym-Triazines and Derivatives", Interscience, New York, 1959, p. 149.
65. E. R. Bissell and R. E. Spenger, *J. Org. Chem.*, 24, 1147 (1959).
66. L. Claisen and F. Matthews, *Ber.*, 16, 308 (1883).
67. L. Gattermann and K. Schnitzspahn, *Ber.*, 31, 1770 (1898).
68. F. B. Dains, *Ber.*, 35, 2496 (1902).
69. L. E. Hinkel and R. T. Bunn, *J. Chem. Soc.*, 1834 (1930).
70. L. E. Hinkel, E. E. Ayling, and J. H. Beynon, *J. Chem. Soc.*, 184 (1936).
71. L. E. Hinkel and T. I. Watkins, *J. Chem. Soc.*, 407 (1940).
72. L. E. Hinkel and T. I. Watkins, *J. Chem. Soc.*, 647 (1944).
73. L. E. Hinkel and R. P. Hullin, *J. Chem. Soc.*, 1033 (1949).
74. L. E. Hinkel and G. H. R. Summers, *J. Chem. Soc.*, 2813 (1952).
75. J. U. Nef, *Annalen*, 287, 265 (1895).
76. A. Michael and H. Hibbert, *Annalen*, 364, 64 (1909).
77. C. Grundmann and A. Kreutzberger, *J. Amer. Chem. Soc.*, 76, 632 (1954).
78. C. Grundmann and A. Kreutzberger, *J. Amer. Chem. Soc.*, 76, 5646 (1954).
79. L. Goubeau, E. J. Jahn, A. Kreutzberger, and C. Grundmann, *J. Phys. Chem.*, 58, 1078 (1954).
80. C. Grundmann and A. Kreutzberger, *J. Amer. Chem. Soc.*, 77, 44 (1955).
81. A. L. Peiker and C. C. Coffin, *Canad. J. Res.*, 8, 114 (1932).
82. R. Schmitt and L. Glutz, *Ber.*, 1, 66 (1868).
83. F. Tiemann and K. Köhler, *Ber.*, 14, 1976 (1881).
84. F. Tiemann and R. Piest, *Ber.*, 14, 1982 (1881).
85. L. Claisen and O. Manasse, *Ber.*, 20, 2194 (1887).
86. J. E. Bucher, *J. Ind. Eng. Chem.*, 9, 233 (1917).
87. W. Tschelinzeff and W. Schmidt, *Ber.*, 62, 2210 (1929).
88. O. Hromatka, *Ber.*, 75B, 814 (1942).
89. O. W. Bauer and J. W. Teter, U. S. P. 2 443 291; *Chem. Abs.*, 42, 7321 (1948).
90. W. Wenner, U. S. P. 2 489 348; *Chem. Abs.*, 44, 2559 (1950).
91. W. Wenner, *J. Org. Chem.*, 15, 548 (1950).
92. R. C. Thomas, Jr., C. H. Wang, and B. E. Christensen, *J. Amer. Chem. Soc.*, 73, 5914 (1951).
93. R. H. Patton and J. H. Simons, *J. Amer. Chem. Soc.*, 79, 894 (1957).
94. R. P. Welcher, M. E. Castellion, and V. P. Vystrach, *J. Amer. Chem. Soc.*, 81, 2541 (1959).
95. H. A. Michael and J. Jeanpretre, *Ber.*, 25, 1678 (1892).
96. B. S. Rabinowitch and C. A. Winkler, *Canad. J. Res.*, 20B, 221 (1942).
97. F. Mauthner, *Ber.*, 42, 188 (1909).
98. L. Claisen and J. Shadwell, *Ber.*, 11, 620 (1878).
99. C. A. Weisgerber, U. S. P. 2 535 245; *Chem. Zentr.*, 4220 (1952).
100. E. L. Carpenter and H. S. Davis, *J. Appl. Chem.*, 7, 671 (1957).
101. A. H. Homeyer, G. B. De Lamater, and L. A. Patterson, U. S. P. 2 734 917; *Chem. Abs.*, 50, 16 859 (1956).
102. H. S. Anker, *J. Biol. Chem.*, 176, 1333 (1948).
103. R. Rambaud, S. Ducher, A. Broche, M. Brini-Fritz, and M. Vessiere, *Bull. Soc. chim. France*, 877 (1955).
104. V. K. Krieble and R. H. Smellie, U. S. P. 2 425 542; *Chem. Abs.*, 41, 7409 (1947).
105. V. Krieble and R. H. Smellie, U. S. P. 2 441 114; *Chem. Abs.*, 42, 8815 (1948).
106. *Jap. P.* 8273 (1956); *Chem. Abs.*, 52, 11 904 (1958).
107. E. N. Zil'berman and A. E. Kulikova, *Russ. P.* 115 895 (1958).
108. E. N. Zil'berman and A. E. Kulikova, *Russ. P.* 117 459 (1958).
109. E. N. Zil'berman, A. E. Kulikova, and N. A. Sazanova, *Khim. Nauka i Prom.*, 4, 135 (1959).
110. A. E. Kulikova, E. N. Zil'berman, and N. A. Sazanova, *Zhur. Obshch. Khim.*, 30, 2180 (1960).
111. E. N. Zil'berman and N. K. Taikova, *Zhur. Obshch. Khim.*, 30, 2506 (1960).
112. E. N. Zil'berman, S. B. Meiman and A. E. Kulikova, *Zhur. Prikl. Khim.*, 33, 2375 (1960).
113. A. Pinner, *Ber.*, 23, 2917 (1890).
114. M. S. Newman, "Steric Effects in Organic Chemistry" (Translated into Russian), *Inostr. Lit.*, 1960, p. 242.
115. E. E. Royals, "Advanced Organic Chemistry", Constable and Co, London, 1955, p. 571.
116. E. N. Zil'berman, A. Ya. Lazaris, G. G. Petukhov, O. D. Strizhakov, and V. I. Ganina, *Dokl. Akad. Nauk SSSR*, 142, 96 (1962).
117. T. L. Brown, *J. Amer. Chem. Soc.*, 81, 3232 (1959).
118. E. N. Zil'berman, A. Ya. Lazaris, and M. A. Chernyshova, *Zhur. Vses. Khim. Obshch. im. Mendeleeva*, 7, 109 (1962).
119. H. L. Wheeler and B. Barnes, *Amer. Chem. J.*, 24, 76 (1900).
120. F. L. James and W. H. Bryan, *J. Org. Chem.*, 23, 1225 (1958).
121. W. Wislicenus, *Ber.*, 20, 589 (1887).
122. R. Roger and D. G. Neilson, *Chem. Rev.*, 61, 179 (1961).
123. A. Pinner, "Die Imidoaether und ihre Derivate", Oppenheim, Berlin, 1892.
124. E. N. Zil'berman and A. E. Kulikova, *Zhur. Obshch. Khim.*, 29, 3039 (1959).
125. J. Stieglitz and H. McKee, *Ber.*, 33, 807 (1900).
126. A. Knorr, *Ber.*, 49, 1735 (1916).
127. A. Pinner, *Ber.*, 16, 352 (1883).
128. C. L. Stevens, *J. Amer. Chem. Soc.*, 70, 165 (1948).
129. S. M. McElvain and R. D. Mullineaux, *J. Amer. Chem. Soc.*, 74, 1811 (1952).
130. T. Cuvigny, *Bull. Soc. chim. France*, 655 (1957).
131. O. Dimroth and H. Feuchter, *Ber.*, 36, 2238 (1903).
132. R. Forsyth and F. L. Pyman, *J. Chem. Soc.*, 397 (1930).
133. J. N. Ashley, H. J. Barber, A. J. Ewins, S. Newbery, and A. D. H. Self, *J. Chem. Soc.*, 103 (1942).
134. F. Linsker and M. T. Bogert, *J. Amer. Chem. Soc.*, 66, 191 (1944).
135. F. H. Newth and L. F. Wiggins, *J. Chem. Soc.*, 397 (1947).
136. C. A. MacKenzie, G. A. Schmidt, and L. R. Webb, *J. Amer. Chem. Soc.*, 73, 4990 (1951).
137. F. H. S. Curd and C. G. Raison, *J. Chem. Soc.*, 160 (1947).
138. D. J. Morgan, *Chem. and Ind.*, 854 (1959).
139. V. I. Zaretskii, *Russ. P.* 115 896; *Byull. Izobret.*, No. 11, 28 (1958).
140. N. I. Shirokova, T. V. Krasnova, and N. V. Aleksandrov, *Zhur. Prikl. Khim.*, 33, 746 (1960).
141. H. J. Barber, P. Z. Gregory, F. W. Major, R. Slack, and A. M. Woolman, *J. Chem. Soc.*, 84 (1947).
142. G. R. Clemo and E. Walton, *J. Chem. Soc.*, 723 (1928).
143. *Soc. An. des Manufactures des Glaces et Produits Chimiques de Saint-Gobain*, *French P.* 1 120 234 (1958).
144. T. M. Laasko and C. S. Unruh, *Ind. Eng. Chem.*, 50, 1119 (1958).
145. Ch. C. Price and J. Zomlefer, *J. Org. Chem.*, 14, 210 (1949).
146. W. K. T. Gleim, U. S. P. 2 747 980; *Chem. Zentr.*, 11 432 (1957).
147. H. M. Woodburn, A. B. Whitehouse, and B. G. Pautler, *J. Org. Chem.*, 24, 210 (1959).
148. Houben-Weyl, "Methoden der organischen Chemie" (Translated into Russian), Vol. IV, Book 1, Goskhimizdat, 1949, p. 537.
149. E. N. Zil'berman and A. E. Kulikova, *Russ. P.* 119 526 (1958).
150. E. N. Zil'berman and A. E. Kulikova, *Russ. R.* 125 553 (1959).
151. E. N. Zil'berman, A. E. Kulikova, and N. M. Teplyakov, "Mezhdunarodnyi Simpozium po Makromolekulyarnoi Khimii" (International Symposium on Macromolecular Chemistry), Moscow, 1960; Reports and Abstracts, Section I, p. 64.
152. E. N. Zil'berman and A. E. Kulikova, *Zhur. Obshch. Khim.*, 30, 3999 (1960).

153. Jap. P. 1276 (1954); Chem. Abs., 49, 11 694 (1955).
154. Jap. P. 4415 (1954); Ref. Zhur., Khim., 5741 (1957).
155. Mase, Yuki Hosi Kagaku Kekaisi, 14, 709 (1956).
156. E. N. Zil'berman and A. E. Kulikova, Russ. P. 123 154 (1959).
157. N. M. Teplyakov, E. N. Zil'berman, and A. E. Kulikova, Trudy Khim. i Khim. Tekhnol. (Gor'kii), 3, 203 (1960).
158. E. N. Zil'berman, N. M. Teplyakov, and A. E. Kulikova, Russ. P. 125 036 (1959).
159. E. N. Zil'berman and N. M. Teplyakov, Vysokomol. Soed., 1, 934 (1959).
160. E. N. Zil'berman and N. M. Teplyakov, Vysokomol. Soed., 2, 133 (1960).
161. R. H. Hartigan and J. B. Cloke, J. Amer. Chem. Soc., 67, 709 (1945).
162. F. Cramer, K. Pawelzik, and H. J. Baldauf, Chem. Ber., 91, 1049 (1958).
163. E. N. Zil'berman and A. M. Sladkov, Zhur. Obshch. Khim., 31, 245 (1961).
164. C. D. Hurd, "Pyrolysis of Carbon Compounds" (Translated into Russian), GONTI, 1938, p. 192.
165. S. M. McElvain and B. E. Tate, J. Amer. Chem. Soc., 73, 2233 (1951).
166. C. L. Stevens, D. Morrow, and J. Lawson, J. Amer. Chem. Soc., 77, 2341 (1955).
167. M. Chapentier-Morize and H. Felkin, Compt. rend., 242, 1186 (1956).
168. F. Cramer and H. J. Baldauf, Chem. Ber., 92, 370 (1959).
169. O. A. Reutov, "Teoreticheskie Problemy Organicheskoi Khimii" (Theoretical Problems of Organic Chemistry), Izd. Akad. Nauk SSSR, 1956, p. 74.
170. W. Steinkopf, Ber., 42, 617 (1909).
171. W. Steinkopf and W. Malinowski, Ber., 44, 2898 (1911).
172. F. Cramer, K. Pawelzik, and F. W. Lichtenhaler, Chem. Ber., 91, 1555 (1958).
173. A. N. Baksheev and N. I. Gavrilov, Zhur. Obshch. Khim., 22, 2021 (1952).
174. C. S. Marvel, P. de Radtitzky, and J. J. Brader, J. Amer. Chem. Soc., 77, 5997 (1955).
175. H. Zahn and W. Lauer, Makromol. Chem., 23, 85 (1957).
176. A. Colson, Compt. rend., 121, 1155 (1895).
177. A. Colson, Ann. Chim. Phys., 12, 231 (1897).
178. R. Walther and P. G. Schickler, J. prakt. Chem., [2], 55, 305 (1897).
179. E. N. Zil'berman, Zhur. Obshch. Khim., 30, 1277 (1960).
180. A. Ya. Lazaris and E. N. Zil'berman, Trudy Khim. i Khim. Tekhnol. (Gor'kii), 3, 434 (1960).
181. V. K. Kriebler and R. H. Smellie, U. S. P. 2 390 106; Chem. Abs., 40, 1868 (1946).
182. W. Franske and F. Boxmann, German P. 930 445; Chem. Abs., 52, 19 955 (1958).
183. F. Cramer and K. Baer, Chem. Ber., 93, 1231 (1960).
184. F. D. Cramer and G. Weimann, Chem. and Ind., 46 (1960).
185. F. Cramer, Angew. Chem., 72, 236 (1960).
186. F. Cramer and H. J. Baldauf, Angew. Chem., 72, 627 (1960).
187. F. Cramer and G. Weimann, Chem. Ber., 94, 996 (1961).
188. R. J. W. Cremlyn, J. Chem. Soc., 1805 (1961).
189. C. L. Stevens and M. E. Munk, J. Amer. Chem. Soc., 80, 4065 (1958).
190. A. U. Rahman, M. A. Medrano, and O. P. Mittal, Rec. Trav. chim., 79, 188 (1960).
191. S. Ishikawa, Sci. Papers Inst. Phys. Chem. Res., Tokyo, 7, 293 (1927); Chem. Zentr., 1, 1764 (1928).
192. S. Ishikawa, Sci. Papers Inst. Phys. Chem. Res., Tokyo, 7, 277 (1927); Chem. Zentr., 1, 1764 (1928).
193. E. C. Taylor and J. A. Zoltewicz, J. Amer. Chem. Soc., 82, 2656 (1960).
194. S. S. Minovici, Ber., 29, 2097 (1896).
195. F. Henle and G. Schupp, Ber., 38, 1369 (1905).
196. E. N. Zil'berman, Trudy Khim. i Khim. Tekhnol. (Gor'kii), 4, 376 (1961).
197. V. Migrdichian, "The Chemistry of Organic Cyanogen Compounds", 1947, p. 95.
198. A. Strecker, Annalen, 103, 321 (1857).
199. A. W. Titherley and T. H. Holden, J. Chem. Soc., 101, 1871 (1912).
200. J. B. Polya and T. M. Spotswood, Rec. Trav. chim., 67, 927 (1948).
201. J. B. Polya and P. L. Tardrew, J. Chem. Soc., 1081 (1948).
202. D. Davidson, Trans. N. Y. Acad. Sci., 20, 316 (1958).
203. G. Fraenkel and C. Niemann, Proc. Nat. Acad. Sci. U. S. A., 44, 688 (1958).
204. G. Fraenkel and C. Franconi, J. Amer. Chem. Soc., 82, 4478 (1960).
205. J. T. Edwards and S. C. R. Meacock, J. Chem. Soc., 2000 (1957).
206. R. Huisgen and H. Brade, Chem. Ber., 90, 1432 (1957).
207. R. J. L. Martin and J. H. Reece, J. Chem. Soc., 4697 (1960).
208. W. Gerrard, M. F. Lappert, H. Pyszora, and J. W. Wallis, J. Chem. Soc., 2144 (1960).
209. C. D. Schmulbach and R. S. Drago, J. Amer. Chem. Soc., 82, 4484 (1960).
210. M. J. Janssen, Spectrochim. Acta, 17, 475 (1961).
211. G. Fraenkel, A. Loewenstein, and S. Meiboom, J. Phys. Chem., 65, 700 (1961).
212. A. R. Osborn, T. C. W. Mak, and E. Whalley, Canad. J. Chem., 39, 1101 (1961).
213. A. R. Katritzky and R. A. Y. Jones, Chem. and Ind., 722 (1961).
214. A. E. Chichibabin, "Osnovnyi Nachala Organicheskoi Khimii" (Fundamentals of Organic Chemistry), Vol. I, Goskhimizdat, 1953, p. 281.
215. N. Ogata, Bull. Chem. Soc. Japan, 32, 813 (1959).
216. E. Spinner, Spectrochim. Acta, 15, 95 (1959).
217. E. Spinner, J. Phys. Chem., 64, 275 (1960).
218. B. I. Sukhorukov, A. I. Finkel'shtein, E. N. Zil'berman, A. E. Kulikova, and V. I. Ganina, Zhur. Fiz. Khim., 35, 1600 (1961) [Russ. J. Phys. Chem., 785 (1961)].
219. Ch. D. Hurd, M. F. Dull, and K. E. Martin, J. Amer. Chem. Soc., 54, 1974 (1932).
220. Sh. Sugawara and H. Shigehara, J. Pharm. Soc. Japan, 62, 531 (1942); Chem. Abs., 45, 2861 (1951).
221. R. E. Miller and R. Rosenthal, U. S. P. 2 809 208 (1959).
222. R. Jaunin, Helv. Chim. Acta, 35, 1414 (1952).
223. R. Jaunin, M. B. Piccoli, and T. Charalambous, Helv. Chim. Acta, 37, 216 (1954).
224. D. Davidson and M. Karten, J. Amer. Chem. Soc., 78, 1066 (1956).
225. D. Klamann, Monatsh., 84, 925 (1953).
226. N. V. Ridel' and M. P. Gerchuk, Khim. Nauka i Prom., 2, 665 (1957).
227. N. V. Ridel' and M. P. Gerchuk, Zhur. Obshch. Khim., 28, 1306 (1958).
228. E. N. Zil'berman and A. E. Kulikova, Zhur. Vses. Khim. Obshch. im. Mendeleeva, 5, 107 (1960).
229. E. N. Zil'berman and A. E. Kulikova, Russ. P. 127 251 (1959).
230. E. N. Zil'berman, A. E. Kulikova, N. M. Teplyakov, and A. A. Rushinskaya, Zhur. Obshch. Khim., 32, 3039 (1962).
231. E. N. Zil'berman and A. E. Kulikova, Zhur. Prikl. Khim., 35, 869 (1962).
232. A. E. Kulikova and E. N. Zil'berman, Zhur. Obshch. Khim., 30, 596 (1960).
233. A. W. Ralston, H. J. Harwood, and W. O. Pool, J. Amer. Chem. Soc., 59, 986 (1937).
234. D. Davidson and H. Skovronek, J. Amer. Chem. Soc., 80, 376 (1958).
235. E. N. Zil'berman, N. K. Taikova, N. A. Rybakova, and

- Feller, Zhur.Obshch.Khim., 31, 2040 (1961).
236. E. N. Zil'berman and N. A. Rybakova, Zhur. Obshch. Khim., 29, 3347 (1959).
237. B. A. Arbuzov and E. A. Pozhil'tsova, Izv. Akad. Nauk SSSR, Otd. Khim. Nauk, 65 (1946).
238. L. Kh. Freidlin, A. A. Balandin, T. A. Sladkova, D. I. Lainer, and L. G. Emel'yanov, Zhur. Prikl. Khim., 32, 1600 (1959).
239. V. D. Polkovnikov and L. Kh. Freidlin, Izv. Akad. Nauk SSSR, Otd. Khim. Nauk, 1488 (1959).
240. L. Kh. Freidlin and T. A. Sladkova, Izv. Akad. Nauk SSSR, Otd. Khim. Nauk, 151 (1961).
241. R. L. Shriner and F. W. Neumann, Chem. Rev., 35, 351 (1944).
242. H. J. Backer and W. L. Wanmaker, Rec. Trav. chim., 70, 638 (1951).
243. D. R. Husted, U. S. P. 2 676 985 (1960).
244. J. C. Grivas and A. Taurins, Canad. J. Chem., 36, 771 (1958).
245. J. C. Grivas and A. Taurins, Canad. J. Chem., 37, 1260 (1959).
246. J. C. Grivas and A. Taurins, Canad. J. Chem., 39, 761 (1961).
247. H. C. Brown, J. Polymer Sci., 44, 9 (1960).
248. W. Treibs, J. Herrmann, and G. Zimmermann, Chem. Ber., 93, 2198 (1960).
249. J. G. Erickson, U. S. P. 2 615 023; Chem. Abs., 47, 9349 (1953).
250. J. G. Erickson, J. Org. Chem., 20, 1569 (1955).
251. P. Oxley, M. W. Partridge, and W. F. Short, J. Chem. Soc., 303 (1948).
252. J. Cymerman, J. W. Minnis, P. Oxley, and W. F. Short, J. Chem. Soc., 2097 (1949).
253. E. Walton, P. Ofner, and R. H. Thorpe, J. Chem. Soc., 648 (1949).
254. W. Wilson, J. Chem. Soc., 3524 (1952).
255. M. T. Gittos and W. Wilson, J. Chem. Soc., 2371 (1955).
256. W. Wilson, Chem. and Ind., 200 (1955).
257. F. F. Blicke, A. J. Zambito, and R. E. Stenseth, J. Org. Chem. 26, 1826 (1961).
258. L. Gattermann, Ber., 31, 1149 (1898).
259. L. E. Hinkel, E. N. Ayling, and W. H. Morgan, J. Chem. Soc., 2793 (1932).
260. L. E. Hinkel and R. P. Hullin, J. Chem. Soc., 1593 (1949).
261. I. V. Machinskaya, Symposium, "Reaktsii i Metody Issledovaniya Organicheskikh Soedinenii" (Reactions and Methods of Study of Organic Compounds), Vol. 7, Goskhimizdat, 1958, p. 307.
262. W. E. Truce, "Organic Reactions" (Translated into Russian), Vol. 9, Inostr. Lit., 1959, p. 45.
263. K. Hoesch, Ber., 48, 1122 (1915).
264. H. Stephen, J. Chem. Soc., 117, 1529 (1920).
265. J. Houben, Ber., 59, 2878 (1926).
266. P. E. Spierri and A. S. Du Bois, "Organic Reactions" (Translated into Russian), Vol. 5, Inostr. Lit., 1951, p. 284.
267. R. Phadke and R. C. Shah, J. Indian Chem. Soc., 27, 349 (1950).
268. E. N. Zil'berman and N. A. Rybakova, Zhur. Obshch. Khim., 30, 1992 (1960).
269. N. A. Rybakova and E. N. Zil'berman, Zhur. Obshch. Khim., 31, 1272 (1961).
270. S. Ebine, Sci. Reports Saitama Univ., 2A, 69 (1955); Chem. Abs., 50, 1197 (1956).
271. M. Strell and E. Kopp, Chem. Ber., 91, 1621 (1958).
272. M. Lora-Tamayo, G. G. Munoz, and R. Madronero, Bull. Soc. chim. France, 1331 (1958).
273. J. Houben and W. Fischer, J. prakt. Chem., [2], 123, 89 (1929).
274. J. Houben and W. Fischer, J. prakt. Chem., [2], 123, 262 (1929).
275. J. Houben and W. Fischer, J. prakt. Chem., [2], 123, 313 (1929).
276. E. N. Zil'berman and A. Ya. Lazaris, Russ. P. 133 892 (1960).
277. S. M. McElvain and C. L. Stevens, J. Amer. Chem. Soc., 69 2663 (1947).
278. H. Stephen, J. Chem. Soc., 127, 1874 (1925).
279. G. Wittig and H. Hartmann, Ber., 72, 1387 (1939).
280. J. W. Williams, J. Amer. Chem. Soc., 61, 2248 (1939).
281. E. Mosettig, "Organic Reactions" (Translated into Russian), Vol. 8, Inostr. Lit., 1956, p. 316.
282. O. Bayer, in Houben-Weyl, "Methoden der organischen Chemie", Vol. VII, Book I, G. Thieme Verlag, Stuttgart, 1954, p. 299.
283. M. M. Shemyakin, V. I. Maimind, B. V. Tokarev, and V. I. Karpov, Zhur. Obshch. Khim., 28, 978 (1958).
284. Chang Pang, Chiang Kweiche, Hsu Ray-chin, Acta Chim. Sinica, 24, 141 (1958); Ref. Zhur. Khim., 19310 (1959).
285. T. Stephen and H. Stephen, J. Chem. Soc., 4695 (1956).
286. J. A. Knight and H. D. Zook, J. Amer. Chem. Soc., 74, 4560 (1952).
287. E. Lieber, J. Amer. Chem. Soc., 71, 2862 (1949).
288. J. P. Wibaut, J. Overhoff, and P. J. Hubers, Rec. Trav. chim., 52, 55 (1933).
289. A. Sonn and E. Müller, Ber., 52, 1927 (1919).
290. T. S. Work, J. Chem. Soc., 429 (1942).
291. W. H. Hartung, J. Amer. Chem. Soc., 50, 3370 (1928).
292. K. Kindler and W. Peschke, Annalen, 485, 113 (1931).
293. K. Kindler and F. Hesse, Arch. Pharm., 271, 439 (1933).
294. K. Kindler and K. Schrader, Annalen, 564, 49 (1949).
295. V. A. Preobrazhenskii, A. M. Polyakova, and S. R. Rafikov, Zhur. Obshch. Khim., 12, 518 (1942).

CHEMICAL ADVANCES IN SYNTHESIS AND REACTIONS OF DIARYLUREAS

D. F. Kutepov

CONTENTS

I. Introduction	634
II. Synthesis of diarylureas by phosgenation of aromatic amines	634
III. Conditions for and mechanism of the phosgenation of aromatic amines	636
IV. Synthesis of diarylureas and arylalkylureas by the reaction of amines with aryl isocyanates	639
V. Synthesis of dialkylureas and diarylureas by the reaction of amines with urea	641
VI. Conditions for and mechanism of the reaction of diarylureas with aromatic amines	644
VII. Synthesis of substituted diarylureas by nitration, chlorination, and sulphonation of diphenylurea	646
VIII. Hydrolysis and other reactions of diarylureas	648
IX. Synthesis of aromatic amines <i>via</i> diarylureas and by other new methods	650