

One-Step Conversion of Oximes to *Gem*-Chloro-Nitro Derivatives

Paolo Ceccherelli,[†] Massimo Curini,^{*} Francesco Epifano, Maria Carla Marcotullio and Ornelio Rosati


Istituto di Chimica Organica, Facoltà di Farmacia, Università degli Studi, Via del Liceo, 06123 Perugia, Italy

Received 17 February 1998; revised 3 April 1998; accepted 8 April 1998

Abstract: a new method for one-pot conversion of oximes to *gem*-chloro-nitro compounds using Oxone and sodium chloride is described. © 1998 Elsevier Science Ltd. All rights reserved.

Keywords: oximes; oxidation; nitro compounds; diastereoselection.

Gem-chloro-nitro compounds are versatile intermediates in organic synthesis for the preparation of molecules containing nitro groups. The preparation of these compounds is usually achieved, with good results, by chlorination-oxidation of oximes.¹ The oxime is initially transformed by chlorination into the *gem*-chloro-nitroso derivative, which is then oxidized to the desired *gem*-chloro-nitro compound.



The first step of the transformation is achieved by the use of elemental chlorine² aqueous hypochlorous acid,² *t*-butyl hypochlorite.³ The resulting chloro-nitroso compound is further oxidized with nitric,⁴ trifluoroperoxyacetic⁵ or *m*-chloroperbenzoic acids,⁶ ozone,⁷ aqueous sodium² or *n*butylammonium hypochlorite.³ The one-pot conversion can be obtained using N,N,N,-trihalo-1,3,5-triazines⁸ or chloroperoxidase in the presence of H₂O₂ and KCl.⁹

The methods reported have some limitations such as: use of strong and non-selective oxidizing agents, toxic and expensive reagents, low yields and long reaction times.

We wish to report that commercial potassium hydrogen monopersulfate (Oxone[®]) in the presence of NaCl can be used for selective conversion of oximes into *gem*-chloro-nitro derivatives under mild conditions in a one-pot procedure. When oximes are treated with Oxone[®] and NaCl supported on wet basic alumina¹⁰ in chloroform at 45 °C, *gem*-chloro-nitro compounds are produced in good yields (Table).

[†] Deceased on January 1, 1998.

^{*} Corresponding author. Email: curini@unipg.it

Table. Chlorination-oxidation of oximes

Entry	Product	Time (h)	Yield (%) ^a
1	1-chloro-1-nitro-cyclohexane	4	79
2	5-chloro-5-nitro-nonane	2	84
3	1-chloro-1-nitro-4- <i>t</i> -butyl-cyclohexane	4	80
4	1-chloro-1-nitro-cyclooctane	1	83
5	2-chloro-2-nitro-adamantane	1	74
6	1 β -chloro-1-nitro- <i>trans</i> -decaline	2	77
7	(1R,3R,5S)-3-chloro-3-nitro-menthane	7	66
8	3 α -acetoxy-17 β -chloro-17-nitro-androstane	1	73

^a Yields in pure isolated products, characterized by IR, GC-MS, ¹H NMR and ¹³C NMR.

The process is carried out by the species produced as a result of the Oxone[®]-oxidation of the chloride anion.¹¹ It is important to note that only in the case of chlorination-oxidation of a phenyl-methyl-oxime does the reaction occurs with the formation of the parent ketone and 1-chloro-1-phenyl-ethene (8:1), being unprecedented. The reaction is diastereoselective (entries 6, 7, 8) as shown by GC-MS and comparison of NMR data with those reported in the literature,¹² giving the diastereoisomer in which the nitro group occupies an axial position.

The procedure described is equally effective or better than any procedure yet reported and is also effective for the transformation of sterically hindered substrates.

Typical procedure. Oxone[®] was added to a well stirred suspension of NaCl (3 mmol) and wet basic alumina (6 g) in chloroform (15 mL) and the mixture heated at 45 °C for 5 min. A solution of oxime (0.6 mmol) in chloroform (1.5 mL) was the added and the reaction mixture stirred until the deep blue color disappeared (1-7 h, see Table). The mixture was filtered under vacuum and the solution evaporated under reduced pressure. The crude material was purified by chromatography on Al₂O₃ (III) (eluent petroleum ether) affording the desired *gem*-chloro-nitro compound.

Acknowledgement. Financial support from the CNR, Rome and the Ministero della Pubblica Istruzione, Italy, is gratefully acknowledged.

References

- [1] Russell GA, Dedolph DF. *J. Org. Chem.* 1985;50:2498-2502. Lechevallier A, Beugelmans RA, Amarollah-Madjdabadi A. *Synthesis* 1986:826-827. Lechevallier A, Beugelmans RA, Amarollah-Madjdabadi A. *Ibid.* 1986:828-830. Ranganathan S, Raman H, Srinivasan CV. *Tetrahedron* 1978;34:3129-3132. Fishwick BR, Rowles DK, Stirling CJ. *J. Chem. Soc. Perkin Trans. 1* 1986:1171-1179. Wade PA, Dailey WP, Carroll PJ. *J. Am. Chem. Soc.* 1987;109: 5452-5461.
- [2] Archibald TG, Garver LG, Baum K, Cohen MC. *J. Org. Chem.* 1989;54:2869-2873.
- [3] Corey EJ, Estreicher H. *Tetrahedron Lett.* 1980;21:1117-1120.
- [4] Iffland DC, Criner GX, Koral M, Lotspeich FJ, Papanastassiou ZB, White SM Jr. *J. Am. Chem. Soc.* 1953;75:4044-4047.
- [5] Marchand AP, Arney BE, Dave PR Jr. *J. Org. Chem.* 1988;53:443-446. Marchand AP, Suri S C. *J. Org. Chem.* 1984;49:2041-2043.
- [6] Ibne-Rasa KM, Edwards JO, Kost MT, Gallopo AR. *Chem. Ind. (London)* 1974;23:964-966.
- [7] Barnes MW, Patterson JM. *J. Org. Chem.* 1976;41:733-735.
- [8] Walters RT, Zajac WW, Woods JM. *J. Org. Chem.* 1991;56:316-321.
- [9] Zaks A, Yabannavar AY, Dodds DR, Evans AC, Pradip RD, Malchow R. *J. Org. Chem.* 1996;61:8692-8695.
- [10] Greenhalgh RP. *Synlett* 1992:235-236.
- [11] Dieter RK, Nice LE, Velu SE. *Tetrahedron Lett.* 1996;37:2377-2380.
- [12] Kresze G, Mayer NM, Winkler J. *Liebigs Ann. Chem.* 1971;747:171-190.