

An ELEMENTARY
LABORATORY EXPERIMENT
INVOLVING *the* HOFMANN
REARRANGEMENT

*THE PREPARATION OF METHYLAMINE HYDROCHLORIDE FROM ACETAMIDE
BY MEANS OF CALCIUM HYPOCHLORITE*

C. R. HAUSER AND W. B. RENFROW, JR.

Duke University, Durham, North Carolina

THE well-known Hofmann reaction by which an amide, in the presence of chlorine or bromine and an alkali, is converted into a primary amine with one less carbon atom, is of considerable interest in organic chemistry. The reaction may be used in going "down series," and it serves as a laboratory method for the preparation of a primary aliphatic amine. The reaction is of interest also because it involves a molecular rearrangement.

The Hofmann reaction is often illustrated in the laboratory by the preparation of methylamine hydrochloride from acetamide. It has been our experience,

however, that when this experiment is carried out with bromine and alkali according to the directions given in many laboratory manuals, a considerable portion of the product consists of ammonium chloride.

In this paper directions are given for the preparation of methylamine hydrochloride from acetamide using commercial calcium hypochlorite* and sodium hydroxide. The product obtained with these reagents is contaminated with only a small amount of ammonium

* The Mathieson Alkali Works, Niagara Falls, N. Y., manufacture calcium hypochlorite under the trade name of "H.T.H."

alcohol distilled off until crystals begin to form in the hot liquid. The solution is then allowed to cool to room temperature and is finally cooled in an ice-bath. The crystals are filtered with suction, washed with 10 cc. of absolute alcohol, and dried on a watch glass set on a hot steam coil. The yield of methylamine hydrochloride should be about fifty-five per cent. of the theoretical amount, melting at 228–230°.

The use of air in this experiment serves to agitate

the mixture and prevent it from bumping. If desired, a mercury-sealed mechanical stirrer may be used for this purpose and the air omitted. In this case one receiver will be sufficient. A short-stem funnel is fitted to the adapter, the larger end of the funnel dipping about 0.5 cm. below hydrochloric acid contained in a beaker. By using a mechanical stirrer the reaction may be carried out on a larger scale with no essential modification in procedure.