

S0040-4039(96)00524-2

Total Synthesis of (*ent*)-Korupensamine D

Thomas R. Hoye* and Minzhang Chen

Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455


Abstract The first total synthesis of the enantiomer of the natural product, korupensamine D (**16R**), is described. The key steps include a highly efficient preparation of the enantiomerically pure primary amine **4** via the ring opening of aziridine **2** with an arylcuprate reagent and the development of a one-pot selective functionalization of a hindered secondary amine in the presence of phenolic hydroxyl groups (i.e., **8** to **9**).
Copyright © 1996 Elsevier Science Ltd

Because of the promising anti-HIV profile of the michellamines, the preparation and evaluation of various analogs, including unnatural antipodes, is of considerable interest. The potential use of various korupensamines as starting materials for this purpose has prompted our efforts toward the synthesis of korupensamine D¹ (structure **4** in the preceding *Letter*), which differs from korupensamines A-C¹ (**1-3** in the preceding *Letter*) by virtue of the *cis* (rather than *trans*) arrangement of the methyl substituents on the tetrahydroisoquinoline (THIQ) substructure. Korupensamine D also possesses an N-methyl substituent rather than the free NH present in congeners A-C. We describe here the synthesis of **16R**, the enantiomer of the natural antipode of korupensamine D, by a route that is also directly applicable to the natural antipode.

The synthesis starts with a new sequence for the preparation of the enantiomerically pure primary amine **4**, a valuable intermediate first used by Bringmann for the construction of non-racemic THIQs.² The ring opening of *R*-N-tosylaziridine **2** with the Grignard reagent derived from **1** in the presence of CuBr•SMe₂³ resulted in the formation of **3** in 100% yield. Either enantiomer of aziridine **2** is readily available from D- or L-alanine in three steps.⁴ Cleavage of the toluenesulfonamide group in **3** delivered primary amine **4** (79%) as a single enantiomer (Mosher amide analysis).

The known acetamide **5** (99%) underwent Bischler-Napieralski cyclization to afford the cyclic imine **6** (82%).² The *cis*-configured THIQ **7** has previously been prepared by reduction of cyclic imine **6** with sodium borohydride in high diastereoselectivity (ds>95%).^{2b} We reduced **6** with H₂ and 10% Pd/C, which also gave the *cis*-configured compound **7** (93%) as the only observable (¹H NMR) diastereomer. Demethylation of **7** with excess boron tribromide gave the resorcinol amine HBr salt **8** (~99 %).

The next key intermediate we envisioned was an N-methylated, doubly OH-protected THIQ derivative. A one-pot procedure was developed to selectively generate the hindered carbamate in **9**, leaving the phenolic hydroxyl groups intact. Sequential silylation of compound **8** (2.1 equivalents of triethylsilyl chloride and Et₃N in CH₂Cl₂), carbamate formation (2 equivalents of ethyl chloroformate and Et₃N), and removal of the silyl groups [-5 equiv of TBAF (1.0M in THF)] gave compound **9** (87% for the three steps). Benzylation with benzyl bromide and potassium carbonate smoothly afforded compound **10** (72%), which was reduced with LiAlH₄ to give the N-methylated compound **11** in 94% yield.



a) Mg° , 5 mol% $\text{BrCH}_2\text{CH}_2\text{Br}$, THF; 10 mol% $\text{CuBr}\cdot\text{SMe}_2$, 0 °C; **2** (100%); b) Na/NH_3 , -78 °C (79%); c) Ac_2O , Et_3N (99%); d) POCl_3 , $\text{MeC}\equiv\text{N}$, reflux (82%); e) H_2 , Pd/C (93%); f) xs BBR_3 , CH_2Cl_2 , -78 °C to RT (~99%); g) 2.1 equiv TESCl , Et_3N , CH_2Cl_2 ; ClCO_2Et , Et_3N ; TBAF/THF (87%); h) BnBr , K_2CO_3 (72%); i) LiAlH_4 (94%); j) I_2 , Ag_2SO_4 , EtOH (80%); k) **13**, Pd(PPh_3)₄, sat'd NaHCO_3 , PhCH_3 , reflux (73%); l) HCl , $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (92%); m) H_2 , Pd/C, $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (100%).

Regiospecific iodination of **11** with iodine and silver sulfate gave the cis-configured, N-methylated, C(5)-activated iodide **12** in 80% yield. Palladium(0) catalyzed coupling of **12** with the naphthalene boronic acid (**13**⁵ or its boronic anhydride) gave an ~4:5 ratio of atropisomers **14R** and **14S** in 73% yield. Hydrolysis of the MOM ethers in the mixture of **14** gave the naphthols **15R** and **15S** (92%). These naphthols were partially separable by MPLC (hexanes:EtOAc-2:1, with 3% Et_3N). Hydrogenolysis of the benzyl groups in the single atropisomer **15R** quantitatively provided the enantiomer of koropensamine D (**16R**), which had identical ¹HNMR spectral data to those reported for koropensamine D.¹ Hydrogenolysis of a mixture of **15R** and **15S** similarly provided a mixture of the enantiomer of koropensamine D (**16R**) and its diastereomer **16S**. The specific rotation of the synthetic **16R** is opposite in sign to that of natural koropensamine D, thus confirming the assigned absolute configuration.¹ Preparation of structural analogs of the koropensamines and michellamines from precursors like **16** continues.⁶

Acknowledgment. This work and that described in the preceding *Letter* were supported by grant CA-60284 awarded by the DHHS. We appreciate suggestions offered by Professor B. H. Lipshutz, J. A. Suriano, O. P. Priest, and M. K. Renner.

References and Notes

- Hallock, Y. F.; Manfredi, K. P.; Blunt, J. W.; Cardellina, II, J. H.; Schaffer, M.; Gulden, K.-P.; Bringmann, G.; Lee, A. Y.; Clardy, J.; Francois, G.; Boyd, M. R. *J. Org. Chem.* **1994**, *59*, 6349.
- (a) Bringmann, G.; Jansen, J. R.; Rink, H.-P. *Angew. Chem. Int. Ed. Engl.* **1986**, *25*, 913. (b) Bringmann, G.; Weirich, R.; Reuscher, H.; Jansen, J. R.; Kinzinger, L.; Ortman, T. *Liebigs Ann. Chem.* **1993**, 877.
- For a review see: Tanner, D. *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 599.
- Daub, G. W.; Heerding, D. A.; Overman, L. E. *Tetrahedron*, **1988**, *44*, 3919.
- Hoye, T. R.; Chen, M.; Mi, L.; Priest, O. P. *Tetrahedron Lett.* **1994**, *35*, 8747.
- All new compounds described here and in the preceding *Letter* have been characterized by ¹HNMR spectroscopy and combustion and/or high resolution mass spectrometric analysis. ¹³CNMR data are available for most.

(Received in USA 12 March 1996; accepted 14 March 1996)