## **HEAT OF COMBUSTION**

The heat of combustion of a substance at 25°C can be calculated from the enthalpy of formation ( $\Delta_f H^o$ ) data in the table "Standard Thermodynamic Properties of Chemical Substances" in this Section. We can write the general combustion reaction as

$$X + O_2 \rightarrow CO_2(g) + H_2O(l) + other products$$

For a compound containing only carbon, hydrogen, and oxygen, the reaction is simply

$$C_a H_b O_c + \left(a + \frac{1}{4}b - \frac{1}{2}c\right) O_2 \rightarrow a \ CO_2(g) + \frac{1}{2}b \ H_2 O(l)$$

and the standard heat of combustion  $\Delta_c H^o$ , which is defined as the negative of the enthalpy change for the reaction (i.e., the heat released in the combustion process), is given by

$$\Delta_c H^\circ = -a \ \Delta_{\rm f} H^\circ \left( {\rm CO}_2, \ {\rm g} \right) - \frac{1}{2} b \ \Delta_{\rm f} H^\circ \left( {\rm H}_2 {\rm O}, \ {\rm l} \right) + \Delta_{\rm f} H^\circ \left( {\rm C}_a {\rm H}_b {\rm O}_c \right)$$

393.51 
$$a + 142.915$$
  $b + \Delta_f H^{\circ}(C_a H_b O_c)$ 

This equation applies if the reactants start in their standard states  $(25^{\circ}\text{C})$  and one atmosphere pressure) and the products return to the same conditions. The same equation applies to a compound containing another element if that element ends in its standard reference state (e.g., nitrogen, if the product is  $N_2$ ); in general, however, the exact products containing the other elements must be known in order to calculate the heat of combustion.

The following table gives the standard heat of combustion calculated in this manner for a few representative substances.

Molecular formula	Name	$\Delta_{\rm c} H^{\circ}/{\rm kJ~mol^{-1}}$	Molecular formula	Name	$\Delta_{\rm c} H^{\circ}/{\rm kJ~mol^{-1}}$
	Inorganic substances		C <sub>3</sub> H <sub>8</sub> O	1-Propanol (l)	2021.3
			$C_3H_8O_3$	Glycerol (l)	1655.4
C	Carbon (graphite)	393.5	$C_4H_{10}O$	Diethyl ether (l)	2723.9
CO	Carbon monoxide (g)	283.0	$C_5H_{12}O$	1-Pentanol (1)	3330.9
$H_2$	Hydrogen (g)	285.8	C <sub>6</sub> H <sub>6</sub> O	Phenol (s)	3053.5
$H_3N$	Ammonia (g)	382.8			
$H_4N_2$	Hydrazine (g)	667.1		Carbonyl compounds	
$N_2O$	Nitrous oxide (g)	82.1			
			CH <sub>2</sub> O	Formaldehyde (g)	570.7
	Hydrocarbons		C <sub>2</sub> H <sub>2</sub> O	Ketene (g)	1025.4
			C <sub>2</sub> H <sub>4</sub> O	Acetaldehyde (l)	1166.9
$CH_4$	Methane (g)	890.8	C <sub>3</sub> H <sub>6</sub> O	Acetone (l)	1789.9
$C_2H_2$	Acetylene (g)	1301.1	C <sub>3</sub> H <sub>6</sub> O	Propanal (1)	1822.7
$C_2H_4$	Ethylene (g)	1411.2	C <sub>4</sub> H <sub>8</sub> O	2-Butanone (1)	2444.1
$C_2H_6$	Ethane (g)	1560.7			
$C_3H_6$	Propylene (g)	2058.0		Acids and esters	
$C_3H_6$	Cyclopropane (g)	2091.3			
$C_3H_8$	Propane (g)	2219.2	$CH_2O_2$	Formic acid (l)	254.6
$C_4H_6$	1,3-Butadiene (g)	2541.5	$C_2H_4O_2$	Acetic acid (l)	874.2
$C_4H_{10}$	Butane (g)	2877.6	$C_2H_4O_2$	Methyl formate (1)	972.6
$C_5H_{12}$	Pentane (1)	3509.0	$C_3H_6O_2$	Methyl acetate (l)	1592.2
$C_6H_6$	Benzene (1)	3267.6	$C_4H_8O_2$	Ethyl acetate (l)	2238.1
$C_6H_{12}$	Cyclohexane (l)	3919.6	$C_7H_6O_2$	Benzoic acid (s)	3226.9
$C_6H_{14}$	Hexane (1)	4163.2			
$C_7H_8$	Toluene (l)	3910.3		Nitrogen compounds	
$C_7H_{16}$	Heptane (1)	4817.0			
$C_{10}H_{8}$	Naphthalene (s)	5156.3	CHN	Hydrogen cyanide (g)	671.5
			CH <sub>3</sub> NO <sub>2</sub>	Nitromethane (l)	709.2
	Alcohols and ethers		CH <sub>5</sub> N	Methylamine (g)	1085.6
			$C_2H_3N$	Acetonitrile (1)	1247.2
$CH_4O$	Methanol (l)	726.1	C <sub>2</sub> H <sub>5</sub> NO	Acetamide (s)	1184.6
$C_2H_6O$	Ethanol (1)	1366.8	$C_3H_9N$	Trimethylamine (g)	2443.1
$C_2H_6O$	Dimethyl ether (g)	1460.4	$C_5H_5N$	Pyridine (1)	2782.3
$C_2H_6O_2$	Ethylene glycol (l)	1189.2	C <sub>6</sub> H <sub>7</sub> N	Aniline (l)	3392.8