


Me<sub>3</sub>SiCl-NaI-CH<sub>3</sub>CN AS AN EFFICIENT AND PRACTICAL REDUCING AGENT  
FOR BENZYLIC ALCOHOLS

Takashi Sakai\*, Kazuyoshi Miyata, Masanori Utaka, and Akira Takeda  
Department of Synthetic Chemistry, School of Engineering,  
Okayama University, Tsushima, Okayama 700, Japan

Abstract: Secondary and tertiary benzylic alcohols are reduced conveniently to the corresponding aryl alkanes by using Me<sub>3</sub>SiCl-NaI-CH<sub>3</sub>CN reagent in hexane. The reaction was successfully applied to a short step synthesis of (±)-ar-turmerone.

Me<sub>3</sub>SiCl-NaI-CH<sub>3</sub>CN reagent<sup>1</sup> has been recognized as a Me<sub>3</sub>SiI equivalent and widely used as a versatile reagent in organic synthesis.<sup>2</sup> We now wish to report a new entry of its usefulness as an efficient reducing agent of secondary and tertiary benzylic alcohols. Although such transformation can be accomplished conventionally by using many other reagents, such as P<sub>2</sub>I<sub>4</sub>,<sup>3a</sup> Me<sub>2</sub>SiI<sub>2</sub>,<sup>3b</sup> NaBH<sub>4</sub>/CF<sub>3</sub>CO<sub>2</sub>H,<sup>3c</sup> Me<sub>3</sub>SiH-BF<sub>3</sub>,<sup>3d</sup> Ph<sub>3</sub>P-I<sub>2</sub>,<sup>3e</sup> Li/liq. NH<sub>3</sub>,<sup>3f</sup> and so on,<sup>3g</sup> the present method has advantages that the reaction can be done conveniently by using the inexpensive and readily available reagent under mild conditions. Clean products can be obtained without special purification process. The reagent does not damage the other functional group in the molecule.



A typical procedure for the present reduction is as follows: To a mixture of Me<sub>3</sub>SiCl (1.54 ml, 12 mmol), NaI (1.8 g, 12 mmol), and acetonitrile (0.6 ml, 12 mmol), was added a solution of 1-phenylethanol (1b) (244 mg, 2 mmol) in hexane (2 ml). The mixture was stirred for 24 h at room temperature. Dilution with water, extraction with ether and subsequent isolation process gave ethylbenzene (2b) (158 mg) with sufficient purity in 75% yield.


The reaction was applicable to various types of benzylic alcohols as listed in Table 1. Although further investigation is required for the elucidation of the mechanism, the presence of the carbonium ion intermediate stabilized by the aromatic moiety is supported by the fact that optically active 1l<sup>4</sup> (70% e.e.) was converted into racemic 2l. Formation of benzyl iodide in the case of benzylalcohol (1a) may be explained by the difficulty of formation of the primary carbonium ion intermediate.

Table 1. Reduction of Benzylic Alcohols with  $\text{Me}_3\text{SiCl-NaI-CH}_3\text{CN}$  Reagent

	Alcohol <u>1</u>			Product <u>2</u> <sup>a</sup>		Alcohol <u>1</u>			Product <u>2</u> <sup>a</sup>
	Ar	R <sup>1</sup>	R <sup>2</sup>	Yield (%) <sup>b</sup>		Ar	R <sup>1</sup>	R <sup>2</sup>	Yield (%) <sup>b</sup>
<u>1a</u>	Ph	H	H	c	<u>1g</u> <sup>d</sup>	Pipe- ronyl-	H	Et	57 <sup>d</sup>
<u>1b</u>	Ph	H	Me	75	<u>1h</u>	Ph	Me	Et	76
<u>1c</u>	Ph	H	Et	99	<u>1i</u>	Ph	Me	Pr	82
<u>1d</u>	Ph	H	Pr	75	<u>1j</u>	p-Tolyl	Me	CH <sub>2</sub> CO <sub>2</sub> Et	91
<u>1e</u>	Ph	H	Bu	74	<u>1k</u>	p-Tolyl	Me	CH <sub>2</sub> CO <sub>2</sub> H	66
<u>1f</u>	Ph	H	-CH <sub>2</sub> CO <sub>2</sub> Et	91	<u>1l</u> <sup>e</sup>	Ph	Me	CH <sub>2</sub> CO <sub>2</sub> Me	60

<sup>a</sup>The known compounds were identified by using IR and <sup>1</sup>H NMR spectra. <sup>b</sup>Yields after distillation. <sup>c</sup>Benzyl iodide was obtained in 94% yield. <sup>d</sup>Reference 7. <sup>e</sup>The optical purity was determined to be 70% e.e. by <sup>1</sup>H NMR with Eu(hfc)<sub>3</sub>.

The combination of the aldol reaction with the present reduction would become a useful synthetic method for the extension of an alkyl chain of aromatic ring. As one of the effective applications of the reaction sequence, (±)-ar-turmerone (8)<sup>5</sup> was conveniently synthesized as shown below. Thus, the aldol reaction of cerium enolate<sup>6</sup> of mesityl oxide (4) (LDA/THF, CeCl<sub>3</sub>) with p-methylacetophenone (5) gave an alcohol (6)<sup>7</sup> in 54% yield. A similar treatment of 6 with  $\text{Me}_3\text{SiCl-NaI-CH}_3\text{CN}$  afforded iodide 7, which without isolation was successfully converted to 8 by the treatment with K<sub>2</sub>CO<sub>3</sub>/THF in 82% yield from 6.



## References

1. a) G. A. Olah, S. C. Narang, B. G. B. Gupta, and R. Malhotra, *J. Org. Chem.*, **44**, 1247 (1979); b) G. A. Olah, A. Husain, B. P. Singh, and K. Mehrotra, *ibid.*, **48**, 3667 (1983).
2. For a review: G. A. Olah and S. C. Narang, *Tetrahedron*, **38**, 2225 (1982).
3. a) H. Suzuki, H. Tani, H. Kubota, N. Sato, J. Tsuji, and A. Osuka, *Chem. Lett.*, **1983**, 247; b) W. Ando and M. Ikeno, *Tetrahedron Lett.*, **1979**, 4941; c) G. W. Gribble, R. M. Leese, and B. E. Evans, *Synthesis.*, **1977**, 172; d) M. G. Adlington, M. Orfanopoulos, and J. L. Fry, *Tetrahedron Lett.*, **1976**, 2955; e) F. Bohlmann, J. Staffeldt, and W. Skuballa, *Chem. Ber.*, **109**, 1586 (1976); f) G. H. Small, A. E. Minnella, and S. S. Hall, *J. Org. Chem.*, **40**, 3151 (1975); g) M. Hudlicky, "Reduction in Organic Chemistry," Ellis Horwood, West Sussex (1984), p79.
4. S. Mitsui, K. Konno, I. Onuma, K. Shimizu, *J. Chem. Soc. Jpn.*, **85**, 437 (1964); *Chem. Abstr.*, **61**, 13167 (1964).
5. Our previous synthesis: T. Sakai, K. Miyata, M. Ishikawa, and A. Takeda, *Tetrahedron Lett.*, **26**, 4727 (1985).
6. T. Imamoto, T. Kusumoto, and M. Yokoyama, *Tetrahedron Lett.*, **24**, 5233 (1983). Alcohol 6 was obtained in a very low yield without CeCl<sub>3</sub>.
7. All new compounds showed satisfactory spectral (IR and <sup>1</sup>H NMR) and analytical data.